

## Tryptophan synthetase levels in hist-1 mutants

M. Carsiotis

A. M. Lacy

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Carsiotis, M., and A.M. Lacy (1962) "Tryptophan synthetase levels in hist-1 mutants," *Fungal Genetics Reports*: Vol. 1, Article 2. <https://doi.org/10.4148/1941-4765.1016>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## Tryptophan synthetase levels in hist-1 mutants

### Abstract

Tryptophan synthetase levels in *hist-1 mutants*

Carsiotis, M. and A. Lacy. Tryptophan synthetase levels in hist-1 mutants.

401 (1952)). The equivalent mutant of S. typhimurium has been found to produce normal levels of tryptophan synthetase.

Mutant C-84 (hist-1) of N. crassa produces elevated levels of tryptophan synthetase (Hogness and Mitchell, J. Gen. Microbiol. 11, 401 (1952)).