

An improved plasmid for transformation of *Neurospora crassa* using the pan-2 gene as a selectable marker

Wilson Low
National University of Singapore

Gregory Jedd
National University of Singapore

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Low, W., and G. Jedd (2008) "An improved plasmid for transformation of *Neurospora crassa* using the pan-2 gene as a selectable marker," *Fungal Genetics Reports*: Vol. 55, Article 11. <https://doi.org/10.4148/1941-4765.1091>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

An improved plasmid for transformation of *Neurospora crassa* using the *pan-2* gene as a selectable marker

Abstract

The pOKE103 plasmid can be used to transform *Neurospora crassa* using the *pan-2* gene (NCU10048.3) as a selective marker. We mapped the *pan-2* fragment to the right arm of LG VI, contig 4, and found that it carries a *pan-2* linked gene, *atg-12* (NCU10049.3). We report the removal of *atg-12* to produce an improved plasmid, pOKE104.

An improved plasmid for transformation of *Neurospora crassa* using the *pan-2* gene as a selectable marker

Wilson Low and Gregory Jedd. Temasek Life Sciences Laboratory and Department of Biological Sciences, National University of Singapore, Singapore 117604 gregory@tll.org.sg

Fungal Genetics Reports 55:44-45

The pOKE103 plasmid can be used to transform *Neurospora crassa* using the *pan-2* gene (NCU10048.3) as a selective marker. We mapped the *pan-2* fragment to the right arm of LG VI, contig 4, and found that it carries a *pan-2* linked gene, *atg-12* (NCU10049.3). We report the removal of *atg-12* to produce an improved plasmid, pOKE104.

Genetic transformation is a key tool to study fungi and the ability to create strains transformed with multiple plasmids is especially important for mechanistic cellular and developmental studies. In *Neurospora crassa*, a number of options are available: For reproducible expression, transgenes can be targeted as single copies to the *his-3* locus by vectors designed to repair rarely reverting alleles at *his-3* (Aramayo and Metzberg, 1996; Margolin et al., 1997). Recently, a system has been reported, where integration at the *csr-1* locus produces cyclosporin A resistance and allows the selection of single copy integrants at *csr-1* (Bardiya and Shiu, 2007). Selection of transformants using Hygromycin B resistance is also common (Staben et al., 1989), however, the use of this marker for systematic deletion of *Neurospora* genes (Colot et al., 2006) is likely to preclude its use in many cases.

The *pan-2* marker provides an additional means of transformation. In this case transformation is ectopic and results in the integration of single or multiple copies of the transgene. The pOKE103 vector constructed by J. Grotelueschen and Bob Metzberg contains a genomic fragment of *pan-2* in the pGEM3Z (+) vector backbone and this vector produces tight selection in *pan-2* mutant backgrounds (Wu and Glass, 2001). We examined the *pan-2* fragment and found that it carries an additional gene, *atg-12*, a conserved ubiquitin-related protein associated with cellular autophagy (Hanada and Ohsumi, 2005). Thus, transformation with pOKE103 may also result in variable overproduction of ATG-12. There is currently no evidence to suggest that transformation with pOKE103 produces any phenotypic transformation. Nevertheless, because the effects of additional copies of *atg-12* may depend on the genotype and may emerge only in certain mutant backgrounds, we have removed *atg-12* from pOKE103 to produce a simplified derived plasmid, pOKE104.

Plasmid pOKE 103 contains a unique *NdeI* restriction site 5' of the *atg-12* gene and we used site-directed mutagenesis with two complementary primers ("new-*NdeI*"; GTC GGA ACA TAT GTT ATG GAC AGC ATG) to insert a second *NdeI* site 3' of *atg-12* (Fig. 1). Digestion of this vector with *NdeI* removes a 1347 bp fragment containing the entire *atg-12* gene. The vector fragment from this digestion was circularized to produce the pOKE104 vector (Fig. 1). The complete sequence of pOKE104 can be found using NCBI accession number FJ599664 and the plasmid can be obtained from the Fungal Genetics Stock Center (<http://www.fgsc.net/>).




Figure 1. Partial restriction maps (drawn with Vector NTI) of plasmids pOKE103 and pOKE104. The introduced *NdeI* site is shown on the left in pOKE103 (arrow); the right map shows pOKE 104 after removal of *atg-12*. The polylinkers are indicated with unique restriction sites shown in red bold.

Acknowledgements: We thank J. Grotelueschen and Bob Metzberg for constructing the pOKE103 plasmid and the Temasek Life Sciences Laboratory for funding.

References

- Aramayo, R., and R.L. Metzberg. 1996. Gene replacements at the *his-3* locus of *Neurospora crassa*. *Fungal Genet Newslet.* 43:9 -13.
- Bardiya, N., and P.K. Shiu. 2007. Cyclosporin A-resistance based gene placement system for *Neurospora crassa*. *Fungal Genet Biol.* 44:307-14.
- Colot, H.V., G. Park, G.E. Turner, C. Ringelberg, C.M. Crew, L. Litvinkova, R.L. Weiss, K.A. Borkovich, and J.C. Dunlap. 2006. A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors. *Proc Natl Acad Sci U S A.* 103:10352-7.
- Hanada, T., and Y. Ohsumi. 2005. Structure-function relationship of Atg12, a ubiquitin-like modifier essential for autophagy. *Autophagy.* 1:110-8.
- Margolin, B.S., M. Freitag, and E.U. Selker. 1997. Improved plasmids for gene targeting at the *his-3* locus of *Neurospora crassa* by electroporation. *Fungal Genet Newslet.* 44:34-36.
- Staben, C., B. Jensen, M. Singer, J. Pollock, M. Schechtman, J. Kinsey, and E. Selker. 1989. Use of a bacterial Hygromycin B resistance gene as a dominant selectable marker in *Neurospora crassa* transformation. *Fungal Genet Newslet.* 36:79-81.
- Wu, J., and N.L. Glass. 2001. Identification of specificity determinants and generation of alleles with novel specificity at the *het-c* heterokaryon incompatibility locus of *Neurospora crassa*. *Mol Cell Biol.* 21:1045-57.