

## Bird Medium: an alternative to Vogel Medium

Robert L. Metzberg  
*University of California*

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Metzberg, R. L. (2004) "Bird Medium: an alternative to Vogel Medium," *Fungal Genetics Reports*: Vol. 51, Article 8. <https://doi.org/10.4148/1941-4765.1138>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## Bird Medium: an alternative to Vogel Medium

### Abstract

This medium was designed to circumvent some problems that arise in the use of Medium N (Vogel 1964 *Am. Naturalist* 98:435-446). These are, among others, the presence of high levels of citrate, a chelator which leaves the concentration of calcium and trace elements uncertain; the use of ammonium nitrate, which leaves the actual source of nitrogen ambiguous; the use of MgSO<sub>4</sub>, which does not allow the experimenter to vary the concentration of magnesium and sulfur independently; the high activity coefficient for the pKa values of citrate, which makes the pH unnecessarily sensitive to ionic strength; the use of sucrose, which leaves uncertain the nature and relative amounts of the hexose(s) being used at any particular moment; the need to use chloroform as a preservative, which results in the gradual depletion of the aqueous phase of complexes of trace elements.

**Bird Medium: an alternative to Vogel Medium**

Robert L. Metzenberg, Department of Chemistry and Biochemistry, University of California  
Los Angeles CA 90095

Fungal Genetics Newsletter 51:19-20

This medium was designed to circumvent some problems that arise in the use of Medium N (Vogel 1964 Am. Naturalist 98:435-446). These are, among others, the presence of high levels of citrate, a chelator which leaves the concentration of calcium and trace elements uncertain; the use of ammonium nitrate, which leaves the actual source of nitrogen ambiguous; the use of  $MgSO_4$ , which does not allow the experimenter to vary the concentration of magnesium and sulfur independently; the high activity coefficient for the  $pK_a$  values of citrate, which makes the pH unnecessarily sensitive to ionic strength; the use of sucrose, which leaves uncertain the nature and relative amounts of the hexose(s) being used at any particular moment; the need to use chloroform as a preservative, which results in the gradual depletion of the aqueous phase of complexes of trace elements. Molybdate ion is excluded from the trace elements used for Solution 1 because, in concentrated stock solutions, it forms water-insoluble complexes with phosphate plus ammonium ion; instead, it is included in Solution 2, where its presence is benign. Finally, concentrations are expressed in moles rather than in grams, which eases the experimenter's task of thinking in terms of stoichiometry and biomass yield. Bird Medium is not meant to supplant Vogel Medium for routine auxanography, stock-keeping, searches for mutants, or growth of *Neurospora* for preparing DNA, mitochondria, etc. However, it should be seriously considered for critical applications such as preparation of samples for microarrays and for analysis of subtle phenotypes of new mutants.

Bird Medium supports rapid germination of conidiospores and rapid growth of mycelium in good yield, with apparently normal morphology. It appears at least equal to, or better than, Vogel Medium in this regard. It should be noted, however, that mycelial pads harvested from Bird Medium have a subtly different texture from those grown on Vogel Medium, being somehow more slippery to the touch. It is not evident that more slippery is less "normal" for *Neurospora*, nor more so, than less slippery.

**Solution 1**INGREDIENT                      CONCENTRATION AT 1X

	<u>g/liter</u>	<u>mM</u>
water	45 ml	
MES(Sigma M-8250)	4.85 g	22.75
$K_2HPO_4$	1.74 g	10
$NH_4Cl$	1.34 g	25
$K_2SO_4$	0.174 g	1.
NaCl	0.058 g	1.
trace elements without molybdate, 10,000x *, 0.1 ml.		

**Solution 2**INGREDIENT                      CONCENTRATION AT 1X

	<u>g/liter</u>	<u>mM</u>
water	38 ml.	
$MgCl_2 \cdot 6H_2O$	0.203 g	1.
$CaCl_2 \cdot 2H_2O$	0.074 g	0.5
glucose	18 g	100
biotin, 10,000x solution *, 0.1 ml.		
sodium molybdate, 10,000x *, 0.1 ml.		

Make up MES,  $K_2HPO_4$ ,  $NH_4Cl$ ,  $K_2SO_4$ ,  $NaCl$ , and trace elements solution without molybdate in 45 ml. of warm water, which will produce a volume of 50 ml. ("**Solution 1; 20X final strength**"). There is no utilizable carbon source in this solution. It should be stored at room temperature, without chloroform.

Make up  $MgCl_2 \cdot 6H_2O$ ,  $CaCl_2 \cdot 2H_2O$ , biotin, molybdate solution, and glucose in 38 ml. of warm water, which will produce a volume of 50 ml. ("**Solution 2; 20X final strength**"). Store at room temperature over a few ml. of chloroform.

Obviously, it will usually be convenient to make up these two solutions on at least ten times the above scale.

Solutions 1 and 2 can be autoclaved separately in their concentrated form, if desired, and diluted into sterile water, or each can be diluted to 2X, autoclaved, and then combined. The pH of the diluted medium, about 5.8, equals that of Vogel medium, and should not be adjusted. Solution 1 can instead be made up to 20 ml. of 50X instead of 50 ml. of 20X by dissolving the ingredients in 15 ml. of warm water rather than in 45 ml. In practice, however, there are likely to be fewer calculational errors in medium-making if both stock solutions are kept as 20X stock solutions for dilution to 2X, rather than as 50X and 20X. The minor savings in space for Solution 1 are probably not be worth the increased risk of error.

**\* BIOTIN, 10,000X**

	<u>mg/100 ml</u>	<u>mM</u>
water	100 ml	
biotin	6.1	0.25

**\* TRACE ELEMENTS WITHOUT MOLYBDATE, 10,000X**

INGREDIENT                      CONCENTRATION AT 10,000X

	<u>mg/100 ml</u>	<u>mM</u>
water	95 ml	
citric acid·H <sub>2</sub> O	4200	20
ZnSO <sub>4</sub> ·7H <sub>2</sub> O	5750	20
Fe(NH <sub>4</sub> ) <sub>2</sub> (SO <sub>4</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	980	2.5
CuSO <sub>4</sub> ·5H <sub>2</sub> O	250	1.0
H <sub>3</sub> BO <sub>3</sub>	62	1.0
MnSO <sub>4</sub> ·H <sub>2</sub> O	33.3	0.2

**\* SODIUM MOLYBDATE, 10,000X**

	<u>mg/100 ml</u>	<u>mM</u>
water	100 ml	
Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O	48.3	0.2