

# Kansas Agricultural Experiment Station Research Reports

---

Volume 2

Issue 7 Southwest Research-Extension Center Reports

Article 11

---

January 2016

## Tillage Intensity in a Long-Term Wheat- Sorghum-Fallow Rotation

A. Schlegel

Kansas State University, [schlegel@ksu.edu](mailto:schlegel@ksu.edu)

Follow this and additional works at: <http://newprairiepress.org/kaesrr>

 Part of the [Agronomy and Crop Sciences Commons](#)

---

### Recommended Citation

Schlegel, A. (2016) "Tillage Intensity in a Long-Term Wheat- Sorghum-Fallow Rotation," *Kansas Agricultural Experiment Station Research Reports*: Vol. 2: Iss. 7. <https://doi.org/10.4148/2378-5977.1256>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright January 2016 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. K-State Research and Extension is an equal opportunity provider and employer.



---

# Tillage Intensity in a Long-Term Wheat- Sorghum-Fallow Rotation

## Abstract

Grain yields of wheat and grain sorghum increased with decreased tillage intensity in a wheat-sorghum-fallow (WSF) rotation. In 2015, available soil water at wheat planting was greater for no-till (NT) and reduced till (RT) than for conventional till (CT). Similarly, for grain sorghum in 2015, available soil water at planting was increased with NT or RT and least with CT. Averaged across the 15-yr study, available soil water at wheat and sorghum planting was similar for RT and NT and about 1 inch greater than CT. Averaged across the past 15 years, NT wheat yields were 5 bu/a greater than RT and 7 bu/a greater than CT. Grain sorghum yields in 2015 were 42 bu/a greater with long-term NT than short-term NT. Averaged across the past 15 years, sorghum yields with long-term NT have been nearly twice as great as short-term NT (64 vs. 35 bu/a).

## Keywords

tillage intensity, wheat-sorghum-fallow rotation, reduced till, conventional till, no till

## Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

## Cover Page Footnote

This research project was partially supported by the USDA-ARS Ogallala Aquifer Program.

## Tillage Intensity in a Long-Term Wheat-Sorghum-Fallow Rotation<sup>1</sup>

*A. Schlegel*

### Summary

Grain yields of wheat and grain sorghum increased with decreased tillage intensity in a wheat-sorghum-fallow (WSF) rotation. In 2015, available soil water at wheat planting was greater for no-till (NT) and reduced till (RT) than for conventional till (CT). Similarly, for grain sorghum in 2015, available soil water at planting was increased with NT or RT and least with CT. Averaged across the 15-yr study, available soil water at wheat and sorghum planting was similar for RT and NT and about 1 inch greater than CT. Averaged across the past 15 years, NT wheat yields were 5 bu/a greater than RT and 7 bu/a greater than CT. Grain sorghum yields in 2015 were 42 bu/a greater with long-term NT than short-term NT. Averaged across the past 15 years, sorghum yields with long-term NT have been nearly twice as great as short-term NT (64 vs. 35 bu/a).

### Procedures

Research on different tillage intensities in a WSF rotation at the Tribune unit of the Southwest Research-Extension Center was initiated in 1991. The three tillage intensities in this study are conventional (CT), reduced (RT), and no-till (NT). The CT system was tilled as needed to control weed growth during the fallow period. On average, this resulted in four to five tillage operations per year, usually with a blade plow or field cultivator. The RT system originally used a combination of herbicides (one to two spray operations) and tillage (two to three tillage operations) to control weed growth during the fallow period; however, in 2001, the RT system was changed to using NT from wheat harvest through sorghum planting (short-term NT) and CT from sorghum harvest through wheat planting. The NT system exclusively used herbicides to control weed growth during the fallow period. All tillage systems used herbicides for in-crop weed control.

### Results and Discussion

#### *Soil Water*

The amount of available water in the soil profile (0 to 8 ft.) at wheat planting varied greatly from year to year (Figure 1). In 2015, available soil water at wheat planting was greater for NT and RT than for CT. Averaged across the 15-yr study, available soil water at wheat planting was similar for RT and NT (about 7 inches) and 1 inch greater than CT.

---

<sup>1</sup> This research project was partially supported by the USDA-ARS Ogallala Aquifer Program.

Similar to wheat, the amount of available water in the soil profile at sorghum planting varied greatly from year to year (Figure 2). In 2015, available soil water at sorghum planting was increased with NT or RT and least with CT. On average, available soil water at sorghum planting was similar for RT and NT and about 1 inch more than CT.

### *Grain Yields*

Since 2001, wheat yields have been depressed in 10 of 15 years, primarily because of lack of precipitation, while winterkill reduced yields in 2015. Reduced tillage and NT increased wheat yields (Table 1). On average, wheat yields were 7 bu/a higher for NT (20 bu/a) than CT (13 bu/a). Wheat yields for RT were 2 bu/a greater than CT even though both systems had tillage prior to wheat. NT yields were significantly less than CT or RT in only 1 of the 15 years.

The yield benefit from RT was greater for grain sorghum than wheat. Grain sorghum yields for RT averaged 13 bu/a more than CT, whereas NT averaged 29 bu/a more than RT (Table 2). For sorghum, both RT and NT used herbicides for weed control during fallow, so the difference in yield could be attributed to short-term compared with long-term NT. In 2015, sorghum yields were 42 bu/a greater with long-term NT than short-term NT. This consistent yield benefit with long-term vs. short-term NT has been observed since the RT system was changed in 2001. Averaged across the past 15 years, sorghum yields with long-term NT have been nearly twice as great as with short-term NT (64 vs. 35 bu/a).

**Table 1. Wheat response to tillage in a wheat-sorghum-fallow rotation, Tribune, Kansas, 2001–2015.**

Year	Tillage			LSD (0.05)	ANOVA ( $P > F$ )		
	Conventional	Reduced	No-till		Tillage	Year	Tillage × year
	----- bu/a -----						
2001	17	40	31	8	0.002		
2002	0	0	0	---	---		
2003	22	15	30	7	0.007		
2004	1	2	4	2	0.001		
2005	32	32	39	12	0.360		
2006	0	2	16	6	0.001		
2007	26	36	51	15	0.017		
2008	21	19	9	14	0.142		
2009	8	10	22	9	0.018		
2010	29	35	50	8	0.002		
2011	22	20	20	7	0.649		
2012	0	1	5	1	0.001		
2013	0	0	0	---	---		
2014	10	11	18	12	0.336		
2015	10	9	9	9	0.966		
Mean	13c	15b	20a	2	0.001	0.001	0.001

**Table 2. Grain sorghum response to tillage in a wheat-sorghum-fallow rotation, Tribune, Kansas, 2001–2015.**

Year	Tillage			LSD (0.05)	ANOVA ( $P > F$ )		
	Conventional	Reduced	No-till		Tillage	Year	Tillage × year
	----- bu/a -----						
2001	6	43	64	7	0.001		
2002	0	0	0	---	---		
2003	7	7	37	8	0.001		
2004	44	67	118	14	0.001		
2005	28	38	61	35	0.130		
2006	4	3	29	10	0.001		
2007	26	43	62	42	0.196		
2008	16	25	40	20	0.071		
2009	19	5	72	31	0.004		
2010	10	26	84	9	0.001		
2011	37	78	113	10	0.001		
2012	0	0	0	---	---		
2013	37	51	78	32	0.053		
2014	38	72	94	28	0.008		
2015	56	60	102	55	0.153		
Mean	22c	35b	64a	5	0.001	0.001	0.001




Figure 1. Available soil water in 8-ft profile at planting of wheat in a WSF rotation as affected by tillage intensity, Tribune, Kansas, 2001–2015. The last set of bars (Mean) is the average across years.




Figure 2. Available soil water in 8-ft profile at planting of grain sorghum in a WSF rotation as affected by tillage intensity, Tribune, Kansas, 2001–2015. The last set of bars (Mean) is the average across years.