

A non-radioactive electrophoretic mobility shift assay for the detection of heat shock element (HSE)-binding activity in *Neurospora crassa*

U Meyer
University of Bremen

L RENSING
University of Bremen

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Meyer, U., and L. RENSING (1998) "A non-radioactive electrophoretic mobility shift assay for the detection of heat shock element (HSE)-binding activity in *Neurospora crassa*," *Fungal Genetics Reports*: Vol. 45, Article 10. <https://doi.org/10.4148/1941-4765.1257>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

A non-radioactive electrophoretic mobility shift assay for the detection of heat shock element (HSE)-binding activity in *Neurospora crassa*

Abstract

A non-radioactive electrophoretic mobility shift assay based on the digoxigenin detection system had been developed and applied to the analysis of the interaction between the heat shock transcription factor and the heat shock element of *N. crassa*.

A non-radioactive electrophoretic mobility shift assay for the detection of heat shock element (HSE)-binding activity in *Neurospora crassa* - U. Meyer and L. Rensing Institute of Cell Biology, Biochemistry and Biotechnology, University of Bremen, NW2, Leobener Str., 28334 Bremen, Germany.

A non-radioactive electrophoretic mobility shift assay based on the digoxigenin detection system had been developed and applied to the analysis of the interaction between the heat shock transcription factor and the heat shock element of *N. crassa*.

The electrophoretic mobility shift assay (EMSA) is often used to examine DNA-binding proteins. At present most DNA-probes are labeled with the [³²P]-radioisotope and therefore highly radioactive. In addition, the lifetimes of these probes are limited due to the self-destroying radiation and the short half-life of ³²P. In this study, we describe a non-radioactive method based on the digoxigenin-detection system of Boehringer (Mannheim, Germany). The labeled DNA-probe can be stored for one year at -20°C without detectable loss of sensitivity. The aim of this study with *N. crassa* was to analyze the interaction between the heat shock transcription factor (HSF) and the heat shock element (HSE). The DNA-probe used is a slightly modified version of the heat shock element (HSE) of *Saccharomyces cerevisiae* present in the *ssa1* promoter (Figure 1a).

The labeling reaction was done with a thermostable Taq-polymerase (Biotherm, Genecraft, Germany) in a thermocycler under the following conditions. 1) 10 x polymerase-buffer 2 μl, 2) stock solution of dATP(2mM), dCTP(1mM), dGTP(1mM) 4 μl, 3) digoxigenin-dUTP(1mM) 1 μl, 4) HSE (0.5 μg/μl) 1 μl, 5) H₂O 11.5 μl, 6) polymerase (5U/μl) 0.5 μl.

70x	[96°C	2 min
		85°C	10sec
		82.5°C	20 sec
		72°C	2 sec
		4°C	

Protocol for the thermocycler : A hotstart is recommended; add 50 μl mineral oil to cover the sample. In this protocol the annealing temperatures are higher than the elongation temperature (72°C). Choosing the annealing temperature lower than 72°C would lead to a DNA-probe, which is not HSE-like. During this labeling-reaction dTTP is substituted by digoxigenized dUTP (Boehringer Mannheim, Germany). It is important to use a polymerase that accepts dUTP as a dTTP analog. Following the label reaction, the mineral oil was discarded, and the probe was precipitated by adding ethanol (80 μl) in the presence of LiCl (2.5 μl of a 4 M solution) and stored overnight at -20°C. The probe was then centrifuged at 18,000 g (4°C) for 30 minutes. The supernatant was discarded and the precipitate resolubilized in 50 μl bidistilled water. Serial dilutions were dot-blotted onto a DNA-binding membrane to check the label-efficiency. The probe should be detectable even at amounts as low as 0.2 fmol. Our probe was usually detectable at even lower amounts (0.02 fmol). The probe was stored in aliquots at -20°C. We used 1 μl (200 fmol HSE) of the above mentioned solution for one EMSA sample preparation.

N. crassa (*bd, mat A* strain, FGSC-stock no.1858) homogenate was prepared according to the modified procedure for the *S. cerevisiae* heat shock transcription factor (Sorger and Pelham 1987 EMBO J. 6:3035-3041). *N. crassa* mycelium was harvested from liquid cultures (9 cm petri dish, 40 h after inoculation with 1.3x10⁶

conidia/25ml), the medium was then removed, and the mycelium quickly frozen in liquid nitrogen. The mycelium was ground in a precooled mortar with sea sand for about 2-3 minutes. The resulting powder was placed into a small glass vessel, and 1 ml of homogenization buffer (200 mM Tris-HCl pH 8.0, 10% glycerol, 10 mM EDTA, 10⁻¹ M DTT) was added.

The mixture was stirred on a magnetic stirrer for 20 minutes at 4°C and the homogenate centrifuged for 30 minutes at 18,000 g. The supernatant was removed and recentrifuged at the same conditions and the resulting supernatant used for the gel shift assay after determining the protein concentration (25 µg of whole cell extract was used to perform one assay). The homogenate was stored at -20°C. 5 x Binding buffer was prepared according to Sorger and Pelham 1987 EMBO J. 6:3035-3041, modified; 1x : 20 mM HEPES-KOH pH 7.9, 60 mM KCl, 12 % glycerol, 2 mM EDTA, 2 mM DTT. The nonspecific competitor consisted of poly d(I-C) 0.5µg/µl or plasmid DNA, e.g. pBR322, the specific competitor of 400 ng/µl unlabeled HSE-probe (Figure 1b).

The solutions were mixed in the described way (Table 1), vortexed and centrifuged briefly. The binding reactions were incubated for 20 min at room temperature and the samples loaded onto a 4% native 1x TAE-polyacrylamide gel. The samples were electrophoresed at 200 Volts for about 1.5 h in a cold room and the gel was electroblotted onto a membrane (a positively charged nylon membrane is strongly recommended, e.g. Amersham Hybond-N+) at 400 mA for 60 min. This is followed by UV-crosslinking and/or baking (80-100°C for 30 min). The digoxigenized probe was detected by an anti-digoxigenin-fab-fragment conjugated with alkaline-phosphatase according to the manual of Boehringer Mannheim. We always used the BCIP/NBT dye system to detect signals. The results showed that the non-radioactive EMSA is suitable to examine HSE-specific band shifts in *N. crassa*. The experiments revealed a gel shift pattern similar to that of *S. cerevisiae*, showing a specific constitutive HSE-binding activity under normal growth conditions (Figure 2: Lane 1). This signal decreased in heat shocked mycelia (41°C for 1 h). Instead, a more retarded specific signal appeared (possibly due to HSF-phosphorylation during heat shock) (Figure 2: Lane 2), which is again similar to *S. cerevisiae*. A heat shock of 43°C totally erases the constitutive signal but leads to an increased signal strength of the above mentioned slower migrating band (data not shown). We assume that these signals represent interactions between the HSE-probe and the HSF of *N. crassa*.

Table 1: Scheme of binding reactions^a.

Lanes	1	2	3	4
dig.-HSE-probe	+	+	+	+
poly d(I-C)	+	+	+	+
specific competitor	-	-	+	+

^a The 5 x binding buffer (3µl) is first pipetted into a reaction tube, followed by distilled water (ad 15 µl), poly d(IC) (1 µl; 500 ng) and the digoxigenized-HSE-probe (1 µl; 200 fmol). The specific competitor (1 µl; 400ng) was used in binding reaction 3 and 4. Finally 25 µg (different volume) of whole cell extract was added.




Figure 1: a) HSE-probe from *S. cerevisiae*. The nucleotide sequence within the bracket is the HSE itself. Instead of C and T, two adenosines (both are underlined) were inserted into the sticky end on the right, to enable the incorporation of digoxigenin-dUTP. b) Unlabeled HSE was used as a specific competitor to distinguish between specific and nonspecific signals.




Figure 2: Non-radioactive electrophoretic mobility shift assay with cell extracts from *N. crassa*. Lanes 1+3 = cell extract from control cells; lanes 2+4 = cell extract from heat shocked cells; fp = free probe; c = constitutive HSE-binding activity in control mycelia; hs = HSE-binding activity in heat shocked mycelia. ns = non-specific signal; HSE specific signals are no longer detectable in lanes 3+4 due to the presence of specific competitor.