

## Convenient scoring of deoxyglucose resistance in *Neurospora*

V. C. Pollard  
*Stanford University*

David D. Perkins  
*Stanford University*

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Pollard, V. C., and D.D. Perkins (1994) "Convenient scoring of deoxyglucose resistance in *Neurospora*," *Fungal Genetics Reports*: Vol. 41, Article 23. <https://doi.org/10.4148/1941-4765.1385>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## Convenient scoring of deoxyglucose resistance in *Neurospora*

### Abstract

Allen et al. (1989 *J. Bacteriol.* 171:53-58 and personal communication) have described scoring *dgr* mutants on media prepared by adding filter-sterilized fructose and deoxyglucose solutions to medium that has already been autoclaved. This is inconvenient if tests are to be made on tubed slants, as we prefer to do.

# Convenient scoring of deoxyglucose resistance in *Neurospora*

*V. C. Pollard and D. D. Perkins, Department of Biological Sciences, Stanford University, Stanford CA 94305-5020*

Allen et al. (1989 *J. Bacteriol.* 171:53-58 and personal communication) have described scoring *dgr* mutants on media prepared by adding filter-sterilized fructose and deoxyglucose solutions to medium that has already been autoclaved. This is inconvenient if tests are to be made on tubed slants, as we prefer to do.

We are interested in *dgr-1* because it is the closest marker to the nucleolus organizer. In scoring *dgr-1*, we have found it unnecessary to filter-sterilize the sugar. Autoclaving in the medium appears to have no adverse effect. 0.3% 2-deoxyglucose, 0.1% fructose, and 1.5% agar are added to Vogel's minimal salts and dispensed into 10 x 75 tubes, which are then plugged, autoclaved, and slanted. Tests on the slants are perfectly clear, with resistant strains fully grown and sensitives clearly negative. Growth is considerably faster on agar medium than on liquid, with tests on agar readable after two days at 34 deg. or three or four days at 25 deg..