

A new culture method for biochemical studies of the circadian clock

J. Perlman

University of California at Santa Cruz

J. F. Feldman

University of California at Santa Cruz

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Perlman, J., and J.F. Feldman (1979) "A new culture method for biochemical studies of the circadian clock," *Fungal Genetics Reports*: Vol. 26, Article 18. <https://doi.org/10.4148/1941-4765.1706>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

A new culture method for biochemical studies of the circadian clock

Abstract

A new culture method for biochemical studies of the circadian clock

A new culture method for biochemical studies of the circadian clock.

The band strain of *N. crassa* exhibits a well-characterized circadian rhythm of conidiation under conditions of constant temperature and darkness (Sargent et al., 1966 Plant Physiol. 41: 1343-1349). Biochemical and physiological studies of the circadian clock have been very difficult because most cultures used in such experiments have been grown on solid medium to observe the bonding pattern. We have now been able to show that homogeneous, non-bonding cultures also exhibit normal circadian rhythmicity. Included in our studies were lawn-inoculated cultures on solid medium, shaking liquid cultures, and standing liquid cultures. Methods and results are presented only for the latter because that system has been characterized the most extensively and potentially may be of the most use.

Cultures were maintained on Horowitz complete slants. All other medium, both liquid and solid, was Vogel's salts containing 1.2% sodium acetate and 0.05% casamino acids [solid medium contained 1.5% agar]. A conidia from 6-B day old slants were suspended in distilled water or (in later experiments) liquid medium, and filtered through glass wool. The concentration of conidia in the filtrate was measured (Klett-Summerson colorimeter, blue filter), and an aliquot immediately added to a large volume of stirring liquid medium to give a final concentration of 2×10^5 spores/ml. Using an automatic pipetter, 25 ml of the stirring suspension were added to each of several dozen 100 x 15 mm plastic disposable Petri dishes. Six growth tubes with solid medium were inoculated at one end with about 50 microliters of undiluted filtrate. All plates and growth tubes were put in constant light at 25°C. After about a day, they were transferred to constant darkness in an environmental growth chamber, also at 25°C. The Petri dish cultures had visible growth by 24 hours, and subsequently formed a mycelial mat which covered the surface. At several different times after the cultures were placed in the dark, six pieces of mycelium from each of three plates were cut with a cork borer (1.1 mm diam.) and transferred to fresh growth tubes. (The size of the transferred piece has been varied greatly with identical results.) The growth fronts of control and experimental growth tubes were marked in red safelight (G. E. BCJ, 60 watt) at the same time each subsequent day. After about 7 days in the dark, the time of occurrence, or phases, of the first conidial bands of the experimental growth tubes were determined by linear regression analysis and compared to the corresponding band of the control tubes.

The phases observed in the growth tubes inoculated with pieces of the standing liquid cultures were very close to those of the controls at all times sampled. (In some experiments small and consistent phase advances were seen in the experimental tubes.) The sampling manipulations therefore do not affect the phase of the clock, and it may be concluded that the liquid cultures have a normal circadian clock whose phase is set, like that of the controls, by the light-to-dark transition. The phases of experimental growth tubes start to differ from that of the controls at approximately the time when the liquid cultures reach stationary phase (about 55-60 hours of age). It seems likely that either the clock of older cultures "runs down," or that older cultures are susceptible to phase resetting when cut and transferred. This appears to be largely independent of the length of time they spend in constant light. Preliminary experiments suggest that a 3-hour period in constant light (possibly even less) suffices to set the phase of these cultures.

Experiments are underway to relate the age of the cultures and nuclear division timer to the functioning of the clock. Other types of experiments, particularly the addition of various agents to determine their effects on the clock, are now possible.

• • • Thimmon Laboratories, University of California at Santa Cruz, Santa Cruz, CA 95064.