

## A method for labeling DNA in vitro using nicked translation

R. Krumlauf  
*Ohio State University*

C. A. Marzluf  
*Ohio State University*

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Krumlauf, R., and C.A. Marzluf (1978) "A method for labeling DNA in vitro using nicked translation," *Fungal Genetics Reports*: Vol. 25, Article 8. <https://doi.org/10.4148/1941-4765.1729>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## A method for labeling DNA in vitro using nicked translation

### Abstract

A method for labeling DNA in vitro using nicked translation

Krumlauf, R. and C.A. Marzluf.

A method for

labeling DNA in vitro using nicked translation.

The method used to label *N. crassa* DNA in vitro for hybridization reactions is a modification of the technique used by Maniatis et al. (1975, P.N.A.S. 72:1184-1188) to label lambda DNA. Using the ability of *E. coli* DNA polymerase I to translate nicks, we artificially nicked DNA with DNase I and used polymerase I and  $^3\text{H}$ -dTTP to label DNA, with this tech-

nique we have labelled *Neurospora* DNA to  $3.3 \times 10^7$  cpm/ $\mu\text{g}$ , which represents 26% incorporation of label into DNA. Higher specific activities may be obtained by using more label and other labelled bases. The size of the labelled DNA is 247 bases. DNA up to 400 bases long may be made this way.

The reaction buffer contains 50 mM Tris-HCl, pH 7.8, 1  $\mu\text{g}$  DNA, 5 mM MgCl<sub>2</sub>, 10 mM  $\beta$ -mercaptoethanol, 50  $\mu\text{g}/\text{ml}$  BSA, 5  $\mu\text{M}$  cold dNTP's, 2.5  $\mu\text{M}$  dTTP containing .750 mCi  $^3\text{H}$ -dTTP. The reaction mixture is incubated at 15 $^\circ$  C for 10 minutes and the reaction is started by adding  $1 \times 10^{-7}$  mg DNase I. The mixture is incubated one minute with the nuclease when 12.5 units of DNA polymerase I is added. Samples are taken over several hours to monitor the % incorporation. The reaction reaches a plateau after about two hours, but incorporation continues up to four hours.

The reaction is stopped by adding 3 ml of .03 M phosphate buffer, .135 M NaCl and .1% SDS, then boiling for 10 minutes. At this point the reaction mixture contains unincorporated  $^3\text{H}$ -dTTP, labelled DNA and foldback DNA. The foldback DNA is a result of the polymerase displacing DNA strands (instead of hydrolyzing them) and then using the displaced strand as a template. The foldback DNA and unincorporated material may both be removed using hydroxyapatite. The boiled reaction mix is poured on a column of HAP at 50 $^\circ$  C. At 50 $^\circ$  C only single stranded and double stranded DNA will bind in .03 M phosphate buffer. Extensive washing with .03 M PB will elute the unincorporated material and small fragments. Since foldback DNA contains short double stranded regions it may be separated from the labelled DNA which is single stranded (100 $^\circ$  C, 5 min) by eluting the later from the column with .14 M PB. The foldback DNA may be eluted from the column using .5 M PB.

The labelled DNA should then be sired on alkaline sucrose gradients. The DNA we have labelled and isolated in this manner contains very little foldback DNA and represents 75% of the DNA labelled. (Supported by Grant GM-23367 from the National Institutes of Health) - - - Department of Biochemistry and Developmental Biology Program, Ohio State Univ., Columbus, Ohio 43210.