

New ocriflovin-resistant mutants and a gene affecting conidiation, which are expressed only in the presence of a morphological mutant

K. S. Hsu

Taiwan Banana Research Institute

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Hsu, K. S. (1973) "New ocriflovin-resistant mutants and a gene affecting conidiation, which are expressed only in the presence of a morphological mutant," *Fungal Genetics Reports*: Vol. 20, Article 45.
<https://doi.org/10.4148/1941-4765.1853>

This Note on Stocks is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

New ocriflovin-resistant mutants and a gene affecting conidiation, which are expressed only in the presence of a morphological mutant

Abstract

New acr mutants and a gene affecting conidiation

Hsu, K. S. New acriflovin-resistant mutants and a gene affecting conidiation, which are expressed only in the presence of a morphological mutant.

The mutant strains listed below were isolated at the Instituto Interamericano de Ciencias Agrícolas de la OEA, Turrialba, Costa Rica, in 1967. Although it was not possible to complete the study, sufficient information was obtained that they may be useful to other investigators, and cultures have been deposited in the Fungal Genetics Stock Center.

All originated in the strain cr^L; cot-1; ylo-1 A (FGSC#191) obtained from D. D. Perkins. The cr; cot; ylo strain was already heterokaryotic for two apparently spontaneous morphological mutations, KH160 and KH161, when on experiment was begun on induction of acriflavine-resistant mutants by gamma rays. Thus a special type of acr mutant could be detected, where the resistance phenotype is manifested only in the presence of the morphological mutant. Neither morphological KH160 or KH161 is by itself resistant. The morphological mutation segregates from acriflavine resistance in all cases except one, designated acr-6 mo (K19 KH160). (Since the two traits have not been separated by recombination, an alternative interpretation would be to consider ocr-6 as a second-step mutant of mo(KH160) to a resistant allele.)

KH27 and KH165 were isolated following gомmo irradiation; the other new mutants received no mutagenic treatment. The study was terminated before it could be determined whether morphological mutants other than KH160 or KH161 would allow expression of resistance, or whether acr-4 or acr-5 were expressed with either KH160 or KH161. All three acr mutants are capable of growth in the presence of 50 µg acriflavine per ml, when in combination with the appropriate morphological mutant.

Table 1. New acriflavine-resistant strains.

Stock	Isolation Number	Linkage Group
<u>cr; cot-1, grey A</u>	<u>cr^L</u> ; C102, KH165	IR; IVR, IVR
<u>cr; cot-1, grey; ylo-1 a</u>	<u>cr^L</u> ; C102, KH165; Y30539y	IR; IVR, IVR; VI
Note 1. <u>grey</u> : grey conidiation in the presence of crisp. A cross of wild type x <u>cr; grey</u> gave 65 ++; 23 <u>cr</u> +; 23 <u>cr grey</u> .		
Note 2. <u>grey</u> appeared to be linked with <u>cot-1</u> . Among <u>cr</u> isolates of cross wild type x <u>cr; cot grey</u> , 1 <u>cr</u> ++, 14 <u>cr cot grey</u> , 0 <u>cr</u> + <u>grey</u> and 1 <u>cr cot</u> + were observed.		
<u>acr-4; mo</u> (KH160) A	KH16; KH160	II; IIIR
Note 1. <u>acr-4</u> requires simultaneous presence of morphological mutant gene KH160 to manifest acriflavine resistance phenotype		
Note 2. <u>acr-4</u> is linked to mating-type, but is not allelic to <u>acr-3</u> . Two sensitive isolates were found among 78 tested from the cross KH16; KH160 x <u>acr-3</u> (KH8); KH160.		
Note 3. A cross with alcoy indicated that KH160 is in IIIR or VI. Since independent segregation between KH160 and <u>ylo-1</u> had been observed in another cross, KH160 appears to be in IIIR.		
<u>acr-5, mo</u> (KH161) A	KH27, KH161	I or II
<u>acr-5, mo</u> (KH161) a	KH27, KH161	I or II
Note 1. <u>acr-5</u> requires linked morphological mutation KH161 for phenotypic manifestation of acriflavine resistance.		
Note 2. A cross with alcoy indicated that KH161 is in I or II. Hence, <u>acr-5</u> is also in I or II.		
<u>acr-6, mo</u> (KH160) A	KH19, KH160	IIIIR
<u>ocr-6, mo</u> (KH160) a	KH19, KH160	IIIIR
Note 1. <u>acr-6</u> is inseparable from KH160 in IIIIR (no recombinants among 360 progeny from a cross with wild type).		