

## Dominance modification

P. J. Russel  
*Cornell University*

A. M. Srb.  
*Cornell University*

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Russel, P. J., and A.M. Srb. (1971) "Dominance modification," *Fungal Genetics Reports*: Vol. 18, Article 5.  
<https://doi.org/10.4148/1941-4765.1884>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## Dominance modification

### Abstract

Dominance modification

in heterozygous crosses (+/pk), show different characteristic frequencies of linear asci. Thus, it seemed of interest to look for modifiers that act on dominance relations in such heterozygotes. A technique of mutagenesis was employed in order to isolate mutants of wild type (74A) resistant to the colonising action of sorbose, a chemical known to phenocopy the *ascus* abnormality when wild type crosser are subjected to its action. The idea was to determine whether or not on "indirect selection method" would provide strains capable of modifying the dominant peak phenotype at the *ascus* level, in heterozygous crosses.

By this method, 209 pk-boss-resistant mutants were isolated, and 56 of these were crossed to the most dominant peak allele (17-088), which give 96.5% abnormal asci when crossed to 74A. Modifiers were then identified or occurring in those strains that gave a repeatable, statistically significant increase in the percentage of linear asci over that observed in control crosses. Four genic modifiers (pk-mod-1 - pk-mod-4) have been identified in this way. The results of crossing 3 of the modifiers, and 74A, to 17-088 and to the other 4 dominant peaks are shown in Table 1.

Table 1.

Strain	Dominant Peaks									
	17-088		19-722		21-804		22-335		22-384	
	L/T	%L	LIT	%L	L/T	%L	L/T	%L	L/T	%L
+	9712650	3.66	254/1471	17.27	78411013	77.39	2521950	26.53	145/497	29.18
ok-mod-	1290/11,521	11.20*	124/588	21.1	593/642	92.5*	355/605	58.7*	2661442	60.2*
ok-mod-2	502/4106	12.23*	48/264	18.18	3391341	99.41*	216/451	47.89*	1631239	68.20*
pk-mod-3	506/5186	9.76*	135/381	35.43*	2251270	83.33	55/472	11.65	1391239	58.16*

\* Indicates a significant increase over the control value L/T = number linear asci/total number asci scored  
%L = percentage linear asci

It is clear from these results that modifiers first identified in reference to one of the dominant peak alleles do not affect the dominance relations of the other dominant peak alleles in the same way. A difference in specificity of the modifying effect is seen between pk-mod-1 and pk-mod-4 and between pk-mod-2 and pk-mod-3, whereas pk-mod-1 and pk-mod-2 seem to have the some specificity.

Table 2.

Strain	Dominant Pk (17-088) L/T	%L
<u>pk-mod-1; pk-mod-3</u>	102/822	12.41
<u>pk-mod-3; pk-mod-3</u>	30/822	12.88

were in the homozygous condition. Crosses of the type (pk-mod-1; pk/pk-mod-1; pk<sup>+</sup>) were set up and scored for linear versus non-linear asci. With dominant pk (17-088) the result was 175 linears out of a total of 2731 asci; that is, 6.4% linears, indicating a decrease over the effect of the modifier in heterozygous condition.

Further experiments are under way to test whether or not any of the modifiers are allelic, and to amplify existing results. (This work was supported by grant GM-12953, National Institutes of Health, USPHS). ■ ■ ■ Section of Genetics, Development and physiology, Cornell University, Ithaca, New York 14850.

In order to investigate whether the modifiers might act in an additive manner or not, double modifier strains were set up and these were tested against 17-088, as shown in Table 2. A comparison of the figures for the double modifiers with those of the modifiers crossed singly with 17-088 (see Table 1) indicates that there is no significant difference in the modification.

It was then considered that a modifier might have an effect if it