Effect of malachite green in presence of surface actants

D. Mücke

M. Popp

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Mücke, D., and M. Popp (1969) "Effect of malachite green in presence of surface actants," *Fungal Genetics Reports*: Vol. 15, Article 20. <https://doi.org/10.4148/1941-4765.1923>

This Response of Neurospora to Various Antibiotics and other Toxic Chemicals is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Effect of malachite green in presence of surface actants

Abstract

Effect of malachite green in presence of surface actants

Mücke, D. and M. Popp. Effect of malachite green on growth in presence of surface actants.

The LD50 of malachite green was determined to be 3×10^{-5} g per 100 ml, using N. crassa 3a6A, cultured for 5 days at 27°C. At 18°C the same concentration of malachite green inhibits the mycelial growth more than 60% (control without malachite green = 100%) and at 30°C more than 80%. The inhibitory effect of malachite green was increased in the presence of the following surface actants (Tween 40 (10^{-4} g/100 ml), Tween 80 (10^{-4} g/100 ml), Lauryl pyridinium chloride (10^{-5} g/100 ml), and Dimethyl-benzylamino-acetic-dodecylamid (10^{-5} g/100 ml)) at 18°C and at 36°C (except for LK at 36°C). At 27°C the inhibitory effect of the surface octonh tested did not appear. The concentrations of surface actants tested in these investigations did not influence the growth of mycelium if the medium was free of malachite green. - - - Institut für Physiologische Chemic der Universität Rortock, 25 Rortock 1, Leninallee 70, DDR.