Preliminary mapping of nineteen new translocations

J. L. Beske

R. L. Phillips

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Beske, J. L., and R.L. Phillips (1968) "Preliminary mapping of nineteen new translocations," *Fungal Genetics Reports*: Vol. 13, Article 6. <https://doi.org/10.4148/1941-4765.1939>

This Linkage Data is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Preliminary mapping of nineteen new translocations

Abstract

Preliminary mapping of nineteen new translocations

Beske, J. L. and R. L. Phillips. Preliminary mapping of nineteen new translocations with the alcy multiple translocation tester strain.

groups involved. Crosses were made on a 1.7% Difco corn meal agar medium by simultaneously inoculating both parent strains. All crosses were maintained at 25°C. Random ascospores were isolated to a solid Fries minimal or complete medium. For certain crosses, 100 ascospores were isolated on two occasions approximately two weeks apart. The same pattern of re-

The nineteen translocation strains listed in Table 1 were generously given to us by D. D. Perkins and are presently available from the Fungal Genetics Stock Center (see Revised Stock List, *Neurospora Newsl.*, this issue). These strains were crossed with the alcy triple translocation tester strain (T(I;II) al-1; T(IV;V) 2355, cot; T(III;VI) 1, ylo-1) to obtain information on the linkage

Table 1. Linkage data from 19 new translocation strains crossed with the alony multiple translocation tester strain [T(I;II) al-1; T(IV;V) 2355, cot; T(III;VI) 1, ylo-1].

Phenotype	** Genotype	NM 131	NM 121	NM 114	NM 112	NM 111	NM 107	P 2640	NM 163	NM 161	NM 141	NM 170	ALS 6	AR 9	AR 12	AR 1,	NM 109	NM 127	NM 150	NM 180
		A. Independence*						B. <u>al-ylo</u> linkage				C. <u>al-cot</u> linkage	D. Complex results							
<u>cot al</u>	<u>cot al ylo</u> <u>cot al +</u>	3	13	13	13	20	10	18	15	7	16	2,	26	15	42	12	19	31	14	34
<u>cot ylo</u>	<u>cot + ylo</u>	2	7	3	7	7	3	3	0	0	6	2	3	926		1	1		0	0
<u>cot +</u>	<u>cot • +</u>	2	7	6	4	9	0	9	7	8	2	2	6	23	911		2	5	9	0
<u>+ al</u>	<u>+ al ylo</u> <u>+ al +</u>	18	36	19	16	19	10	32	19	26	31	2	33	33	28	7	10	36	22	28
<u>+ ylo</u>	<u>+ + ylo</u>	7	17	8	12	11	9	6	5	2	2	10	10	5	26	10	3	7	6	14
<u>+ +</u>	<u>+ + +</u>	8	13	5	13	15	7	25	20	22	32	32	8	51	17	10	6	20	2	18
Total		40	93	54	65	81	39	93	66	65	89	75	86	136	124	56	41	100	53	94
% Germination		40	47	54	65	81	20	4,	66	65	89	75	43	68	62	56	46	53	53	52
% Recombinants**		Independence						19	15	6	18	8								

* These data were tested for goodness of fit to a ratio of 2 al:1 ylo:1 wild-type in the cot⁺ and cot class. A satisfactory fit was obtained in each case.

** The % recombinants for group B translocations was calculated by doubling the frequency of al⁺ ylo recombinants.

*** cot is now known as cot-1 and ylo is now known as ylo-1.

sults was obtained from the two isolations in every case.

The mechanics of utilizing the *alcoy* strain have been described in detail by Perkins (1964 *Neurospora News*, 6: 22) for mapping new mutants to linkage groups. Perkins (1966 *Neurospora News*, 9: 11) stated that *tns* locations phenotypically indistinguishable from wild type also may be mapped using the *alcoy* tester strain. Normally independent *alcoy* markers will show linkage to each other if the new *tns* location has breaks close to the breakpoints of two of the marked *alcoy* *tns* locations. Therefore, a linkage between *al* and *ylo* would indicate that the new *tns* location involved linkage groups I or II and III or VI. Similarly, a linkage between *al* and *cat* would indicate the involvement of linkage groups I or II and IV or V, while a linkage between *cat* and *ylo* would indicate involvement of IV or V and III or VI. If the *alcoy* markers remain independent, one of the following situations exists:

(1) Linkage group VII is involved in the new *tns* location; (2) The new *tns* location involves linkage groups I and II, III and VI, or IV and V; or (3) One of the two linkage groups involved in the new *tns* location is common to one *alcoy* translocation and the other linkage group is common to another *alcoy* *tns* location, but with the two breaks widely separated in at least one of the common linkage groups. Independence is indicated by a ratio of 2 *al*:1 *ylo*:1 wild type in the *cot*⁺ and *cot* class, since *al* is epistatic to *ylo*.

The linkage results (Table 1) are grouped into four categories; (A) Independence, (B) Linkage of *al* and *ylo*; (C) Linkage of *al* and *cat*; and (D) Complex results not expected of simple reciprocal *tns* locations (note the *al*: non-*al* ratios). The recombination values calculated from the data in categories B and C give a measure of the total genetic length of the two differential (between breaks) segments separating the linked *alcoy* markers and are not extremely valuable in mapping the actual breakpoints of the new translocations.

Fewer *cot* than *cot*⁺ germinants were obtained from crosses involving all but two of the translocation strains (AR17 and NM109). NM150 and NM161 were "morphs" and NM141 and NM170 were "peach", but progeny with these phenotypes are considered as wild types for the purposes of Table 1. An interaction of "peach" with some of the *alcoy* markers is suspected.

The results from NM180 crosses are particularly intriguing since they indicate independence between the *alcoy* markers in the *cot*⁺ class but an *al-cot* linkage in the *cot* class. This unusual genetic behavior might be expected if NM180 were the result of two translocations involving three linkage groups (IV, V, and I or II) with breaks located such that an association of six chromosomes plus a "pair" carrying only the *cot*⁺ allele would result from a cross with the *alcoy* strain instead of an association of eight. This strain will be investigated farther.

In summary, *tns* locations NM107, 111, 112, 114, 121, and 131 are independent of the *alcoy* translocations, NM141, 161, 163, and P2648 involve linkage groups I or II and III or VI, and NM170 involves linkage groups I or II and IV or V. Translocations ALS6, AR9, 12, 17, NM109, 127, 150 and 180 appear to be more complicated than simple reciprocal *tns* locations. (Undergraduate Research Problem by the first author under the direction of the second author conducted as part of Special Problems Course No. 25. ■ ■ ■ Department of Agronomy and Plant Genetics, University of Minnesota, St. Paul, Minnesota 55101.