

Cyanocobalamin in mycelium of *Neurospora sitophila*

H. P. Kleber

H. Aurich

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Kleber, H. P., and H. Aurich (1967) "Cyanocobalamin in mycelium of *Neurospora sitophila*," *Fungal Genetics Reports*: Vol. 12, Article 9. <https://doi.org/10.4148/1941-4765.1954>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Cyanocobalamin in mycelium of *Neurospora sitophila*

Abstract

Cyanocobalamin in *N. sitophila*

Kleber, H.-P. and H. Aurich. Cyanocobalamine
in mycelium of Neurospora sitophila.

fungus was cultured at 30°C in 300 ml Erlenmeyer flasks (30: 784), containing pyridoxine at a concentration of 150 µg/l and glucose as the sole carbon source (20 g/l). Cobalt was not added to the medium additionally. Nevertheless, using the Zeiss ultra-spectrophotometer Q24 we found about 30 µg Co/l medium. This quantity was brought into the culture medium with the trace elements. We followed the growth of the fungus by measuring the weights of mycelial pods after drying at 80°C. For cyanocobalamine determinations, the mycelial pods were washed, dried, homogenized by grinding in a mortar and then extracted with distilled water for 20 min. at 100°C in the presence of 0.05% NaCN. The vitamin B₁₂ concentrations in these extracts were determined by the method described by Muecke and Dummer (1960 Pharmazie 15: 305) using P. stipitata as test organism.

Cyanocobalamine has not yet been described as an essential component of Neurospora. Therefore we studied this compound in mycelium by microbiological assay with Pteriochromonas stipitata. N. sitophila 299 (FGSC#348), a pyridoxine requirer, was used for our experiments. The

for different times on the medium of Ryan et al. (1943 Am. J. Botany 30: 784), containing pyridoxine at a concentration of 150 µg/l. Ammonium tartrate was used as the sole nitrogen source (5 g/l)

Table 1. Growth and cyanocobalamine content of N. sitophila 299.

Age of culture (days)	Mycelial dry wt. (mg/flask)	Cyanocobalamine of mycelium (pg/mg dry wt.)	(pg/flask)
1	2	6	12
2	30	16	480
3	61	31	1891
4	111	35	3885
5	136	16	2176
6	159	12	1908
9	175	7	1225

Vegetative cultures of N. sitophila produce demonstrable amounts of cyanocobalamine, as shown in Table 1. As growth proceeds, the concentration of cyanocobalamine increases. Maximal content was found at the 4th day. At this time only somewhat more than 10% of the exogenous cobalt is incorporated into the cyanocobalamine molecules, calculated from the B₁₂ content. In this case, therefore, the cobalt concentration of the medium is sufficient for cyanocobalamine biosynthesis. At the end of the active growth phase, cyanocobalamine concentration showed a marked decline. — Institute of Physiological Chemistry, Karl Marx University, Leipzig, Germany.