Prototroph frequencies from crosses between pyridoxine auxotrophs

A. Radford

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Radford, A. (1967) "Prototroph frequencies from crosses between pyridoxine auxotrophs," *Fungal Genetics Reports*: Vol. 11, Article 2. <https://doi.org/10.4148/1941-4765.1967>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Prototroph frequencies from crosses between pyridoxine auxotrophs

Abstract

Crosses between pyridoxine auxotrophs

Radford, A. Prototroph frequencies from crosses between pyridoxine auxotrophs.

ascus, being assigned to a separate but closely linked locus, pdx-2. Four other pyridoxine auxotrophs, plus a fifth subsequently lost, were reported but not assigned to a locus (Tatum et al. 1950 Am. J. Botany 37: 38).

Allele	Prototrophs/10 ⁵ spores
37803	1.1, 2.0*
39106	181
39706	110,44*
44204	46, 109*
44502	65. 46*
46904	13
Y2329	1.2
Y 12274	very poor fertility
Y30978	none in 40,000
Y3 1393	none in 12,000

• Results of crosses between alternate strains of these alleles

At present there are eleven pyridoxine auxotrophs of Neurospora crassa (see Stock Lists in NN's #9 and 10). Seven of these have been previously described (Barratt et al. 1954 Adv. Genet. 6:1), six being classified as pdx-1 and the seventh, on the basis of one recombinant

The crossing of pdx-1 allele 35405 to all the other pyridoxine auxotrophs gave the prototroph frequencies indicated in the Table.

It seems clear from these results that the initial assignment of allele 44204 to a second locus was incorrect. This allele, in addition to its behavior in recombination, also shows complementation behavior consistent with its being an integral part of the pdx-1 locus (Radford 1966 Can. J. Genet. Cytol. 8:672). Strains Y2329, Y30978 and Y31393 also show the low prototroph frequencies with 35405 which would be expected from alleles of pdx-1. Strain Y12274 has so far given insufficient spores, when crossed to 35405, for satisfactory analysis. It does show fairly close linkage to me-1 and col-4 on linkage group IV, and is therefore located in the region of pdx-1. - - - Department of Biological Sciences, Stanford University, Stanford, California 94305.