RNA and protein synthesis in *Neurospora crassa* conidia

S. E. Hitchcock

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Hitchcock, S. E. (1967) "RNA and protein synthesis in *Neurospora crassa* conidia," *Fungal Genetics Reports*: Vol. 11, Article 18. <https://doi.org/10.4148/1941-4765.1983>

This Spores is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

RNA and protein synthesis in *Neurospora crassa* conidia

Abstract

RNA and protein synthesis in conidia

In a study of the germination of conidio of wild type strain Em 5297a (ATCC#10816), the synthetic capacities of conidia incubated in minimal medium with and without a carbon source were investigated. Conidia were grown on Vogel's medium N + 1% glucose and 2% agar. Conidia were harvested at 7 days, washed, and incubated in Vogel's liquid medium N with (called "germinating") or without (called "non-germinating") 2 % glucose at a spore concentration of 10 mg FW/ml.

Figure 1 shows RNA synthesis, as measured by umcil-2-C¹⁴ incorporation into hot alcohol-extracted, cold TCA-insoluble material, under these two conditions. There is no difference until 30 minutes, when there is a marked stimulation of incorporation into the conidia incubated in glucose-supplemented medium. This is about the time germ tube formation begins. The difference is nearly fifty-fold after 3 hours and represents a large increase in specific activity as well as in total activity, which is shown here. This stimulation is inhibited by cycloheximide (100 µg/ml).

Protein synthesis, as measured by phenylalanine-U-C¹⁴ incorporation into hot alcohol-extracted, cold TCA-insoluble material, seems to be different from RNA synthesis (Figure 2). While the label incorporated into conidia incubated in supplemented medium takes on immediate lead over those incubated in mineral salts only, the difference is only two-fold after 5 hours. Since the protein in "germinating" conidia has about doubled during this time, there is little, if any, increase in specific activity.

Fractionation of cells into hot alcohol-soluble fractions (containing free amino acids, nucleotides, neutral compounds, and organic acids), RNA, and protein fractions gave interesting results which influence the interpretation of Figures 1 and 2.

Table 1. Incorporation of uracil and phenylalanine.

Glucose (%)	t (hr)	Uracil		t (hr)	Phenylalanine	
		0	2		0	2
Alcohol fractions cpm x 10 ⁻³	1	30.8	48.5	2	888.4	238.2
	2	49.8	313.4	4	877.7	251.8
% of cell activity*	1	77.1	81.9	2	91.7	40.8
	2	67.0	56.4	4	74.2	28.6
RNA (protein)** cpm x 10 ⁻³	1	9.1	24.5	2	70.8	338.9
	2	10.7	214.3	4	292.9	616.2
% of cell activity	1	22.8	32.9	2	7.3	58.1
	2	18.0	43.5	4	24.8	70.1

Flasks contained 1 µc uracil-2-C¹⁴, 2.100 x 10⁶ cpm, final conc. 0.261 µM; 1 µc L-phenylalanine-U-C¹⁴, 2.160 x 10⁶ cpm, final conc. 0.77 µM (10 ml). * Cell activity = total counts recovered in cell fractionation. Supernatant and washings not included.

** RNA = uracil-2-C¹⁴ incorporated into hot TCA-soluble material; protein = phenylalanine-U-C¹⁴ incorporated into hot NaOH-soluble material.

Figure 1. Incorporation of uracil-2-C¹⁴ 5 µc, 0.075 µmolar, 10 ml) into RNA in "germinating" (G) and "non-germinating" (U) conidia. SG = spore germination.

Figure 2. Incorporation of phenylalanine-U-C¹⁴ (1 µc, 1.27 µmolar, 10 ml) into protein. Symbols same as for Figure 1.

Table 1 shows that the large difference in uracil incorporated into RNA shown in Figure 1 reflects a difference in the abilities of the cells to take umcil into the cell, rather than gross differences in synthetic ability. In 2 hours there is a 6-fold difference in uptake of uracil by conidia incubated in minimal plus glucose, while there is only a slightly increased uptake by the conidia incubated in mineral salts only. In the "non-germinating" conidia, much of the uracil taken up is incorporated, indicating that these cells are capable of RNA synthesis. On the other hand, phenylalanine is taken up more readily by the "non-germinating" conidia than by the "germinating" conidia. A considerable fraction of this is incorporated into protein (24.8 % at 4 hours), but not as much as in the "germinating" conidia (70.1% at 4 hours).

These results show that conidia incubated either in mineral salts or in mineral salts plus glucose are capable of RNA and protein synthesis and stress the importance of looking at uptake of the precursor as well as its incorporation into RNA or protein.

This research was supported by National Science Foundation Grant No. GB 1401. - - - Department of Biology, Wesleyan University, Middletown, Connecticut 06457.