

Determination of protein in conidia

G. Schneider

W. Klingmuller

W. Sebald

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Schneider, G., W. Klingmuller, W. Sebald, and F. Kaudewitz (1969) "Determination of protein in conidia," *Fungal Genetics Reports*: Vol. 14, Article 14. <https://doi.org/10.4148/1941-4765.2045>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Determination of protein in conidia

Abstract

Determination of protein in conidia

Authors

G. Schneider, W. Klingmuller, W. Sebald, and F. Kaudewitz

Schneider, G., W. Klingmüller, W. Sebald

and F. Kaudewitz. Protein determination
in conidia of Neurospora crassa.

During experiments in which the rate of uptake of different sugars by conidia of a number of different strains of Neurospora crassa was to be estimated and referred to their protein content, a quick and reliable method for measuring the protein content of ungerminated conidia was needed. For technical reasons, we first tried to measure their protein content directly by applying a modification of the Biuret method as described by Beisenherz et al. (1953 Z. Naturforschung 8b: 555) to whole conidia.

Individual steps of this modification are as follows: (1) Prepare conidial suspensions of 5×10^7 conidia/ml. (2) Precipitate the protein in 1 ml of the conidial suspension with 0.2 ml 50% TCA, shake well and spin down. (3) Wash two times with a mixture of 3 parts ethyl alcohol and one part ethyl ether and a third time with ethyl ether to remove carotenoid color and other lipids. (4) Let the ether evaporate from the pellet and resuspend in 1 ml Biuret reagent. (5) Incubate for 30 min. at room temperature on a shaker. Centrifuge and transfer the supernatant into a 1 cm cuvette. (6) Read the absorption at 546 m μ . Add a small amount of KCN powder to the cuvette, stir and wait 1-2 min. until the remaining absorption is constant, and read again. The difference in absorption is due to the protein in the solution (A_p). (7) Measure the absorption of the Biuret reagent alone before and after addition of KCN. The difference in the blank absorption (A_b) is to be subtracted from A_p . (8) Refer $A_p - A_b$ to a calibration curve, obtained for bovine serum albumin to obtain mg protein/ml.

Applying this procedure and increasing the incubation time at room temperature from 30 min. to 120 min., a marked increase of the resulting absorption was observed (Fig. 1, curve a). This finding was thought to indicate that not all of the protein in a whole conidium was available freely to the Biuret-reagent. This was confirmed by increasing the temperature of incubation up to 50°C, when a further increase in absorption was observed (Fig. 1, curve b). Neither elongation of incubation-time up to 120 min. nor increase of incubation-temperature up to 50°C had a significant effect on absorption of calibration samples containing bovine serum albumin.

In additional experiments, conidia were therefore disrupted either by treating them 1 to 3 times with the X-press (AB Biox, Box 235, Nacka 2, Sweden) with a 25 ml volume operated at -25 to -35°C at ca. 2000 kg/sq cm, or by smashing them with glass beads (d = 0.45-0.50 mm) for 5 to 10 minutes in a homogenizer




Fig. 1. Values for whole or disrupted conidia (5×10^7 /ml) after incubation with the reagent for different periods of time.

(Vibrogen-Zellmühle, Fa. E. Bühler, Tübingen, Germany). The results obtained by these disintegration methods document a further increase in protein content of the supernatant (Fig. 1, curve c for 3 times X-pressing, curve d for glass-bead disintegration). They demonstrate that in the broken cells either protein is available faster to the reagent or additional protein is set free to react with the reagent. Visual observation of the state of the cells by means of phase contrast microscopy showed less than 5% intact conidia after glass-bead disintegration, but co. 50-25% after 1-3 X-pressings, respectively, a finding which may explain the fact that the latter procedure gives a lower protein value than the former. To obtain reliable data on the protein content of *Neurospora* conidia, the complete disruption of the cells is therefore an essential prerequisite.

We are grateful to Dipl. Biol. R. Schweyen and Miss D. Hacker for valuable technical suggestions. This work was supported by the Deutsche Forschungsgemeinschaft. ■ ■ ■ Institut für Genetik, Universität München, München, Germany.