

Decreased activity of all tryptophan biosynthetic enzymes due to zinc deficiency

M. Carsiotis

A. M. Meyers

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Carsiotis, M., and A.M. Meyers (1965) "Decreased activity of all tryptophan biosynthetic enzymes due to zinc deficiency," *Fungal Genetics Reports*: Vol. 8, Article 2. <https://doi.org/10.4148/1941-4765.2073>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Decreased activity of all tryptophan biosynthetic enzymes due to zinc deficiency

Abstract

Decreased activity of all tryptophan biosynthetic enzymes due to zinc deficiency

Carsiotis, M. and A. Meyers. Decreased activity of all tryptophan biosynthetic enzymes due to zinc deficiency.

decreased two- to five-fold when zinc is omitted from the growth medium (Vogel's Medium N). The effect is more marked in a histidine mutant than in the wild type. An explanation of this effect is being sought currently. * * * Department of Microbiology, University of Cincinnati College of Medicine, Cincinnati, Ohio.

A reinvestigation of the effect of zinc deficiency noted by Nason et al. (1951 J. Biol. Chem. 188:397) has revealed that all enzymes of the tryptophan biosynthetic pathway are