Ureaseless mutants in *Neurospora crassa*

H. G. Kolmark

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Kolmark, H. G. (1965) "Ureaseless mutants in *Neurospora crassa*," *Fungal Genetics Reports*: Vol. 8, Article 7. <https://doi.org/10.4148/1941-4765.2086>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Ureaseless mutants in *Neurospora crassa*

Abstract

Ureaseless mutants in *Neurospora crassa*

Kølmær, H. G. Ureaseless mutants in Neurospora crassa.

The presence of the enzyme urease in *Neurospora* has been known for many years (Srb and Horowitz 1944 J. Biol. Chem.

154: 129). Mutation to an ureaseless condition is therefore to be expected.

In the present experiments it was first found that urease could very easily be demonstrated directly on a growing colony, simply by rooking a small piece of pH-indicator paper (range about pH 6-8) in an 8% solution of urea in water and placing it in contact with the mycelium or conidia. A color change towards the alkaline takes place in a few minutes due to liberation of ammonia when urea is enzymatically hydrolysed.

This change in pH was utilized in the screening methods finally worked out for isolation of ureaseless mutants.

Strains of Neurospora: Some earlier experiments with macroconidiating strains of colonial morphology did not succeed in the isolation of ureaseless mutants. A microconidiating, small colonial strain, 398-28 A, was then tried in the expectation that mutants might more easily become phenotypically expressed when induced in mononucleate conidia. A morphological mutant, m-25, derived from this strain after treatment with ultraviolet light, was used in some of the later experiments. m-25 forms very small and extremely dense colonies.

Experimental Procedures: Suspensions of conidia in saline were irradiated with UV and, after appropriate dilutions, incubated in 25 ml of liquid minimal medium in petri dishes kept at 21-22°C. Under these conditions the colonies grow below the surface and adhere to the gloss bottom of the plate. After 6-7 days they are ready to be tested for urease activity.

The medium is then decanted off. With careful handling the colonies stick to their growth place on the gloss. They are rinsed twice with saline. After the last rinsing the plates are placed in a slanted position for a few minutes to let superficial saline drain off. A filter paper is moistened with a mixture of 4% (or stronger) urea and a pH indicator (brom cresol purple or brom thymol blue) in the range 5.5-7.5, adjusted to the lower pH with a weak phosphate buffer (ionic strength 0.01). When the paper is pressed against the colonies with a glass plate, all normal colonies change the indicator color in a few minutes. Non-reacting colonies are isolated into tubes with minimal medium. Established cultures are retested later in comparison with the original strain.

In an alternative procedure, the UV-treated conidia are spread on the surface of solid medium. A white net of nylon fabric, ca. 3 threads per mm, is placed on top of the inoculated medium. The colonies, appearing between the meshes, are after 3 days overlaid with 15 ml of liquid minimal medium and the incubation continued for 5 days more. The liquid medium is added because of the finding that the enzymatic reaction proceeds more readily when the mycelium has grown submerged. The net is stripped off with part of the colonies attached, when ready for testing. The screening test is done on this replica as described above. Isolations are made from the untested part of the colonies left on the plate.

Two cultures out of a total of 58 isolates were by retesting found to be entirely lacking in urease activity, as evidenced by no shift towards alkalinity during 4-5 days when a small amount of mycelium was incubated at 34°C in 0.25 ml of 4% urea, dissolved in 0.01 ionic strength phosphate buffer. The reaction is started at clear yellow (pH ca. 6) with the indicator brom thymol blue. The original strain gives a strongly positive reaction in 2 hours.

Several other isolates appear to have a decreased urease activity. The frequency of ureaseless mutants was found to be about 1 per 10⁴ survivors. The ureaseless mutants are unable to grow on urea as the sole source of nitrogen, while the original strains grow well. Otherwise, the mutants show no decreased vitality with regard to germination of conidia or ascospores nor in mycelial growth under the conditions observed.

Genetic testing: One of the ureaseless mutants, u-9 A, was crossed to the wild type strain 74-ORB-1a. The ureaseless character was found to segregate regularly at 1st or 2nd reduction division in dissected asci. Ascospore cultures with ureaseless recombined into wild type morphology or fully fertile also as protoperithecial parents. Genetic testings to identify the linkage group are in progress, but so far it can only be stated that it seems not to be situated in the linkage groups I, II or VII. It is proposed that the mutant phenotype described above be designated by the symbol ur, since this symbol seems not previously to be in use as a locus symbol in *Neurospora*.

This work was supported by grants from the British Council and from O. E. C. D. during a stay at The Hartley Botanical Laboratories, Liverpool University, England and, presently, by grants from the Swedish Research Council, - - - Institute of Physiological Botany, University of Uppsala, Sweden.