Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1469

1968

The effects of feeding a high concentrate ration containing 25% ground beef manure to fattening heifers in concrete and soil-surfaced lots

C.L. Drake

L.I. Smart

E.F. Smith

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Drake, C.L.; Smart, L.I.; Smith, E.F.; and Lipper, R.I. (1968) "The effects of feeding a high concentrate ration containing 25% ground beef manure to fattening heifers in concrete and soil-surfaced lots," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2872>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1968 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

The effects of feeding a high concentrate ration containing 25% ground beef manure to fattening heifers in concrete and soil-surfaced lots

Authors

C.L. Drake, L.I. Smart, E.F. Smith, and R.I. Lipper

The Effects of Feeding a High Concentrate Ration
Containing 25% Ground Beef Manure to Fattening
Heifers in Concrete and Soil-surfaced Lots (Project 660)

C.L. Drake, L.I. Smart, E.F. Smith and R.I. Lipper¹

This is a cooperative project with the Department of Agricultural Engineering to study surface runoff, in addition to animal performance.

Purpose

A. To compare a ration containing 25% ground beef manure with a conventional ration.

B. Study the influence of concrete or soil surfacing on animal performance.

Procedure

Twenty Hereford heifers were randomly allotted to four equal sized lots: 2 surfaced with concrete and 2 with soil. Self-feeders in soil-surfaced lots have concrete aprons.

Manure was collected from pens of cattle receiving a high energy ration, dried and processed through a hammer mill with the screen removed. It was fed mixed with other ingredients in the self-feeders. The ration containing manure consisted of 25% manure, 67.5% ground sorghum grain and 7.5% soybean meal. The control ration con-

¹ Department of Agricultural Engineering

tained 33.5% alfalfa crumbles and 66.5% ground sorghum grain. Chemical analyses of feed ingredients are in table 29 and feedlot and carcass data, in table 30.

Results

Although differences were small, heifers on concrete-surfaced lots gained slightly more and were more efficient. Concrete-surfaced lots are faster and easier to clean than unsurfaced lots, especially during wet weather.

Feed efficiency was the same for control heifers in both concrete- and soil-surfaced lots; however, heifers receiving 25% beef manure required more feed per pound of gain. Differences in feed costs per hundred weight gain were small in all cases.

Heifers receiving manure tried to separate it. A lower percentage of manure and pelleting the entire ration might have increased gains and reduced waste in the manure ration.

Samples of manure were analyzed in the veterinary diagnostic laboratory and found free of Salmonella organisms.

Table 29

Chemical Analyses of Feed Ingredients

	Feed ingredient		
	<u>Sorghum grain</u>	<u>Beef manure</u>	<u>Dehydrated¹ alfalfa crumbles</u>
Analysis, %:			
Protein (nitrogen x 6.25)	10.19	18.69	20.19
Ether extract (fat)	2.87	2.74	3.72
Crude fiber	1.87	13.79	20.68
Moisture	11.96	12.01	7.32
Ash	1.56	26.96	11.22
Nitrogen-free extract	71.55	25.81	36.87

¹ Dehydrated alfalfa crumbles were analyzed for protein only.

Table 30

The Influence of 25% Ground Beef Manure
In Rations of Heifers on
Concrete or Soil-surfaced Lots

Ration	Type of surface			
	Concrete		Soil	
	Manure	Control	Manure	Control
Initial wt., lbs. July 18, 1967	713	724	721	716
Final wt., lbs. Oct. 5, 1967	877	944	876	925
Total gain, lbs.	164	220	155	209
Av. daily gain, lbs.	2.08	2.78	1.96	2.65
Lbs. feed per lb. gain	11.95	9.09	12.64	9.09
Cost per cwt.# ¹	20.53	20.63	21.72	20.63
Carcass Data:				
Hot carcass wt., lbs.	546	577	545	558
Dressing, % ²	62.3	61.1	62.2	60.3
Fat thickness, 12th rib, in.	0.33	0.34	0.39	0.35
Loin eye area, sq.in.	11.70	11.57	10.10	10.67
U.S.D.A. grade ³	17	17	17	18

¹ Manure valued at \$5 per ton.

² Based on hot carcass weight

³ Carcass grade score: Average good, 17; High good, 18.