

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 2 *Dairy Research (1984-2014)*

Article 89

1988

Bovine recombinant interleukin-2 enhances resistance to bovine herpesvirus-1: Dose response trial (1988)

P.G. Reddy

Frank Blecha

J.L. Morrill

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Dairy Science Commons](#)

Recommended Citation

Reddy, P.G.; Blecha, Frank; Morrill, J.L.; and Minocha, H.C. (1988) "Bovine recombinant interleukin-2 enhances resistance to bovine herpesvirus-1: Dose response trial (1988)," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 2. <https://doi.org/10.4148/2378-5977.3014>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1988 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



**Bovine recombinant interleukin-2 enhances resistance to bovine herpesvirus-1:
Dose response trial (1988)**

Authors

P.G. Reddy, Frank Blecha, J.L. Morrill, and H.C. Minocha

K**S****U****BOVINE RECOMBINANT INTERLEUKIN-2 ENHANCES
RESISTANCE TO BOVINE
HERPESVIRUS-1: DOSE RESPONSE TRIAL****P.G. Reddy¹, F. Blecha¹,
J.L. Morrill, and H.C. Minocha²**

Summary

Twenty-five calves were allotted to five groups: controls that did not receive bovine recombinant interleukin-2 (rIL-2) and four groups that received 5 daily injections of rIL-2 at 11.4, 1.1, 0.11, or 0.011 $\mu\text{g}/\text{lb}/\text{day}$. On day 0 of the experiment, all calves received bovine herpesvirus-1 (BHV-1) vaccine and the first of the 5 daily injections of bovine rIL-2. All calves were infected with BHV-1 on day 21 of the experiment. Calves treated with 11.4 $\mu\text{g}/\text{lb}/\text{day}$ had elevated rectal temperatures and mild diarrhea during administration of rIL-2. All other calves were normal. Compared to control calves, those treated with 11.4, 1.1, and 0.11 $\mu\text{g}/\text{lb}/\text{day}$ had higher ($P<0.05$) serum antibody titers to BHV-1 and following challenge lower ($P<0.05$) BHV-1 titers in nasal secretions. Additionally, clinical disease as evidenced by nasal and ocular discharge was less severe. Cytotoxic responses against BHV-1-infected bovine kidney cells were increased ($P<0.05$) in calves treated with rIL-2 in a dose dependent manner. These data suggest that bovine rIL-2 at doses of 0.11 to 1.1 $\mu\text{g}/\text{lb}/\text{day}$ for 5 days may enhance immunity against BHV-1 without causing adverse side effects.

Introduction

Interleukin-2 (IL-2) is a glycoprotein secreted by a subset of T cells and large granular lymphocytes after stimulation with mitogen or antigen. This lymphokine induces the clonal expansion of activated T cells and B cells and activates natural killer cells.

Recently, we evaluated the use of bovine rIL-2 in bovine herpesvirus-1 (BHV-1)-vaccinated and -challenged calves. Our data indicated that bovine rIL-2 (11.4 $\mu\text{g}/\text{lb}$ body weight) used in conjunction with BHV-1 vaccination enabled calves to withstand a challenge with virulent BHV-1 better than a vaccination alone, suggesting that bovine rIL-2 may be an effective adjuvant to immunization against viral diseases. However, the dose of bovine rIL-2 used in that study caused mild fever and diarrhea, which abated immediately after rIL-2 treatment was stopped.

The objective of the present study was to determine if a lower dose of bovine rIL-2 would enhance resistance to BHV-1 without causing adverse side effects.

Procedures

Twenty-five Holstein or crossbred beef calves, 4 to 6 months old and seronegative for BHV-1, were used. The calves were allotted by weight to one of five treatment groups; controls that did not receive rIL-2 and four groups that received intramuscular injections of bovine rIL-2 at 11.4, 1.1, 0.11, or 0.011 $\mu\text{g}/\text{lb}/\text{day}$ for 5 days.

¹Department of Anatomy and Physiology.

²Department of Laboratory Medicine.

At the start of the experiment (day 0), calves were vaccinated with a modified-live BHV-1 vaccine (Norden Laboratories, Lincoln, NE) and their respective doses of bovine rIL-2 were administered intramuscularly. Bovine rIL-2 was administered daily for 4 subsequent days. Calves that did not receive bovine rIL-2 were injected intramuscularly with an equivalent volume of saline for 5 consecutive days. On day 21, all calves were inoculated intranasally (2 ml) and conjunctivally (1 ml) with 10^7 PFU/ml of BHV-1.

Results and Discussion

Treatment of calves with 11.4 $\mu\text{g/lb/day}$ of bovine rIL-2 resulted in elevated rectal temperatures and mild diarrhea, but these signs subsided immediately after the last injection of bovine rIL-2. Calves in other rIL-2 treatment groups remained normal and healthy during the administration of rIL-2.

Following challenge with virulent BHV-1, calves that received 11.4, 1.1, and 0.11 $\mu\text{g/lb/day}$ showed significantly lower ($P < 0.05$) rectal temperatures on days 2 to 5 postinfection than control calves. There was a similar trend with regard to clinical signs of BHV-1 infection, with nasal and ocular discharges from rIL-2-treated calves being significantly less severe ($P < 0.05$) than those from control calves.

Cytotoxic responses against BHV-1-infected bovine kidney cells by peripheral blood mononuclear cells (PBMC) from control calves remained low at all sampling times, whereas the cytotoxic response by PBMC from calves treated with rIL-2 increased in a dose dependent manner. Following rIL-2 injections, cytotoxic responses were significantly higher for PBMC from calves receiving 11.4, 1.1, and 0.11 $\mu\text{g/lb/day}$ at all sampling times than for PBMC from control calves.

Calves treated with rIL-2 at 11.4 and 1.1 $\mu\text{g/lb/day}$ had higher titers of ($P < 0.05$) serum neutralizing antibodies to BHV-1 on days 10, 15, 21, and 25, and calves treated with 0.11 $\mu\text{g/lb/day}$ had higher titers on days 15 and 21 ($P < 0.05$), compared to control calves.

Shedding of BHV-1 in nasal secretions was significantly lower on days 1 to 6 postinfection in calves treated with 11.4 and 1.1 $\mu\text{g/lb/day}$ and on days 3 and 4 postinfection in calves treated with 0.25 $\mu\text{g/lb/day}$, compared to control calves.

Discussion

Interleukin-2 augmentation of antiviral immunity has been demonstrated by adoptive transfer of immune lymphocytes from IL-2-treated animals and treatment of cattle with bovine rIL-2. The present study, while indicating a dose response of calves to bovine rIL-2 administered in conjunction with BHV-1 vaccine, also demonstrated that augmentation of immunity is possible with a dose that would not cause any visible adverse side effects. Although the optimum dose required should be confirmed by field trials, this study and our earlier study clearly demonstrate the potential benefit of bovine rIL-2 as an immunoadjuvant to vaccines against the bovine respiratory disease complex (BRD).

Some concern has been expressed about the effectiveness of vaccines alone as a means to combat BRD, which according to current estimates is causing an annual economic loss in the range of \$250 million to \$1 billion. The efficacy of vaccines may often be compromised by colostrally acquired maternal antibodies and certain states of stress-induced immunosuppression. This suggests the need for additional ways to modulate the immunity of cattle to resist BRD, and our study indicates the practical value of immunotherapeutic use of bovine rIL-2.