Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 182

1978

Using oral E. Coli milk vaccine to control colibacillosis

D A. Schoneweis

V Gaeth

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Schoneweis, D A. and Gaeth, V (1978) "Using oral E. Coli milk vaccine to control colibacillosis," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6022>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1978 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Colibacillosis is still a costly disease to swine producers in spite of new antibiotics and other products and procedures used to prevent or treat the disease. A recent aid in controlling the disease is the oral E. coli milk vaccine developed by Dr. Irvin Kohler. It consists of isolating a pathogenic E. coli from the herd and feeding the organism to pregnant sows to stimulate colostral antibodies. It is a giant step forward from the earlier recommendation of the late Dr. Howard Dunne who suggested taking the bedding and feces from the farrowing house and feeding it to pregnant sows to elevate the antibody content of the colostrum. Advantages of oral vaccine include:

1. Stimulates antibodies against the E. coli serotype on the farm. (Do not use a serotype on any farm except where isolated.)
2. Oral vaccine apparently stimulates more IgA and IgM antibodies than the parenteral injection of a bacterin does.
3. Less chance of spreading other diseases such as Salmonellosis from feeding feces of sows to other sows.

The procedure requires the isolation of the pathogenic E. coli from the herd and the subsequent feeding of large populations of the organisms to sows in late gestation. The production of the oral vaccine should only be attempted by persons with a

knowledge of microbiology; and, it is recommended that a veterinarian be consulted to set up the program which can be readily adapted to most swine operations. The oral E. coli vaccine has effectively reduced the colibacillosis problem on many swine operations.

Research recently conducted at Kansas State indicates that non-fat dry milk can be reconstituted with chlorinated tap water with no deleterious effects upon the E. coli population and the use of milk previously inoculated and incubated is not a satisfactory substitute for inoculated BHI broth. It is recommended that the temperature be held at 37-38 C (98-100 F); however, a lower temperature will still result in growth of the E. coli. The use of milk as a preinoculation media resulted in lowered populations of E. coli in the oral vaccine.