Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 228

1981

Duration of plasma oxytetracycline levels in neonatal pigs

D Schoneweis

S Hummels

L Schulteis

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Schoneweis, D; Hummels, S; and Schulteis, L (1981) "Duration of plasma oxytetracycline levels in neonatal pigs," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6068>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1981 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

K

Duration of Plasma Oxytetracycline Levels in Neonatal Pigs

S

Dave Schoneweis¹, Susan Hummels¹ and Lee Schulteis¹

U

Summary

Samples from neonatal pigs were evaluated to determine the persistence of plasma oxytetracycline after injecting LA 200[®]. Significant (minimum of .2 mcg/ml) amounts of oxytetracycline were found in all samples 96 hours after the injection, and oxytetracycline still persisted in many pigs after 168 hours.

Procedure

Pigs were injected with 1 ml of LA 200 at 3, 12, and 21 days of age. Blood samples were collected at 4, 12, 24, 48, and 96 hours after the oxytetracycline injection. The range and number of samples exceeding .2 mcg/ml for the different-age pigs are shown in Table 39.

Table 39. Duration of Plasma Oxytetracycline Levels in Neonatal Pigs

Age at injection	Range (mcg/ml)				
	Number of samples >.2 mcg/number of samples				
	Hours post-injection (PI)				
	4	12	24	96	168
3 days	<u>10.0 - 32.9</u> 9/9	<u>6.6 - 30.2</u> 9/9	<u>5.6 - 23</u> 8/8	<u>.5 - 1.6</u> 9/9	<u>0 - .5</u> 4/7
12 days	<u>7.4 - 10.3</u> 8/8	<u>4.1 - 8.4</u> 8/8	<u>1.7 - 4.8</u> 8/8	<u>.4 - .9</u> 8/8	<u>0 - .4</u> 5/8
21 days	<u>3.0 - 5.2</u> 8/8	<u>2.0 - 5.8</u> 8/8	<u>1.2 - 2.1</u> 6/6	<u>.5 - .9</u> 8/9	<u>0 - .3</u> 2/9

* One sample showed trace at 96 hours, but had a .3 mcg/ml at 168 hours.

Results

Significant plasma levels of oxytetracycline were noted in samples at 96 hours post injection. The one pig having only a trace in the 96-hour test exhibited .3 mcg/ml at 168 hours. All pigs were injected with 1 ml of LA 200, and consequently the 4-, 12-, and 24-hour samples were highest in the 3-day-old pigs and lowest in 21-day-old pigs; however, the difference was not so great at the later bleedings. The effectiveness of oxytetracycline depends on the susceptibility of the infective agent to the antibiotic, and a satisfactory response cannot be anticipated solely because of a continuous plasma level.

¹Department of Surgery and Medicine

[®]Chas. Pfizer and Co.