Kansas Agricultural Experiment Station Research Reports

Volume 0 Issue 10 Swine Day (1968-2014)

Article 229

1981

Value of adding fat to finishing diets to alleviate heat stress (1981)

David A. Nichols

D R. Ames

G L. Allee

See next page for additional authors

Follow this and additional works at: https://newprairiepress.org/kaesrr



Part of the Other Animal Sciences Commons

Recommended Citation

Nichols, David A.; Ames, D R.; Allee, G L.; and Hines, Robert H. (1981) "Value of adding fat to finishing diets to alleviate heat stress (1981)," Kansas Agricultural Experiment Station Research Reports: Vol. 0: Iss. 10. https://doi.org/10.4148/2378-5977.6069

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1981 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



| Value of adding fat to finishing diets to alleviate heat stress (1981) | | | | |
|--|--|--|--|--|
| Authors David A. Nichols, D R. Ames, G L. Allee, and Robert H. Hines | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |



Value of Adding Fat to Finishing Diets to Alleviate Heat Stress





D.A. Nichols, D.R. Ames, R.H. Hines and G.L. Allee

Summary

Crossbred barrows , observed at temperatures of 62 and $95^{0}F$, were fed either a control or a fat added diet. Pigs housed at $95^{0}F$ ate significantly less feed, gained slower, and were less efficient than littermates housed at $62^{0}F$.

Fat addition to the diet did not significantly influence performance when fed at $62 \text{ or } 95^{0}\text{F}$. The addition of fat did not appear to reduce heat stress or improve performance when fed at the higher temperature.

Introduction

The major environmental factor influencing swine performance is temperature. During exposure to heat stress, energy for maintenance increases due to increased energy demands for panting and other methods of heat loss.

Manipulating diet during heat stress, such as by adding fat, might help alleviate the production problems of restricted growth, reduced appetite and reduced feed efficiency. We attempted to determine if fat was beneficial in reducing heat stress.

Procedure

Thirty-two crossbred barrows of similar genetic background were obtained for the study. They averaged 160 lbs. at the start of each trial.

The experiment was divided into two 28 day feeding trials. Temperatures of $62^{0}F$ and $95^{0}F$ were studied. Pigs were fed either a control sorghum soy-diet or the control diet with 5% fat added. All pigs were housed at $59^{0}F$ five days prior to the start of each trial. Temperature and lighting were constant and water and feed were supplied ad libitum. Temperatures were controlled in two environmental rooms (10 ft. x 12 ft.) divided into four equal sized pens with two pigs per pen housed on concrete slatted floors.

Results and Discussion

Table 34. Effect of Temperature and Fat Addition on Performance

| Diet | | 62 ⁰ Temperature F | |
|-------------------|---|-------------------------------|---------------------|
| Control Fat Added | Daily feed intake (lbs.) Avg. daily gain (lbs.) F/G | 7.00 1.81 3.87 | 3.54 .75 4.72 |
| | Daily feed intake (lbs.) Avg. daily gain (lbs.) F/G | 6.84 1.78 3.84 | 3.49 .79 4.45 |

Temperature and fat addition effects on performance are shown in Table 34. Pigs housed at 95° F ate significantly (P<.01) less feed and gained significantly (P<.01) slower than those housed at 62° F. Feed to gain ratio also increased at the higher temperature, but the difference was not significant in our study due to relatively few observations.

Adding fat at $62^{0}F$ or $95^{0}F$ did not significantly influence daily feed intake, daily gain or feed to gain ratio. Though added fat tended to improve feed to gain ratio more at the higher temperature, the difference was not significant.