

# Kansas Agricultural Experiment Station Research Reports

---

Volume 0  
Issue 10 *Swine Day (1968-2014)*

Article 469

---

1991

## Causes of diarrhea, pneumonia, and septicemia in swine for 1991 submissions to the KSU Veterinary Diagnostic Laboratory (1991)

R K. Frank

M W. Vorhies

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

---

### Recommended Citation

Frank, R K. and Vorhies, M W. (1991) "Causes of diarrhea, pneumonia, and septicemia in swine for 1991 submissions to the KSU Veterinary Diagnostic Laboratory (1991)," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6309>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1991 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



**K**

**S**

**U**

**CAUSES OF DIARRHEA, PNEUMONIA, AND SEPTICEMIA  
IN SWINE FOR 1991 SUBMISSIONS TO THE KSU  
VETERINARY DIAGNOSTIC LABORATORY**

*R. K. Frank and M. W. Vorhies<sup>1</sup>*

---

**Summary**

Causes of pre- and postweaning diarrhea, pneumonia, and bacterial septicemia in pigs were summarized for fiscal year 1991 (July, 1990 to June, 1991) for submissions to the Kansas State University Veterinary Diagnostic Laboratory. *Escherichia coli* was the most common cause of both pre- and postweaning diarrhea in pigs (33.5% and 25.0%, respectively, of submissions for diarrhea). Other commonly diagnosed causes included transmissible gastroenteritis (24.4%) and coccidiosis (16.5%) for preweaning diarrhea, and proliferative enteritis (19.2%) and salmonellosis (13.2%) for postweaning diarrhea. The most commonly diagnosed causes of pneumonia in nursing, growing, and finishing pigs were *Pasteurella multocida*, *Mycoplasma*, and *Actinobacillus (Haemophilus) pleuropneumoniae*. *Streptococcus* and *Salmonella* were common causes of bacterial septicemia in Kansas pigs.

(Key Words: Disease, Diagnosis, Diarrhea, Pneumonia.)

**Introduction**

Enteric and respiratory diseases account for large economic losses to the swine industry each year. An accurate diagnosis of the cause is essential for effective prevention and control of these diseases. The present summary was performed to demonstrate the relative importance of various causes of enteric, respiratory, and septicemic diseases as determined at the diagnostic laboratory level.

**Procedures**

Diagnoses and age were summarized for cases of diarrhea, pneumonia, and bacterial septicemia in pigs from computer records for submissions to the Kansas State University Veterinary Diagnostic Laboratory for fiscal year 1991 (July, 1990 to June, 1991). Specimens included living or dead pigs and/or tissues. A diagnosis was made following light microscopic, bacteriologic, and virologic examination of tissues and summarized in a computer data base.

**Results and Discussion**

Causes and number of cases of pre- and postweaning diarrhea, pneumonia, and septicemia are summarized in Tables 1 and 2, respectively. In spite of extensive vaccination and effective vaccines for the control of colibacillosis, *E. coli* still remains the most frequent cause of both pre- and postweaning diarrhea in pigs in Kansas. Transmissible gastroenteritis was a much more common cause of preweaning than postweaning diarrhea and frequently recurred in sequential farrowings (endemic or chronic TGE). The highest incidence of coccidiosis was in the months of July and August (12/27 cases). Proliferative enteritis (terminal ileitis) is an important cause of diarrhea in Kansas swine herds. Cases of diarrhea with an idiopathic diagnosis were those for which no cause could be determined by routine laboratory testing. The primary reasons for no diagnosis in cases of diarrhea were 1) submitting tissues from pigs too late in

---

<sup>1</sup>Department of Veterinary Diagnosis.

the disease process, 2) advanced postmortem change in submitted tissues, and 3) improper tissue collection by the referring veterinarian.

*Pasteurella multocida*, *Mycoplasma*, and *Actinobacillus (Haemophilus) pleuropneumoniae* were the most commonly diagnosed causes of respiratory disease in pigs (Table 3). Frequently more than one bacterial species was isolated from the same lung. Anti-

biotic treatment is one likely explanation for the large number of idiopathic diagnoses. Additionally, some lungs were submitted fixed in formalin, with no unfixated tissue available for bacterial culturing.

*Streptococcus* and *Salmonella* were the most common causes of bacterial septicemia in Kansas pigs, accounting for 33.3 and 26.7% of all cases, respectively (Table 4).

**Table 1. Causes of Prewaning (< 3 Weeks of Age) Diarrhea in Pigs for Submissions to the KSU Veterinary Diagnostic Laboratory (July, 1990 to June, 1991)**

Cause/Disease	No. cases	% of total cases
<i>E. coli</i>	55	33.4
TGE	40	24.4
<i>Isospora suis</i> (Coccidia)	27	16.5
Idiopathic <sup>a</sup>	16	9.8
Rotavirus	8	4.9
<i>Clostridium perfringens</i> type C	7	4.3
Viral <sup>b</sup>	7	4.3
Other	4	2.4
<b>Total</b>	<b>164</b>	<b>100.0</b>

<sup>a</sup>Exact cause of the diarrhea could not be determined.

<sup>b</sup>Microscopic intestinal changes were consistent with a viral diarrhea, but a virus was not identified by routine testing.

**Table 2. Causes of Postweaning ( $\geq$  4 weeks of age) Diarrhea in Pigs for Submissions to the KSU Veterinary Diagnostic Laboratory (July, 1990 to June, 1991)**

Cause/Disease	No. cases	% of total cases
<i>E. coli</i>	30	25.0
Proliferative enteritis (Terminal ileitis)	23	19.2
<i>Salmonella</i>	16	13.3
Idiopathic <sup>a</sup>	9	7.5
Swine dysentery	8	6.7
Non-specific colitis	8	6.7
TGE	7	5.8
Necrotic enteritis <sup>b</sup>	7	5.8
Hemorrhagic bowel syndrome	5	4.2
Viral <sup>c</sup>	4	3.3
Rotavirus	3	2.5
Total	120	100.0

<sup>a</sup>Exact cause of the diarrhea could not be determined.

<sup>b</sup>No cause was identified, but often the end result of *Salmonella* infection or swine dysentery.

<sup>c</sup>Microscopic intestinal changes were consistent with a viral diarrhea, but a virus was not identified by routine testing.

**Table 3. Causes of Pneumonia in Nursery, Growing, and Finishing Pigs for Submissions to the KSU Veterinary Diagnostic Laboratory (July, 1990 to June, 1991)**

Cause/Disease	No. cases <sup>a</sup>	% of total cases
<i>Mycoplasma</i>	44	21.2
<i>Pasteurella multocida</i>	42	20.2
<i>Actinobacillus (Haemophilus)</i> <i>pleuropneumoniae</i>	24	11.5
Idiopathic <sup>b</sup>	23	11.0
<i>Streptococcus suis</i>	22	10.6
<i>Streptococcus</i> (not <i>suis</i> )	19	9.1
<i>Bordatella bronchiseptica</i>	5	2.4
Swine influenza	4	1.9
Other bacterial	23	11.0
Misc.	2	1.0
Total	208	100.0

<sup>a</sup>Isolates were from approximately 150 different submissions.

<sup>b</sup>Exact cause of the pneumonia could not be determined.

**Table 4. Causes of Septicemia in Nursery, Growing, and Finishing Pigs for Submissions to the KSU Veterinary Diagnostic Laboratory (July, 1990 to June, 1991)**

Cause/Disease	No. cases	% of total cases
<i>Streptococcus</i> (not <i>suis</i> )	20	33.3
<i>Salmonella</i>	16	26.7
Edema disease	8	13.3
<i>Streptococcus suis</i>	6	10.0
Erysipelas	6	10.0
Other	4	6.7
<b>Total</b>	<b>53</b>	<b>100.0</b>



**Melanie Krause analyzing samples in the swine research lab.**