The production of mosaic mutations in *Neurospora crassa*

J. Royes

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Royes, J. (1962) "The production of mosaic mutations in *Neurospora crassa*," *Fungal Genetics Reports*: Vol. 1, Article 5. <https://doi.org/10.4148/1941-4765.1019>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

The production of mosaic mutations in *Neurospora crassa*

Abstract

The production of mosaic mutations in *Neurospora crassa*

Royes, J. The production of mosaic mutations in Neurospora crassa.
arginineless and methionineless anmycelial.

Nitrous acid, ultra violet light and X-rays have been used for producing recessive lethals in conidia of a balanced heterokaryon between
A modification of Atwood's method showed that the

proportion of mosaics among mutations decreased from nitrous acid (77-87%) to UV (40%) to X-rays (28%). A substantial proportion of the X-ray induced mosaics and a smaller proportion of those induced by the other mutagens may have been derived from conidia with more than 1 amycelial nucleus (only lethals induced in such nuclei are detected). The excess of mosaicism, especially after treatment with nitrous acid, is attributed either to segregation of mutated and non-mutated strands in the treated chromosome or to delayed mutation. In most mosaics, and independent of the mutagen, the lethal sector was much smaller than one half; sectors comprising 20% or less formed between 70 and 78%. Possible reasons for this are being examined.