

## Growth of a histidina strain, hist-2 (C94), of *Neurospora crassa*

C. McNelly

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

McNelly, C. (1962) "Growth of a histidina strain, hist-2 (C94), of *Neurospora crassa*," *Fungal Genetics Reports*: Vol. 1, Article 14. <https://doi.org/10.4148/1941-4765.1028>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## Growth of a histidina strain, hist-2 (C94), of *Neurospora crassa*

### Abstract

Growth of a histidina strain, *hist-2* (C94), of *Neurospora crassa*

McNelly, Caroline. Growth of a histidine strain, hist-2 (C94), of Neurospora crassa.

wild-type. Using a conidial suspension as inoculum, the following media were tested:- minimal reproductive (Westergaard and Mitchell, 1945), pH 6.7, with histidine and nicotinamide added in a ratio 1:1 and 2:1; malt, peptone and yeast medium (Frost, this Newsletter) pH 6.5, with histidine and nicotinamide added in a 1:1 ratio; and complete malt medium, pH 6.5, supplemented with histidine only. All showed sparse growth and slight conidiation. Bacto-peptone medium (Frost, this Newsletter), pH 5.0, with no supplement and with added histidine: nicotinamide ratios of 1:0, 1:1, 1:2, and 2:1, also showed poor growth. However, this medium adjusted to pH 6.5 - 6.7 showed wild-type growth and conidiation in all cases where histidine had been added. The same results were found for ascospore germination from the cross cr. +, + (F945) x +, hist-2, nic-2 (C94, 43002). It has been found that bacto-peptone from different sources varies widely and the ratio of histidine to nicotinamide added must be adjusted to suit the batch of peptone used. In all cases, however, wild-type growth and conidiation is achieved only at pH 6.5 - 7.0.

The abundance of growth and conidiation of the double mutant, hist-2, nic-2 (C94, 43002), was found to decrease as the strain was successively backcrossed to either Lindegren or Abbott