

The genetic production of small, homokaryotic spores in *Neurospora tetrasperma*

J. W. Johnson

A. M. Srb

Follow this and additional works at: <http://newprairiepress.org/fgr>

Recommended Citation

Johnson, J. W., and A.M. Srb (1962) "The genetic production of small, homokaryotic spores in *Neurospora tetrasperma*," *Fungal Genetics Reports*: Vol. 2, Article 7. <https://doi.org/10.4148/1941-4765.1046>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

The genetic production of small, homokaryotic spores in *Neurospora tetrasperma*

Abstract

The genetic production of small, homokaryotic spores in *Neurospora tetrasperma*

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Johnson, Jane W. and Adrian M. Srb

The genetic production of small, homokaryotic spores in Neurospora tetrasperma.

When the colonial mutant pk-2 of Neurospora crassa is crossed into N. tetrasperma, small, homokaryotic ascospores are formed with high frequency. These smaller, homokaryotic spores are easily distinguishable from the large, heterokaryotic, bisexual spores normally formed when N. tetrasperma reproduces sexually. Isolation of the exceptional spores provides a population genetically comparable to random ascospore isolates after crosses in N. crassa or N. sitophila. Consequently, certain kinds of genetic work with N. tetrasperma are greatly facilitated if one parent in a cross includes the peak gene.

Table I shows the frequency of asci containing small, homokaryotic spores in a series of 12th and 13th generation backcrosses of pk to wild-type N. tetrasperma.

Table 1

<u>Parents</u>	<u>No. of asci observed</u>	<u>% of asci with more than 4 spores</u>	<u>Small spores as % of total spores</u>
1. +, A x +, a	1,165	0.86	0.43
2. +, A x <u>pk</u> , a	1,013	21.72	14.62
3. +, a x <u>pk</u> , A	918	32.90	22.81
4. +, +, A x <u>pk</u> , <u>ad</u> , a	1,011	49.55	41.00
5. +, +, A x +, <u>ad</u> , a	214	0.46	0.23

In the crosses designated 4 and 5, one of the parents includes a purple-adenine requiring gene originating in N. crassa and the mating type allele a from N. sitophila. The marker genes from N. crassa were first transferred to N. sitophila by an interspecific cross followed by back-crossing, and were then crossed into N. tetrasperma. ---Department of Plant Breeding, Cornell University, Ithaca, New York.