

## A versatile set of Lifeact-RFP expression plasmids for live-cell imaging of F-actin in filamentous fungi

Alexander Lichius  
*University of Edinburgh*

Nick D. Read  
*University of Edinburgh*

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Lichius, A., and N.D. Read (2010) "A versatile set of Lifeact-RFP expression plasmids for live-cell imaging of F-actin in filamentous fungi," *Fungal Genetics Reports*: Vol. 57, Article 4. <https://doi.org/10.4148/1941-4765.1070>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## A versatile set of Lifeact-RFP expression plasmids for live-cell imaging of F-actin in filamentous fungi

### Abstract

Here we report the construction and application of a range of expression plasmids designed to facilitate live-cell imaging of F-actin dynamics in filamentous fungi simultaneously with other, preferably GFP-tagged fusion proteins. Pros and cons of the use of three different red fluorescent proteins (RFPs), two different promoters and three different selection markers are addressed.

## A versatile set of Lifeact-RFP expression plasmids for live-cell imaging of F-actin in filamentous fungi

Alexander Lichius and Nick D. Read

Fungal Cell Biology Group, Institute of Cell Biology, University of Edinburgh, Rutherford Building, Edinburgh EH9 3JR, UK;

[Alex@fungalculture.org](mailto:Alex@fungalculture.org)

Fungal Genetics Reports 57:8-14


Here we report the construction and application of a range of expression plasmids designed to facilitate live-cell imaging of F-actin dynamics in filamentous fungi simultaneously with other, preferably GFP-tagged fusion proteins. Pros and cons of the use of three different red fluorescent proteins (RFPs), two different promoters and three different selection markers are addressed.

### Live-cell imaging of F-actin dynamics is possible in a wide range of eukaryotes.

Lifeact is a 17 aa peptide derived from the actin-binding protein 140 (Abp140) of *S. cerevisiae* which specifically binds to filamentous actin (F-actin) (Riedl et al., 2008). Functionality of green fluorescent Lifeact reporters (Lifeact-GFP) for the visualisation of F-actin structures in living cells has so far been documented for three of the four eukaryotic phyla, including yeasts and filamentous fungi (Berepiki et al., 2010; Böhmer et al., 2009; Coffman et al., 2009; Delgado-Álvarez et al., 2010; Riedl et al., 2008), plants (Era et al., 2009; Smertenko et al., 2010; Vidali et al., 2009), and mammals (Estecha et al., 2009; Riedl et al., 2008; Riedl et al., 2010). The application of Lifeact-GFP in filamentous fungi has been pioneered in the ascomycete *Neurospora crassa* (Berepiki et al., 2010; Delgado-Álvarez et al., 2010), and is currently being adopted for the use in numerous other fungal species. In basidiomycetes, however, labelling properties of Lifeact appear to be restricted to the visualization of septal rings, as F-actin cables and patches have so far not been explicitly reported (Böhmer et al., 2009). Lifeact-GFP reporters work equally well in fungi that use the CUG codon for serine and not leucine, such as *Candida sp.* (Kawaguchi et al., 1989), once they have been codon corrected, (E. Epp, McGill Univ. Montreal, pers.comm.).

### Why RFP and which one?


The most widely used fluorescent reporter for live-cell imaging analyses of protein dynamics in filamentous fungi is GFP. To allow simultaneous observation of F-actin with any other GFP-tagged protein, the development of a Lifeact probe (Riedl et al., 2008) labelled with a different color was desired. As green and red emission signals can be spectrally well separated, and fluorescence microscopes with the corresponding excitation and emission detection settings are widely available, we aimed to generate a functional Lifeact-RFP reporter construct. The first red fluorescent Lifeact probe that we adopted for the filamentous model fungus *Neurospora crassa* was Lifeact-tdTomato (Roca et al., 2010). Although Lifeact-tdTomato reliably labelled F-actin patches, cables and septal rings (Figure 1), this reporter occasionally produced additional ring-shaped structures (~1-2 µm in diameter, Figures 1B and C), which we suggest may be formed through spontaneous self-assembly of F-actin, destabilized through the interaction with the Lifeact-tdTomato construct, and native septins. Evidence for this comes from previous studies indicating that due to sterical hindrance, mRFP and tdTomato fusion constructs failed to correctly localize microtubules and connexin (Shaner et al., 2004; Shaner et al., 2008). This not well understood property of some RFPs might have potentiated mislocalization artefacts of the overexpressed Lifeact-tdTomato fusion construct leading to the formation of these cytoplasmic rings from destabilized F-actin. Interestingly, similar structures have been reported to form upon Latrunculin A treatment in mammalian cells (Kinoshita et al., 2002) and haploid cells of *Ustilago maydis* (Böhmer et al., 2009). Appearance of these rings in *N. crassa* was resolved by exchanging tdTomato for the monomeric red fluorescent protein TagRFP (Shaner et al., 2008).



**Figure 1.** Lifeact-RFPs labelled F-actin cables, patches and rings. (A) Two fused wt cells expressing Lifeact-TagRFP. F-actin cables stretched throughout the spore body of the upper cell, whilst actin patches were distributed over the surface of the bottom cell. Dense arrays of actin cables were localized at the cell fusion site (arrowhead), and condensed into an actin ring prior to septum formation (arrow). (B to D) Examples of wt conidia expressing Lifeact-tdTomato. (B) Conidial germling with actin cables aligned with the long axis of the germ tube. An extremely bright fluorescent ring (arrow) in the cytoplasm is obvious but not associated

with any known cellular structure. Similar rings have been described in other model systems and are likely the product of spontaneous self-assembly of the Lifeact-tdTomato construct with native septins (see text for details). Also vacuolar accumulation of the fusion construct created minor artefactual background fluorescence. (C) Two fusing germlings with bright fluorescent rings at different localizations within the cells. (D) Conidia in the process of CAT homing (Read et al., 2009), showing intense accumulation of actin patches and cables in the tip of the smaller germling (arrowhead). Formation of a cortical actin ring at the spore neck is again indicated with an arrow. All images show maximum intensity projection of deconvolved z-stacks. Scale bars, 5 µm.

To ensure an equivalent high level of fluorescence regardless of its use as a N- or C-terminal fusion tag the original TagRFP was modified through the addition of GFP-like N- and C-terminal residues (VSKGEE and LNGMDELYK, respectively), thereby improving its utility for 4D live-cell imaging (Berepiki *et al.*, 2010; Shaner *et al.*, 2004; Shaner *et al.*, 2008). Finally, the photostability of this optimized TagRFP version was increased through site-directed mutagenesis (SDM) of S162T, using the oligonucleotides TagRFP\_S162T\_fw 5'-CCTGGAAGGCAGAACCGGACATGGCCCTGAAGC-3' and its reverse complement TagRFP\_S162T\_rv 5'-GCTTCAGGGCCATGTCGGTTCTGCCTTCCAGG-3' (the mismatching nucleotide conferring S162T exchange is underlined; the affected triplet is in bold and italicised). The resulting TagRFP-T displayed a 10-times lower average decay constant ( $k$ ) compared to TagRFP in living cells of *N. crassa* (Figure 2), confirming its status as the most photostable monomeric FP so far described (Shaner *et al.*, 2008). Furthermore, TagRFP-T, in contrast to TagRFP, although showing a small, ~17% decrease in brightness, shows virtually no emission in the green part of the spectrum (Shaner *et al.*, 2008). Therefore, Lifeact-TagRFP-T is ideally suited for co-imaging with GFP-tagged proteins, for example BML-sGFP which reliably labels  $\beta$ -tubulin in microtubules (Berepiki *et al.*, 2010; Freitag *et al.*, 2004).



**Figure 2. Lifeact-TagRFP-T is on average 10x more photostable than Lifeact-TagRFP.** (A) Cells expressing Lifeact-TagRFP or Lifeact-TagRFP-T were imaged continuously over a period of 2 min with 1 frame/sec in one focal plane and 0.5 sec exposure using 60% illumination intensity from a 550 nm LED excitation light source. Bleaching rates were measured as the decrease in grey levels per second in 20  $\mu\text{m}^2$  regions of interest drawn up within three different intracellular areas containing different amounts of F-actin as judged by initial Lifeact-FP signal intensity: cortical F-actin at germ tube tip, cortical F-actin in non-polarized cell, and cytoplasmic F-actin in the cell centre of a polarized growing germling. Graphs represent the means of measurements from five individual cells. (B) On average, Lifeact-TagRFP lost 3.6 grey levels/sec, whereas Lifeact-TagRFP-T only bleached with 0.33 grey levels/sec, corresponding to average decay constants ( $k$ ) of  $0.003 \pm 0.0007$  and  $0.0004 \pm 0.0002$ , respectively

Although no mislocalization of a range of different TagRFP/RFP-T fusion constructs has been reported in mammalian cells (Shaner *et al.*, 2008) - which we have confirmed for a number of other fusion constructs in *N. crassa*, including MAP kinases (OS-2-TagRFP-T) (A. Lichius and N. D. Read, in prep.) and Cdc42-Rac1-Interactive-Binding (CRIB-TagRFP-T) reporters (A. Lichius, A. Goryachev, M. D. Fricker, B. Obara and N. D. Read, in prep.) - C-terminal addition of TagRFP/RFP-T did occasionally result in increased vacuolar accumulation of the fluorescence signal in *N. crassa*. This did not occur with the corresponding GFP-tagged versions of these constructs, suggesting this phenomenon might be associated with the increased stability of TagRFP/RFP-T at low pH compared to other FPs (pKa: TagRFP < TagRFP-T < tdTomato < EGFP, Shaner *et al.*, 2008). In other words, the amount of vacuolar accumulation associated with the normal turn-over of the fluorescent reporter proteins might not be different between GFP and TagRFP/RFP-T tagged constructs, but due to the increased pH stability of RFPs their degradation progresses much slower and thus they are more readily detectable. This unwanted side effect can hamper live-cell imaging significantly as it decreases the signal-to-noise ratio. It can

be minimized by selecting transformants with low plasmid integration number, and through the use of a weaker promoter driving reporter expression (see below). The low binding affinity of the Lifeact peptide to F-actin and its high on/off rate (Riedl *et al.*, 2008) are the likely reasons why these unwanted secondary effects involving (1) aggregation of the reporter construct with endogenous proteins and (2) pronounced vacuolar accumulation, do generally not occur with the Lifeact-TagRFP/RFP-T probes. With our current level of understanding, for co-localization studies it seems advisable to use the Lifeact peptide with a TagRFP-T label and tag the other protein under investigation with GFP. This will ensure optimal performance of both fluorescent fusion constructs.

### Which promoter provides stable expression throughout the fungal lifecycle?

So far, the most commonly used promoter for ectopic expression of FP fusion constructs in *Neurospora crassa* is *Pccg-1* (Freitag *et al.*, 2004; McNally and Free, 1988). *Pccg-1* regulates expression of the clock-controlled gene 1 (*ccg-1*) in a very complex manner, involving at least circadian clock control, light control, metabolic control, and very probably developmental regulation (Arpaia *et al.*, 1995; Loros *et al.*, 1989). Overall, this results in very strong but not necessarily consistent expression throughout development. Heterologous expression under its regulation is induced by light (Arpaia *et al.*, 1995), and highest in conidia and aerial hyphae, including enveloping hyphae during protoperithecial development (A. Lichius and K.M. Lord, C. E. Jeffree and N. D. Read, in prep.). In conidial germlings we consistently observed a noticeable decrease in expression and thus recordable signal intensity after 6-7 h of incubation (about 5-6 h after cell symmetry breaking of the germinating conidium). We also found that *Pccg-1* did not allow satisfactory visualization of Lifeact-FPs in mature vegetative hyphae (Berepiki *et al.*, 2010). To fully overcome these problems and guarantee stable fluorescent signal levels throughout the whole life cycle of the fungus, the promoter of the translation elongation factor-1 (locus NCU02003.4) (*Ptef-1*) of *Neurospora crassa* was alternatively employed (Berepiki *et al.*, 2010). In summary, for studies during the first 6-7 h of conidial development, *Pccg-1* is a very well suited promoter for *N. crassa*, whereas expression under its control appears less consistent during the transition from the germling phase into mature hyphal growth. The use of *Ptef-1*, which gives overall weaker but more consistent expression throughout all developmental stages of *N. crassa* is an alternative worth considering.

### Which selection marker allows transformation of *Neurospora* gene deletion strains?

An interesting question to address is how deletion of specific genes, e.g. components of the cell polarity machinery, influence F-actin architecture and functionality. To facilitate utilization of the Lifeact-RFP expression plasmids in all gene deletion strains generated within the *Neurospora* Genome Project (<http://www.dartmouth.edu/~neurosporagenome/>), it was imperative to employ a selection marker other than hygromycin B, as the hygromycin B resistance gene *hph* has been used as a central part of the gene knock-out cassette. The two resistance genes exploited in this study were *bar*, conferring resistance towards ignite (also known as phosphinothricin or basta) (Pall and Brunelli, 1994), and *nat1*, conferring resistance towards nourseothricin (Kück and Hoff, 2006). Ignite resistance has been employed for the generation of  $\Delta mus-51$  and  $\Delta mus-52$  gene deletion host strains (Colot *et al.*, 2006), thus it cannot be used for the transformation of knock-out strains in which the  $\Delta mus$  deletion cassette has not been removed through backcrossing to the wild type. Furthermore, ignite selection in *Neurospora* is relatively leaky compared to other commonly used selection markers, especially when using the potentially impure chemical extracted from the commercially available “Harvest” herbicide (Bayer Crop Sciences Ltd., Monheim, Germany) (Hays and Selker, 2000; Metzenberger *et al.*, 2000). Therefore, pALx-Lifeact transformant selection with nourseothricin was the preferred option, as it is much more stringent than ignite *per se* and has no reported problems in conferring cross-resistance to other selection markers, such as hygromycin B (Kück and Hoff, 2006). Nourseothricin has been successfully used for the transformation of several yeast species including *Saccharomyces cerevisiae* (Goldstein and McCusker, 1999), *Schizosaccharomyces pombe* (Hentges *et al.*, 2005), *Candida albicans* (Shen *et al.*, 2005) and *Cryptococcus neoformans* (McDade and Cox, 2001), and more recently in an increasing number of filamentous fungal species, including *Acremonium chrysogenum* and *Sordaria macrospora* (Kück and Hoff, 2006), as well as *Neurospora crassa* (Maerz *et al.*, 2009; Maerz *et al.*, 2008).

### Construction of pALx-Lifeact expression vectors.

pAL2-Lifeact, the first vector generated within this project, was assembled by blunt-end ligation of the Lifeact-tdTomato coding sequence amplified from pJT580a into an *EcoRI* linearized backbone of pBARGRG-1 (Pall and Brunelli, 1994; Roca *et al.*, 2010). pAL3-Lifeact was assembled using InFusion PCR cloning (<http://www.clontech.com>) to recombine the insert encoding for Lifeact-TagRFP amplified from pJT603f (Jens Tilsner, University of Edinburgh) with the *EcoRV/BamHI* linearized pBARGRG-1 in a ligation-independent step (Berepiki *et al.*, 2010). Exchange of the ignite resistance cassette (*PtpC::bar*) for the nourseothricin resistance cassette (*PtpC::nat1*) was accomplished using the same technique to recombine the *PtpC::nat1* insert amplified from pG-Nat1 (Kück and Hoff, 2006) with *SpeI/XbaI* linearized pAL3-Lifeact, resulting in pAL4-Lifeact. TagRFP in pAL3-Lifeact and pAL4-Lifeact were modified to TagRFP-T by SDM as described above to yield pAL5-Lifeact and pAL6-Lifeact, respectively. Promoter replacement of *Pccg-1* with *Ptef-1* was performed again using InFusion PCR cloning, but this time integrating the *Ptef-1* insert amplified from pTEFG-2 (Berepiki *et al.*, 2010) into *BglII/BamHI* linearized pAL4-Lifeact to generate pAL10-Lifeact. In a final SDM step, pAL10-Lifeact was modified to yield pAL12-Lifeact, which expresses Lifeact-TagRFP-T under control of the *Ptef-1* promoter, and is selectable with nourseothricin. Table 1 provides an overview of all the plasmids generated; Figure 3 shows examples of the physical vector maps. The sequence of the 17 aa Lifeact reporter (GGVADLIKKFESISKEE) is the same in all constructs and identical to the originally published peptide derived from Abp140 of *S. cerevisiae* (Riedl *et al.*, 2008). Tables 2 and 3 show primers used for InFusion PCR cloning and sequencing of the recombined vectors, respectively.

**Table 1.** List of all pALx-Lifeact expression plasmids currently deposited at the FGSC.

plasmid	promoter	reporter peptide	fluorescent protein	terminator	selection gene/marker	FGSC #
pAL2-Lifeact	<i>Pccg-1</i>	Lifeact	tdTomato	<i>TtrpC</i>	<i>bar/ignite</i>	756
pAL3-Lifeact	<i>Pccg-1</i>	Lifeact	TagRFP	<i>TtrpC</i>	<i>bar/ignite</i>	757
pAL4-Lifeact	<i>Pccg-1</i>	Lifeact	TagRFP	<i>TtrpC</i>	<i>nat1/nourseothricin</i>	758
pAL5-Lifeact	<i>Pccg-1</i>	Lifeact	TagRFP-T	<i>TtrpC</i>	<i>bar/ignite</i>	759
pAL6-Lifeact	<i>Pccg-1</i>	Lifeact	TagRFP-T	<i>TtrpC</i>	<i>nat1/nourseothricin</i>	760
pAL10-Lifeact	<i>Ptef-1</i>	Lifeact	TagRFP	<i>TtrpC</i>	<i>nat1/nourseothricin</i>	761
pAL12-Lifeact	<i>Ptef-1</i>	Lifeact	TagRFP-T	<i>TtrpC</i>	<i>nat1/nourseothricin</i>	762

**Table 2.** List of InFusion PCR cloning primers. 15-18 nt overhangs are italicised.

name	5'-3' sequence
LA_if_BamHI_fw	<i>TTTCCTCGACGGATCC</i> ATGGGTGTCGCAGATTTGATCAAGA
TagRFP_if_EcoRV_rv	<i>ATCGATAAGCTTGATATCTT</i> ACTTGTACAGCTCGTCCATGCCA
PtpC-nat1_if_SpeI_fw	<i>GTTTAGATCCACTAGTCA</i> ACTGATATTGAAGGAGCATTTTTTGG
nat1_if_XbaI_rv	<i>CAGAAATGTGCTCTAGAT</i> CAGGGGCAGGGCATGCTCA
Ptef-1_if_BglII_fw	<i>GCGAGGCAGAGATCTCC</i> GTGACCACTGAACTACACTAGTCTAGAG
Ptef-1_if_BamHI_LA_rv	<i>GACACCCATGGATCC</i> GATATCTTTGACGGTTGATGTGCTG


**Table 3.** List of sequencing primers.

name	5'-3' sequence	binding site
pAL_seq_fw	GATTGCCGATTAAGGGAACCAAATG	end of <i>Pccg-1</i>
pAL_seq_rev	AACACCATTTGTCTCAACTCCGGAG	downstream of FP end
PtpC_seq_fw	AGCTAGCTTGGCTGCAGGTC	upstream of <i>PtpC</i> start
nat1_seq_rv	GGGTTATTGTCTCATGAGCGGA	downstream of <i>nat1</i> end
Tef1_seq_fw	GGTGCTTCACAACTCAGCAAATTCTC	downstream of <i>Ptef-1</i> start
TagRFP_seq_rv	CATGTGAAGCCCTCAGGGAA	inside TagRFP/TagRFP-T

**Lifeact-RFP reporters work in a range of ascomycete fungi.**

Besides *Neurospora crassa* (Roca *et al.*, 2010) and this report), red fluorescent Lifeact reporters have already been successfully expressed and visualized in other ascomycetes, including *Sordaria macrospora* (S. Poeggeler, Univ. Goettingen, pers. comm.), *Magnaporthe oryzae* (N.J. Talbot, Univ. Exeter, pers. comm.) and *Aspergillus nidulans* (B.D. Shaw, Texas A&M Univ., pers. comm.); either by direct or co-transformation of pALx-TagRFP/RFP-T vectors. However, severe overexpression of Lifeact-FP reporters can lead to significant impairment of cell development and colony morphology, and is a problem that currently needs to be resolved for some fungal models, such as *Ashbya gossypii* (A. Gladfelter, Dartmouth College, pers. comm.). Once suitable expression systems have been identified for those species, the Lifeact-RFP reporters described here should assist and facilitate further and comparative studies of F-actin dynamics in a broad range of filamentous fungi.

All plasmids and a selection of *N. crassa* strains co-/expressing the various Lifeact-FP reporters are available from the Fungal Genetics Stock Center (<http://www.fgsc.net>).



**Figure 3.** Physical maps of a selection of pALx-Lifeact plasmids. Sequences and maps of all vectors can be downloaded from the FGSC web site (<http://www.fgsc.net>). Unique restriction sites are indicated in red.

### Acknowledgements

Sincere thanks to the named individuals for sharing unpublished results and providing valuable feedback to the manuscript; especially to Amy Gladfelter and Elias Epp. This research was funded by a Biotechnological and Biological Sciences Research Council grant (grant # BB/E010741/1) to N.D.R.

### References

- Arpaia, G., Loros, J. J., Dunlap, J. C., Morelli, G., Macino, G., 1995. Light induction of the clock-controlled gene *ccg-1* is not transduced through the circadian clock in *Neurospora crassa*. *Molecular and General Genetics*. 247, 157-63.
- Berepiki, A., Lichius, A., Shoji, J.-y., Tilsner, J., Read, N. D., 2010. F-actin dynamics in *Neurospora crassa*. *Eukaryotic Cell*. 9, 547-57.
- Böhmer, C., Ripp, C., Bölker, M., 2009. The germinal centre kinase Don3 triggers the dynamic rearrangement of higher-order septin structures during cytokinesis in *Ustilago maydis*. *Molecular Microbiology*. 74, 1484-1496.
- Coffman, V. C., Nile, A. H., Lee, I. J., Liu, H., Wu, J.-Q., 2009. Roles of formin nodes and myosin motor activity in Mid1p-dependent contractile-ring assembly during fission yeast cytokinesis. *Molecular Biology of the Cell*. 20, 5195-5210.
- Colot, H. V., Park, G., Turner, G. E., Ringelberg, C., Crew, C. M., Litvinkova, L., Weiss, R. L., Borkovich, K. A., Dunlap, J. C., 2006.

A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors. Proceedings of the National Academy of Sciences of the USA. 103, 10352-10357.

Delgado-Álvarez, D. L., Callejas-Negrete, O. A., Gómez, N., Freitag, M., Roberson, R. W., Smith, L. G., Mouriño-Pérez, R. R., 2010. Visualization of F-actin localization and dynamics with live cell markers in *Neurospora crassa*. Fungal Genetics and Biology. 47, 573-586.

Era, A., Tominaga, M., Ebine, K., Awai, C., Saito, C., Ishizaki, K., Yamato, K. T., Kohchi, T., Nakano, A., Ueda, T., 2009. Application of Lifeact reveals F-actin dynamics in *Arabidopsis thaliana* and the liverwort, *Marchantia polymorpha*. Plant and Cell Physiology. 50, 1041-1048.

Estecha, A., Sanchez-Martin, L., Puig-Kroger, A., Bartolome, R. A., Teixido, J., Samaniego, R., Sanchez-Mateos, P., 2009. Moesin orchestrates cortical polarity of melanoma tumour cells to initiate 3D invasion. Journal of Cell Science. 122, 3492-501.

Freitag, M., Hickey, P. C., Raju, N. B., Selker, E. U., Read, N. D., 2004. GFP as a tool to analyze the organization, dynamics and function of nuclei and microtubules in *Neurospora crassa*. Fungal Genetics and Biology. 41, 897-910.

Goldstein, A. L., McCusker, J. H., 1999. Three new dominant drug resistance cassettes for gene disruption in *Saccharomyces cerevisiae*. Yeast. 15, 1541-53.

Hays, S., Selker, E. U., 2000. Making the selectable marker *bar* tighter and more economical. Fungal Genetics Newsletter. 47, 107.

Hentges, P., Van Driessche, B., Tafforeau, L., Vandenhoute, J., Carr, A. M., 2005. Three novel antibiotic marker cassettes for gene disruption and marker switching in *Schizosaccharomyces pombe*. Yeast. 22, 1013-9.

Kawaguchi, Y., Honda, H., Taniguchi-Morimura, J., Iwasaki, S., 1989. The codon CUG is read as serine in an asporogenic yeast *Candida cylindracea*. Nature. 341, 164-166.

Kinoshita, M., Field, C. M., Coughlin, M. L., Straight, A. F., Mitchison, T. J., 2002. Self- and actin-templated assembly of mammalian septins. Developmental Cell. 3, 791-802.

Kück, U., Hoff, B., 2006. Application of the nourseothricin acetyltransferase gene (*nat1*) as dominant marker for transformation of filamentous fungi. Fungal Genetics Newsletter. 53, 9-11.

Loros, J. J., Denome, S. A., Dunlap, J. C., 1989. Molecular cloning of genes under control of the circadian clock in *Neurospora*. Science. 243, 385-388.

Maerz, S., Dettmann, A., Ziv, C., Liu, Y., Valerius, O., Yarden, O., Seiler, S., 2009. Two NDR kinase-MOB complexes function as distinct modules during septum formation and tip extension in *Neurospora crassa*. Molecular Microbiology. 74, 707-723.

Maerz, S., Ziv, C., Vogt, N., Helmstaedt, K., Cohen, N., Gorovits, R., Yarden, O., Seiler, S., 2008. The nuclear Dbf2-related kinase COT1 and the mitogen-activated protein kinases MAK1 and MAK2 genetically interact to regulate filamentous growth, hyphal fusion and sexual development in *Neurospora crassa*. Genetics. 179, 1313-1325.

McDade, H. C., Cox, G. M., 2001. A new dominant selectable marker for use in *Cryptococcus neoformans*. Medical Mycology. 39, 151-4.

McNally, M. T., Free, S. J., 1988. Isolation and characterization of a *Neurospora* glucose-repressible gene. Current Genetics. 14, 545-551.

Metzenberger, R. L., Jacobson, D. J., Bertrand, H., 2000. Making the selective agent for the *Bar* plasmids, phosphinothricin (glufosinate) affordable for routine use. Fungal Genetics Newsletter. 47, 79-80.

Pall, M. L., Brunelli, J. P., 1994. New plasmid and plasmid hybrid vectors and a *Neurospora crassa* genomic library containing the *bar* selectable marker and Cre/lox site-specific recombination system for use in filamentous fungi. Fungal Genetics Newsletter. 41, 63-65.

Read, N. D., Lichius, A., Shoji, J., Goryachev, A. B., 2009. Self-signalling and self-fusion in filamentous fungi. Current Opinion in Microbiology. 12, 608-615.

- Riedl, J., Crevenna, A. H., Kessenbrock, K., Yu, J. H., Neukirchen, D., Bista, M., Bradke, F., Jenne, D., Holak, T. A., Werb, Z., Sixt, M., Wedlich-Söldner, R., 2008. Lifeact: a versatile marker to visualize F-actin. *Nature Methods*. 5, 605-607.
- Riedl, J., Flynn, K. C., Raducanu, A., Gartner, F., Beck, G., Bosl, M., Bradke, F., Massberg, S., Aszodi, A., Sixt, M., Wedlich-Soldner, R., 2010. Lifeact mice for studying F-actin dynamics. *Nature Methods*. 7, 168-9.
- Roca, M. G., Kuo, H.-C., Lichius, A., Freitag, M., Read, N. D., 2010. Nuclear dynamics, mitosis and the cytoskeleton during the early stages of colony initiation in *Neurospora crassa*. *Eukaryotic Cell*. 9, 1171-1183.
- Shaner, N. C., Campbell, R. E., Steinbach, P. A., Giepmans, B. N. G., Palmer, A. E., Tsien, R. Y., 2004. Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein. *Nature Biotechnology*. 22, 1567-1572.
- Shaner, N. C., Lin, M. Z., McKeown, M. R., Steinbach, P. A., Hazelwood, K. L., Davidson, M. W., Tsien, R. Y., 2008. Improving the photostability of bright monomeric orange and red fluorescent proteins. *Nature Methods*. 5, 545-551.
- Shen, J., Guo, W., Kohler, J. R., 2005. CaNAT1, a heterologous dominant selectable marker for transformation of *Candida albicans* and other pathogenic *Candida* species. *Infection and Immunity*. 73, 1239-1242.
- Smertenko, A. P., Deeks, M. J., Hussey, P. J., 2010. Strategies of actin reorganisation in plant cells. *Journal of Cell Science*. 123, 3019-28.
- Vidali, L., Rounds, C. M., Hepler, P. K., Bezanilla, M., 2009. Lifeact-mEGFP reveals a dynamic apical F-actin network in tip growing plant cells. *PLoS ONE*. 4, e5744.