January 2015

Effects of Amino Acid and Energy Intake During Late Gestation on Piglet Birth Weight and Reproductive Performance of Gilts and Sows Housed Under Commercial Conditions

M. A. Goncalves

Kansas State University, Manhattan, marcio@k-state.edu

K. M. Gourley

Kansas State University, Manhattan, kgourley@k-state.edu

S. S. Dritz

Kansas State University, Manhattan, dritz@k-state.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

next page for additional authors

Part of the [Large or Food Animal and Equine Medicine Commons](#), and the [Other Animal Sciences Commons](#)

Recommended Citation

Goncalves, M. A.; Gourley, K. M.; Dritz, S. S.; Tokach, M. D.; Bello, N. M.; DeRouchey, J. M.; Woodworth, J. C.; and Goodband, R. D. (2015) "Effects of Amino Acid and Energy Intake During Late Gestation on Piglet Birth Weight and Reproductive Performance of Gilts and Sows Housed Under Commercial Conditions," *Kansas Agricultural Experiment Station Research Reports*: Vol. 1: Iss. 7. <https://doi.org/10.4148/2378-5977.1112>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2015 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Effects of Amino Acid and Energy Intake During Late Gestation on Piglet Birth Weight and Reproductive Performance of Gilts and Sows Housed Under Commercial Conditions

Cover Page Footnote

Appreciation is expressed to Hord Livestock Company (Bucyrus, OH) for providing the animals and research facilities, and to Ryan Hughes, Jason Dunfee, and Matt Davis for technical assistance.

Authors

M. A. Goncalves, K. M. Gourley, S. S. Dritz, M. D. Tokach, N. M. Bello, J. M. DeRouchey, J. C. Woodworth, and R. D. Goodband

Effects of Amino Acid and Energy Intake During Late Gestation on Piglet Birth Weight and Reproductive Performance of Gilts and Sows Housed Under Commercial Conditions¹

M. A. D. Gonçalves², K. M. Gourley, S. S. Dritz², M. D. Tokach, N. M. Bello³, J. M. DeRouchey, J. C. Woodworth, and R. D. Goodband

Summary

The objective of this study was to determine the effects of amino acid (AA) and energy intake during late gestation on piglet birth weight and reproductive performance of high-performing gilts and sows housed under commercial conditions. At d 90 of gestation, a total of 1,102 females (PIC 1050) were housed in pens by parity group (P1 or P2+), blocked by weight within each pen, and each female was randomly assigned to dietary treatments within weight block. Dietary treatments consisted of combinations of 2 standardized ileal digestible (SID) AA (10.7 or 20.0 g SID Lys intake/d with other AA meeting or exceeding the NRC [2012] recommendations as a ratio to Lys) and 2 energy intakes (4.50 or 6.75 Mcal/d intake of NE) in a 2 × 2 factorial arrangement. Data were analyzed using generalized linear mixed models with parity group and dietary treatments as the linear predictor and random effects of pen as the experimental unit for parity and the individual female as the experimental unit for dietary treatments. With high energy intake, the magnitude of BW gain during late gestation was greater (AA × Energy, $P < 0.001$) with increasing AA intake compared with increasing AA at low energy intake. Gilts gained more weight at low energy intake than sows (parity × energy, $P < 0.001$); however, there was no evidence for differences ($P = 0.601$) in weight gain between gilts and sows at high energy intake. Sows fed high-energy intake had marginally reduced probability of piglets born alive (parity × energy, $P = 0.092$) compared with sows fed low energy, but no evidence for differences in gilts was observed. This was due to the increased probability (parity × energy, $P = 0.014$) of still-borns to be higher in sows fed high energy intake. There was no evidence for differences between the dietary treatments for litter birth weight and individual piglet birth weight of total piglets born. However, individual born-live birth weight was heavier ($P =$

¹ Appreciation is expressed to Hord Livestock Company (Bucyrus, OH) for providing the animals and research facilities, and to Ryan Hughes, Jason Dunfee, and Matt Davis for technical assistance.

² Department of Diagnostic Medicine/Pathobiology, College of Veterinary Medicine, Kansas State University.

³ Department of Statistics, College of Arts and Sciences, Kansas State University.

0.011) for females fed high-energy intake treatments compared to those with low energy intake. Born-alive piglets from sows were heavier ($P < 0.001$) than those from gilts. There was a lower probability ($P = 0.034$) of pre-weaning mortality for females fed high AA intake compared to low AA intake, regardless of energy level. There was no evidence for differences between the dietary treatments on farrowing rate, number of total piglets born, and percent of piglets born alive in the subsequent cycle. In conclusion, 1) body weight gain of gilts and sows depends not only on energy but also AA intake, 2) sows fed an increased amount of energy had increased stillborn rates, 3) the positive effect of increased amount of feed during late gestation on individual piglet birth weight, 30 g per pig, was due to energy rather than AA intake.

Key words: amino acids, birth weight, energy, gilt, sow

Introduction

The increase in litter size over recent decades has reduced the uterine space available for fetal growth and development, thus reducing individual piglet birth weight. Consequently, lower birth weight reduces piglet survivability, wean weight, and market weight. However, few nutritional options have been identified to mitigate reduced birth weight. There is evidence that increasing feed intake in late gestation improves piglet birth weight (Cromwell et al., 1989⁴, Shelton et al., 2009⁵, Soto et al., 2011⁶). However, the effect of increased amounts of feed offered during late gestation on piglet birth weight in high-performing herds (> 14.5 total piglets born/sow) is unclear. Further, because all studies reported increased feed intake, which increased both energy and amino acids (AA), it is unclear if the influences on piglet birth weight are due to dietary AA or energy content. The objective of this study was to determine the effects of AA and energy intake during late gestation on piglet birth weight and reproductive performance of high-performing gilts and sows housed under commercial conditions. The hypothesis was that both maternal dietary AA and energy in the late gestation period would affect piglet birth weight.

Procedures

The Kansas State University Institutional Animal Care and Use Committee approved the protocol used in this experiment. The experiment was conducted at a commercial sow farm in northern Ohio during the summer. Females were individually housed from d 0 to 40 of gestation, and then were group-housed (20 and 18 sq. ft/animal for gilts and sows, respectively) from d 40 to 111 of gestation. Each group pen was equipped with an electronic sow feeding station (Schauer, Prambachkirchen, Austria) and two cup waterers. All females had ad libitum access to water.

From d 0 to 89 of gestation, females were fed a common diet with 0.59% standardized ileal digestible (SID) Lys according to body condition: Thin, ideal, and fat females were

⁴ Cromwell, G. L., D. D. Hall, A. J. Clawson, G. E. Combs, D. A. Knabe, C. V. Maxwell, P. R. Noland, D. E. Orr, and T. J. Prince. 1989. Effects of Additional Feed during Late Gestation on Reproductive Performance of Sows: A Cooperative Study. *J Anim Sci* 67:3–14.

⁵ Shelton, N. W., C. R. Neill, J. M. DeRouchey, M. D. Tokach, R. D. Goodband, J. L. Nelssen, and S. S. Dritz. 2009. Effects of increasing feeding level during late gestation on sow and litter performance. In: Kansas State University. Agricultural Experiment Station and Cooperative Extension Service. Swine Day 2009, SRP 1020:38–50.

⁶ Soto, J., L. Greiner, J. Connor, and G. Allee. 2011. Effects increasing feeding levels in sows during late gestation on piglet birth weights. *J. Anim. Sci.* 89:86 (Abstr.).

fed 7.0, 4.4, and 4.0 lb/d, respectively. To be eligible for enrollment in the experiment, at d 90 of gestation females must not have: 1) returned to estrus during the previous cycle, 2) had an abortion during the previous cycle, 3) moderate or greater severity lameness, or 4) body condition score less than 2. At d 90 of gestation, a total of 1,102 females (PIC 1050, Hendersonville, TN; 741 gilts and 361 sows) were housed in pens by parity group (P1 or P2+), blocked by weight within each pen, and each female was randomly assigned to dietary treatments within weight block. The average parity for P2+ (sows) after farrowing was 4.1 ± 1.9 cycles (mean $\pm$ SD). Dietary treatments consisted of combinations of 2 SID AA intakes (10.7 or 20.0g SID Lys/d and other AA met or exceeded the NRC [2012] recommendations as a ratio to Lys) and 2 energy intakes (4.50 or 6.75 Mcal/d intake of NE) in a 2 x 2 factorial arrangement. All other nutrients met or exceeded the NRC (2012) recommendations.

Two diets were formulated (Table 1) and delivered at 4 different ratios and intake levels to achieve the desired dietary treatments for the 2 x 2 factorial (Table 2). Diets were balanced on a Ca to standardized total tract digestible (STTD) P ratio basis. Phytase was included in both diets with release considered to be 0.12% for STTD P. No AA release was considered for phytase.

At d 111 of gestation, females were moved to the farrowing house and fed 8.0 lb/d of a common lactation diet with 1.25% SID Lys provided until farrowing and then provided the same diet ad libitum thereafter. Both gestation and lactation diets were corn-soybean meal-based and presented in meal form. After weighing individual piglets, litter size was equalized regardless of dietary treatment.

On the subsequent cycle, no dietary treatments were applied, and females were fed a common diet with 0.59% SID Lys according to body condition. Thin, ideal, and fat females were fed 7.0, 4.4, and 4.0 lb/d, respectively, until d 90 of gestation, and then feed allowance was increased by 2.0 lb/d for thin and ideal condition females.

Prior to diet formulation, 5 samples of corn and soybean meal were submitted for proximate and total AA analysis. The analyzed values were used in formulation in conjunction with NRC (2012) digestibility coefficients. Samples of the diets were submitted for analysis of DM, CF, ash, crude fat, Ca, and P (Ward Laboratories, Inc., Kearney, NE). Diet samples were taken from each electronic feeding station twice a week, and then CP and total AA analyses were conducted in duplicate on composite samples (Ajinomoto Heartland Inc., Chicago, IL). Feeding station calibration was monitored twice a week by weighing 10 samples from each dispenser in each station.

Data were analyzed using generalized linear mixed models with parity group and dietary treatments as the linear predictor and random effects of pen as the experimental unit for parity and the individual female as the experimental unit for dietary treatments. Models of non-normal responses were initially fitted with Laplace method of approximation to evaluate for overdispersion (Stroup, 2012)⁷. Then, models were fit using the restricted maximum likelihood estimation for inference. Response variables were fitted accordingly with distribution. Bonferroni adjustment was used for multiple compari-

⁷ Stroup, W. W. 2012. Generalized linear mixed models: modern concepts, methods and applications. CRC Press, Boca Raton, FL.

sons. Statistical models were fitted using the GLIMMIX procedure of SAS (SAS Institute Inc., Cary, NC). Results were considered significant at $P \leq 0.05$ and marginally significant at $0.05 < P \leq 0.10$.

Results and Discussion

Chemical analysis of feed was similar to formulated values (Table 3). In sows and gilts fed high-energy intake, the magnitude of BW gain was greater with increasing AA intake (Fig. 1A) compared to increasing Lys intake at low-energy intake (AA $\times$ energy, $P < 0.001$; Tables 4 and 5). Gilts gained more weight at low energy intake than sows (Fig. 1B; parity $\times$ energy, $P < 0.001$); however, there was no evidence for differences ($P = 0.601$) in weight gain between gilts and sows at high-energy intake.

There was no evidence of differences in the number of piglets total born between the dietary treatments. However, sows had an increased number ($P < 0.001$) of total piglets born, as compared with gilts. Sows fed the high-energy intake had marginally reduced probability of piglets born alive (parity $\times$ energy, $P = 0.001$), but there was no evidence for differences in gilts. This was due to the increased probability (parity $\times$ energy, $P = 0.014$) of stillborns to be greater in sows fed high-energy intake (Fig. 2). The probability of stillborns was reduced ($P = 0.049$) in females fed high AA intake. There was a AA $\times$ energy $\times$ parity ($P = 0.047$) interaction: Sows fed low-energy and high-AA intake had increased probability of mummified fetuses, as compared to sows fed low-energy but low-AA intake.

Considering the total number of piglets born, litter birth weight and individual piglet birth weight were heavier in sows ($P < 0.001$) than gilts, with marginally greater ($P = 0.091$) within-litter birth weight CV in females fed high energy compared to low energy, independent of parity. Within-litter birth weight, CV was greater ($P < 0.001$) in sows than gilts. However, there was no evidence for differences between the dietary treatments for litter birth weight and individual piglet birth weight of total piglets born.

When litter birth weight and individual birth weight for piglets born alive were considered, weights were heavier in sow litters ($P < 0.001$) compared to gilt litters. Individual born-alive birth weight (Fig. 3) was heavier ($P = 0.011$) for females fed the high-energy intake compared to low-energy intake fed females, regardless of AA intake or parity. Born-alive piglets from sows were heavier ($P < 0.001$) than those from gilts. There was no evidence for differences in the within litter birth weight CV between the dietary treatments. Within-litter birth weight CV was greater ($P < 0.001$) in sows than gilts.

Piglets suckling from females fed high AA intake showed a lower percentage ($P = 0.034$) of pre-weaning mortality (PWM) compared to those born to females fed low AA intake during late gestation, regardless of energy level.

There was no evidence of differences in lactation length between dietary treatment or parity. The percentage of females bred 7 d after weaning was greater ($P < 0.001$) for sows than for gilts due to shorter ($P < 0.001$) wean-to-estrus interval (WEI) in sows compared to gilts. However, there was no evidence for differences in percentage of females bred by 7 d after weaning or WEI between the dietary treatments.

There was no evidence for differences on subsequent farrowing rate between the dietary treatments or parity level. Sows had greater ($P < 0.001$) number of total piglets born compared to gilts on the subsequent cycle. However, gilts had increased ($P < 0.004$) probability of more piglets born alive compared to sows on the subsequent cycle. There was no evidence for differences between the dietary treatments on the number of total piglets born, and probability of born-alive piglets on the subsequent cycle. However, when stillborns were analyzed separately from mummified fetuses, females fed high energy had lower ($P = 0.040$) probability of stillborn piglets in the subsequent cycle compared to those fed low energy.

In conclusion, 1) body weight gain of gilts and sows depends not only on energy but also AA intake, 2) sows fed increased amount of energy had an increased stillborn rate, 3) pre-weaning mortality was reduced in piglets suckling from females with high AA intake, and 4) the positive effect of increased amount of feed during late gestation on individual piglet birth weight, 30 g per pig, was due to energy rather than AA intake.

Table 1. Diet composition (as-fed basis)¹

Ingredient	SID Lys, %	
	0.40	1.06
Corn	87.97	62.47
Soybean meal (46% CP)	8.06	33.59
Monocalcium phosphate	1.60	1.25
Limestone	1.50	1.45
Salt	0.50	0.50
L-Lysine HCl	0.045	0.045
DL-Methionine	---	0.200
L-Threonine	0.035	0.210
Choline chloride 60%	0.100	0.100
Vitamin/mineral premix	0.150	0.150
Phytase ²	0.035	0.035
Total	100	100
Calculated analysis		
Standardized ileal digestible (SID) AA, %		
Lys	0.40	1.06
Ile:lys	80	72
Leu:lys	219	144
Met:lys	41	47
Met & cys:lys	79	71
Thr:lys	81	80
Trp:lys	21	22
Val:lys	94	76
NE, kcal/lb	1,144	1,082
CP, %	11.2	21.5
Ca, %	0.85	0.86
P, %	0.62	0.66
Available P, %	0.52	0.48
Stand. Total Tract Dig. (STTD) P, %	0.52	0.52
Ca:STTD P	1.64	1.64

¹ Diets were fed from d 90 to 111 of gestation. Corn and soybean meal were analyzed for total AA content prior to diet formulation, and NRC (2012) SID digestibility values were used in the diet formulation.

² Quantum Blue 2G (AB Vista Feed Ingredients, Marlborough, UK) provided 318 FTU/lb of diet with a release of 0.12% STTD P.

Table 2. Experimental dietary treatment structure¹

Item	Amino acid intake:	Low		High	
	Energy intake:	Low	High	Low	High
	Delivered ratio ² , %:	71/29	100/0	0/100	50.5/49.5
Feed allowance, lb/d		3.99	5.91	4.17	6.06
Energy, Mcal NE/d		4.50	6.75	4.50	6.75
Lysine, g SID Lys/d		10.7	10.7	20.0	20.0

¹ Dietary treatment structure based on the two diets presented in Table 1.

² Delivered ratio between 0.40% and 1.06% SID Lys diets to achieve the desired dietary treatments on an intake/d basis.

Table 3. Chemical analysis of the diets (as-fed-basis)¹

	Amino acid intake:	Low		High	
	Energy intake:	Low	High	Low	High
Proximate analysis, %					
DM		89.3 (87.4)	89.3 (87.2)	90.1 (88.0)	89.8 (87.6)
CP		13.6 (14.2)	10.4 (11.2)	20.9 (21.5)	15.9 (16.3)
Crude fiber		1.6 (2.2)	1.6 (2.1)	2.1 (2.5)	1.9 (2.3)
Ca		0.84 (0.85)	0.74 (0.85)	0.79 (0.85)	0.78 (0.85)
P		0.59 (0.63)	0.59 (0.62)	0.64 (0.66)	0.64 (0.64)
Fat		2.6 (3.0)	2.5 (3.2)	2.8 (2.7)	2.5 (2.9)
Ash		4.5 (5.1)	4.0 (4.9)	5.1 (5.8)	4.5 (5.3)
Total amino acids, %					
Lys		0.66 (0.69)	0.48 (0.48)	1.14 (1.21)	0.81 (0.84)
Ile		0.50 (0.52)	0.38 (0.38)	0.84 (0.87)	0.61 (0.62)
Leu		1.16 (1.22)	0.96 (1.00)	1.67 (1.74)	1.32 (1.37)
Met		0.26 (0.29)	0.18 (0.19)	0.45 (0.54)	0.31 (0.36)
Met & cys		0.48 (0.52)	0.36 (0.38)	0.76 (0.85)	0.56 (0.61)
Thr		0.54 (0.56)	0.40 (0.39)	0.92 (0.98)	0.65 (0.68)
Trp		0.13 (0.14)	0.12 (0.10)	0.24 (0.26)	0.17 (0.18)
Val		0.59 (0.59)	0.47 (0.45)	0.90 (0.94)	0.69 (0.69)
His		0.34 (0.39)	0.26 (0.31)	0.52 (0.58)	0.38 (0.45)
Phe		0.63 (0.69)	0.49 (0.54)	1.00 (1.05)	0.75 (0.79)

¹ Diet samples were taken from each electronic feeding station twice a week, and then CP and total AA analyses were conducted in duplicate on composite samples by Ajinomoto Heartland Inc. (Chicago, IL).

² Values in parentheses indicate those calculated from diet formulation and are based on values from NRC, 2012 (Nutrient Requirements of Swine, 11th ed. Natl. Acad. Press, Washington, D.C.) with the exception of total AA content from corn and soybean meal, which were analyzed prior to diet formulation by Ajinomoto Heartland Inc. (Chicago, IL).

Table 4. Least square mean estimates (and corresponding SEM) for reproductive performance of gilts and sows fed different lysine and energy intake levels during late gestation under commercial conditions¹

Item	Net energy, Mcal/d:	Gilts				Sows				Probability, P< AA × Energy × Parity
		SID Lysine ² , g/d:	10.7	20.0	10.7	20.0	10.7	20.0	10.7	20.0
ADFI d 90 to 111, lb		4.0 ± 0.02	4.0 ± 0.02	5.7 ± 0.02	5.7 ± 0.02	4.0 ± 0.02	4.2 ± 0.02	6.0 ± 0.02	6.0 ± 0.02	0.608
BW d 90, lb		386.9 ± 3.33	385.1 ± 3.33	386.5 ± 3.35	386.5 ± 3.33	501.1 ± 4.87	501.3 ± 4.83	497.8 ± 4.85	504.2 ± 4.89	0.463
BW gain d 90 to d 111, lb		29.5 ± 0.97	36.2 ± 0.97	40.6 ± 0.97	54.0 ± 0.97	23.1 ± 1.48	29.5 ± 1.48	40.8 ± 1.50	50.7 ± 1.48	0.128
Total piglets born, n		14.2 ± 0.22	14.1 ± 0.21	14.1 ± 0.22	14.2 ± 0.21	15.3 ± 0.34	14.8 ± 0.33	15.1 ± 0.32	15.5 ± 0.35	0.249
Born alive, %		94.6 ± 0.5	95.0 ± 0.5	93.6 ± 0.5	94.2 ± 0.5	93.3 ± 0.8	93.1 ± 0.8	89.6 ± 1.0	90.8 ± 1.0	0.569
Mummified fetuses, %		1.8 ± 0.28	1.7 ± 0.27	2.6 ± 0.36	2.5 ± 0.34	1.6 ± 0.36	3.0 ± 0.54	3.4 ± 0.57	2.8 ± 0.54	0.047
Stillborn, %		3.5 ± 0.40	3.2 ± 0.38	3.6 ± 0.40	3.2 ± 0.37	5.1 ± 0.69	3.7 ± 0.58	6.9 ± 0.83	6.1 ± 0.79	0.456
Total born (TB)										
Litter birth weight, lb		39.7 ± 0.53	39.5 ± 0.53	39.2 ± 0.51	39.5 ± 0.51	45.6 ± 0.75	44.5 ± 0.75	45.4 ± 0.75	46.3 ± 0.77	0.453
Piglet birth weight, lb		2.76 ± 0.022	2.80 ± 0.022	2.82 ± 0.022	2.82 ± 0.022	3.00 ± 0.044	3.00 ± 0.044	3.04 ± 0.044	3.00 ± 0.044	0.885
Birth weight CV, %		20.2 ± 0.66	20.4 ± 0.65	21.7 ± 0.68	21.3 ± 0.67	25.6 ± 1.04	26.2 ± 1.05	27.3 ± 1.06	25.9 ± 1.07	0.610
Born alive										
Litter birth weight, lb		37.9 ± 0.51	37.9 ± 0.51	37.7 ± 0.51	38.1 ± 0.51	43.2 ± 0.75	43.0 ± 0.75	42.1 ± 0.75	44.1 ± 0.77	0.405
Piglet birth weight, lb		2.82 ± 0.022	2.82 ± 0.022	2.87 ± 0.022	2.89 ± 0.022	3.00 ± 0.044	3.06 ± 0.044	3.09 ± 0.044	3.11 ± 0.044	0.489
Birth weight CV, %		18.0 ± 0.47	18.2 ± 0.47	18.5 ± 0.48	18.3 ± 0.47	23.9 ± 0.77	23.2 ± 0.75	23.2 ± 0.76	21.9 ± 0.77	0.955
Litter size after equalization, n		14.5 ± 0.30	14.4 ± 0.30	14.6 ± 0.30	14.3 ± 0.30	14.2 ± 0.42	13.7 ± 0.40	13.9 ± 0.44	14.0 ± 0.43	0.462
Piglets weaned, %		84.3 ± 0.82	86.5 ± 0.77	86.2 ± 0.78	86.4 ± 0.78	80.7 ± 1.31	81.5 ± 1.26	82.0 ± 1.35	83.2 ± 1.26	0.365
Pre-weaning mortality (PWM), %		10.3 ± 0.69	8.0 ± 0.61	8.9 ± 0.64	8.4 ± 0.63	13.7 ± 1.15	13.1 ± 1.11	13.3 ± 1.21	12.1 ± 1.11	0.254
Piglet removal rate, %		5.0 ± 0.55	5.2 ± 0.57	4.5 ± 0.52	4.9 ± 0.55	5.1 ± 0.82	5.00 ± 0.79	4.3 ± 0.77	4.3 ± 0.75	0.963
Lactation length, d										
Lactation length, d		24.9 ± 0.27	24.9 ± 0.27	24.7 ± 0.46	24.0 ± 0.46	24.4 ± 0.41	24.1 ± 0.40	24.2 ± 0.72	24.2 ± 0.68	0.363
Wean-to-estrus interval, d		6.8 ± 0.43	5.9 ± 0.44	6.6 ± 0.45	6.2 ± 0.44	4.4 ± 0.71	4.2 ± 0.68	4.8 ± 0.77	4.9 ± 0.71	0.873
Females bred until 7 d after weaning, %		87.8 ± 2.88	88.9 ± 2.81	85.1 ± 3.26	89.1 ± 2.77	98.3 ± 1.72	98.3 ± 1.68	94.0 ± 3.37	96.2 ± 2.68	0.913

continued

Table 4. Least square mean estimates (and corresponding SEM) for reproductive performance of gilts and sows fed different lysine and energy intake levels during late gestation under commercial conditions¹

		Gilts				Sows				Probability, P< AA × Energy × Parity
SID Lysine ² , g/d:		10.7	20.0	10.7	20.0	10.7	20.0	10.7	20.0	
Item	Net energy, Mcal/d:	4.50	4.50	6.75	6.75	4.50	4.50	6.75	6.75	
Subsequent performance										
	Farrowing rate, %	88.3 ± 2.88	88.4 ± 2.91	84.2 ± 3.36	88.6 ± 2.86	93.7 ± 3.20	93.9 ± 3.13	91.3 ± 3.96	87.1 ± 4.67	0.436
	Total piglets born, n	13.2 ± 0.35	13.2 ± 0.35	13.0 ± 0.36	13.4 ± 0.35	14.7 ± 0.56	15.5 ± 0.56	15.5 ± 0.62	15.0 ± 0.59	0.208
	Born alive, %	93.9 ± 0.64	93.4 ± 0.68	93.9 ± 0.67	94.3 ± 0.61	91.1 ± 1.12	91.6 ± 1.07	92.2 ± 1.1	92.0 ± 1.11	0.459
	Mummified fetuses, %	1.8 ± 0.33	1.8 ± 0.34	2.4 ± 0.40	2.0 ± 0.35	1.7 ± 0.46	2.9 ± 0.60	3.1 ± 0.67	2.3 ± 0.57	0.250
	Stillborn, %	4.3 ± 0.54	4.7 ± 0.58	3.8 ± 0.52	3.7 ± 0.48	7.0 ± 1.01	5.5 ± 0.86	4.7 ± 0.84	5.5 ± 0.92	0.172

¹ A total of 1,105 females (PIC 1050) were used in a 21-d trial with 274 to 278 females per dietary treatment combination. Treatments were fed starting at d 90 of gestation and continued until d 111, at which time the sows were moved to the farrowing house and fed a common diet.

² Other AA met or exceeded the NRC (2012) recommendations as a ratio to Lys.

Table 5. Probability values of the effects of different lysine and energy intake levels during late gestation on reproductive performance of gilts and sows under commercial conditions¹

	AA × Energy	Parity × AA	Parity × Energy	Parity	AA	Energy
ADFI d 90 to 111, lb	0.834	0.050	0.707	0.001	0.001	0.001
BW d 90, lb	0.230	0.187	0.856	0.001	0.438	0.926
BW gain d 90 to d 111, lb	0.001	0.131	0.001	0.028	0.001	0.001
Total piglets born, n	0.154	0.938	0.492	0.001	0.901	0.552
Born alive, %	0.483	0.718	0.092	0.002	0.261	0.001
Mummified fetuses, %	0.068	0.134	0.910	0.199	0.461	0.001
Stillborn, %	0.628	0.471	0.014	0.001	0.049	0.013
Total born (TB)						
Litter birth weight, lb	0.189	0.795	0.241	0.001	0.904	0.489
Piglet birth weight, lb	0.546	0.446	0.643	0.001	0.993	0.365
Birth weight CV, %	0.266	0.792	0.533	0.001	0.678	0.091
Born alive						
Litter birth weight, lb	0.145	0.459	0.954	0.001	0.184	0.945
Piglet birth weight, lb	0.602	0.641	0.743	0.001	0.292	0.011
Birth weight CV, %	0.674	0.466	0.204	0.001	0.522	0.564
Litter size after equalization, n	0.761	0.987	0.986	0.103	0.516	0.904
Piglets weaned, %	0.516	0.789	0.781	0.001	0.120	0.087
Pre-weaning mortality (PWM), %	0.494	0.443	0.882	0.001	0.034	0.356
Piglet removal rate, %	0.804	0.653	0.670	0.724	0.830	0.155
Lactation length, d						
Lactation length, d	0.578	0.735	0.338	0.448	0.341	0.310
Wean-to-estrus interval, d	0.581	0.467	0.529	0.001	0.395	0.455
Females bred until 7 d after weaning, %	0.700	0.990	0.284	0.001	0.595	0.192
Subsequent performance						
Farrowing rate, %	0.927	0.456	0.428	0.167	0.981	0.163
Total piglets born, n	0.578	0.859	0.819	0.001	0.710	0.830
Born alive, %	0.808	0.893	0.875	0.004	0.904	0.284
Mummified fetuses, %	0.080	0.501	0.975	0.220	0.976	0.212
Stillborn, %	0.450	0.682	0.921	0.012	0.974	0.040

¹ A total of 1,105 females (PIC 1050) were used in a 21-d trial with 274 to 278 females per dietary treatment combination.

Figure 1A. Effects of different lysine and energy intake levels fed from d 90 to d 111 of gestation on BW gain of gilts and sows. Means with a different superscript within SID Lys level differ ($P < 0.05$).

Figure 1B. Effects of different energy intake levels fed from d 90 to d 111 of gestation on BW gain of gilts and sows. Means with a different superscript within parity level differ ($P < 0.05$).

Figure 2. Effects of different energy intake levels fed from d 90 to d 111 of gestation on stillborn rate of gilts and sows. Means with a different superscript within parity level differ ($P < 0.05$).

Figure 3. Effects of different energy intake fed from d 90 to d 111 of gestation on individual born live piglet birth weight of gilts and sows.