

January 2023

# The Indirect Effect of Smoking Level in the Association Between Urban Stress and Readiness to Quit Smoking among Adults Experiencing Homelessness

Carol Wang

*University of Texas MD Anderson Cancer Center, carolwangsf@gmail.com*

Tzu-An Chen

*University of Houston - Main, tchen3@Central.UH.EDU*

Maggie Britton

*University of Texas MD Anderson Cancer Center, MBritton@mdanderson.org*

*See next page for additional authors*

Follow this and additional works at: <https://newprairiepress.org/hbr>



This work is licensed under a [Creative Commons Attribution-Noncommercial 4.0 License](https://creativecommons.org/licenses/by-nc/4.0/)

## Recommended Citation

Wang, Carol; Chen, Tzu-An; Britton, Maggie; Businelle, Michael S. PhD; Kendzor, Darla E. PhD; McNeill, Lorna; and Reitzel, Lorraine R. (2023) "The Indirect Effect of Smoking Level in the Association Between Urban Stress and Readiness to Quit Smoking among Adults Experiencing Homelessness," *Health Behavior Research*: Vol. 6: No. 1. <https://doi.org/10.4148/2572-1836.1175>

This Research Brief is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Health Behavior Research by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

# The Indirect Effect of Smoking Level in the Association Between Urban Stress and Readiness to Quit Smoking among Adults Experiencing Homelessness

## Abstract

Over 70% of adults experiencing homelessness are cigarette smokers, a fivefold greater rate than in the general U.S. population. Consequently, tobacco-related conditions are the leading causes of disease and death for this group. Adults experiencing homelessness tend to seek shelter in urban areas. Thus, they not only experience the daily stressors of being homeless, but they may additionally experience unique or additive urban stressors (e.g., stress related to using public services, crime and violence, and/or cultural conflicts with others). For some smokers, stress is known to increase smoking rates and decrease readiness to quit smoking. Likewise, increased smoking rates alone may lead to a lower likelihood of making a quit attempt. The current study examined the potential mediating role of smoking level in the association of urban stress and quit readiness among adults experiencing homelessness (N = 411). Two multinomial logistic regression analyses revealed that urban stress was positively associated with smoking level ( $p = 0.02$ ). The odds ratio for one-unit increase in stress was 1.047 (CI<sub>95</sub>:1.014, 1.082) for being a heavy vs. non-daily smoker. Furthermore, analyses revealed smoking level mediated the effect of stress on quit readiness ( $ab = -0.005$ , CI<sub>95</sub>:-0.010, -0.002]). Homeless smokers who report high levels of stress might smoke at higher levels, which could attenuate quit readiness.

## Keywords

smoking, tobacco, homelessness, urban stress, smoking level

## Acknowledgements/Disclaimers/Disclosures

Research reported in this publication was supported in part by the TSET Health Promotion Research Center at the University of Oklahoma Health Sciences Center and the Oklahoma Tobacco Settlement Endowment Trust (092-016-0002) (to M.S.B.), with additional support from the University of Houston (to L.R.R.). The execution of this manuscript was also supported by the NIMHD (U54MD015946, on which L.R.R. was supported, and T.A.C. and M.S.B. are supported), by a NIMHD/NIDA supplement award (U54MD015946-03S1, on which M.B. is MPI of the science and on which L.R.R. is supported), and by NIDA (R25DA054015 to L.R.R. as MPI, on which T.A.C. was supported and M.B. functioned as an early career scholar). C.W. was supported by the Cancer Prevention Research Training Program (CP RTP) at MD Anderson Cancer Center and the Cancer Prevention and Research Institute of Texas (CPRIT) Postdoctoral Fellowship in Cancer Prevention (RP 170259, Drs. Shine Chang and Sanjay Shete, MPIs), with research mentorship provided by L.H.M. and L.R.R. Training and education grants at MD Anderson benefit from the Cancer Center Support grant, P30CA016672. The study was approved by relevant institutions and informed consent was obtained from all participants. The authors have no conflicts of interest to declare, financial or otherwise.

## Authors

Carol Wang, Tzu-An Chen, Maggie Britton, Michael S. Businelle PhD, Darla E. Kendzor PhD, Lorna McNeill, and Lorraine R. Reitzel

## **The Indirect Effect of Smoking Level in the Association Between Urban Stress and Readiness to Quit Smoking among Adults Experiencing Homelessness**

**Carol Wang, Ph.D.**  
**Tzuan A. Chen, Ph.D.**  
**Maggie Britton, Ph.D.**  
**Michael S. Businelle, Ph.D.**  
**Darla E. Kendzor, Ph.D.**  
**Lorna H. McNeill, Ph.D, M.P.H.**  
**Lorraine R. Reitzel, Ph.D., FAAHB, FSRNT\***

### **Abstract**

Over 70% of adults experiencing homelessness are cigarette smokers, a fivefold greater rate than in the general U.S. population. Consequently, tobacco-related conditions are the leading causes of disease and death for this group. Adults experiencing homelessness tend to seek shelter in urban areas. Thus, they not only experience the daily stressors of being homeless, but they may additionally experience unique or additive urban stressors (e.g., stress related to using public services, crime and violence, and/or cultural conflicts with others). For some smokers, stress is known to increase smoking rates and decrease readiness to quit smoking. Likewise, increased smoking rates alone may lead to a lower likelihood of making a quit attempt. The current study examined the potential mediating role of smoking level in the association of urban stress and quit readiness among adults experiencing homelessness ( $N = 411$ ). Two multinomial logistic regression analyses revealed that urban stress was positively associated with smoking level ( $p = 0.02$ ). The odds ratio for one-unit increase in stress was 1.047 (CI<sub>95</sub>:1.014, 1.082) for being a heavy vs. non-daily smoker. Furthermore, analyses revealed smoking level mediated the effect of stress on quit readiness ( $ab = -0.005$ , CI<sub>95</sub>:-0.010, -0.002]). Homeless smokers who report high levels of stress might smoke at higher levels, which could attenuate quit readiness.

\*Corresponding author can be reached at: [LReitzel@mdanderson.org](mailto:LReitzel@mdanderson.org)

### **Introduction**

Over 580,000 people reported experiencing homelessness in the United States on a single night in 2020, with over one quarter experiencing ongoing, chronic ( $\geq 1$  year) homelessness (HUD, 2021). As many as 70% of adults experiencing homelessness are conventional cigarette smokers, a prevalence 5 times higher than their housed counterparts (Baggett & Rigotti, 2010; Creamer et al., 2019); and tobacco-related

conditions have been some of the top leading causes of elevated morbidity and mortality rates within this group (Baggett, Hwang, et al., 2013; Soar et al., 2020; Vijayaraghavan, et al., 2020). For example, one study cited cancer as the second leading cause of death in a large sample of adults experiencing homelessness, with lung cancer constituting more than one third of cancer deaths (Baggett, Tobey, et al., 2013). Consequently, facilitating smoking cessation among adults experiencing homelessness is crucial to

reduce their smoking-related health/cancer disparities.

Cigarette smoking is linked with the chronic stress associated with experiencing homelessness (Businelle et al., 2013; Okuyemi, et al. 2006). Soar and colleagues' (2020) systematic review on smoking cessation among homeless smokers highlighted the importance of the social environment as a key barrier to smoking cessation. Common stressors experienced by this group include fear of victimization, food insecurity, and difficulty obtaining adequate shelter and accessing healthcare (Baggett, Hwang, et al., 2013; Baggett, Tobey, et al., 2013; Baggett & Rigotti, 2010). Moreover, there are potentially unique and additional stressors for those in urban areas (Lecic-Tosevski, 2019), including difficulty accessing public services, high crime and violence, exposure to gang activity, and elevated racism and discrimination (Agrawal et al., 2019; Murphy, 2019; Palepu et al., 2012; Wrighting et al., 2019).

Adults experiencing homelessness report smoking as a way of coping with acute stressors, including urban stress (Businelle et al., 2013; Okuyemi et al., 2006). In the same vein, previous studies have shown a strong link between stress and nicotine withdrawal (Lawless et al., 2015). More specifically, stress may increase the urge to smoke. As acute stressors may transition to the experience of chronic stress in adults experiencing homelessness, individuals who smoke may increase the number of cigarettes they smoke as a perceived solution to relieve the physiological consequences of stress dysregulation (i.e., smoking to “relieve stress”). A parallel to this is seen in research among domiciled smokers during the COVID-19 pandemic: sizeable portions of

smokers reporting high or increased stress during this time also reported increases in smoking rates (Nagawa et al., 2022; Popova et al., 2023; Rigotti et al., 2021), and for some, decreased readiness to quit (Klemperer, 2020). Therefore, it may be that for individuals experiencing homelessness, experiencing greater urban stress may translate to heavier cigarette consumption as a perceived means to cope, while also decreasing the likelihood of quitting (Brown et al., 2022; Lindson-Hawley et al., 2016).

Research on heaviness of smoking (smoking level) and how it links to factors related to quitting has been somewhat limited in homeless groups. Extant work suggests that light smoking is more prevalent among homeless vs. domiciled smokers, and that light homeless smokers report more past-year quit attempts compared to moderate/heavy homeless smokers (Nguyen et al., 2015). Furthermore, work done among domiciled smokers showed that non-daily smokers reported the highest desire to quit smoking, compared to moderate/heavy smokers (Savoy et al., 2014). Further, non-daily smokers were more likely to make a quit attempt within the past year and successfully quit, compared to heavy smokers (Swayampakala et al., 2013). Thus, indirect evidence may suggest that heavier smokers experience less readiness to quit than those who smoke fewer cigarettes per day.

Despite studies reporting associations between urban stress and heaviness of smoking (i.e., smoking level) on the one hand, and heaviness of smoking and less readiness to quit smoking on the other, it is *unclear* whether heaviness of smoking may mediate an association between urban stress and homeless smokers' readiness to quit smoking. However, it is *possible* given that

readiness to quit is a potential precursor to making a quit attempt and achieving smoking abstinence (Smit et al., 2010). A recent review indicated that much more research is needed regarding how to best support individuals experiencing homelessness who use tobacco, particularly in light of their daily stressors (Vijayaraghavan et al., 2020). This study was meant to incrementally add to this literature base. Results may have implications for targeted treatment efforts in homeless shelters where the capacity and resources to address smoking may be limited. It can also suggest the potential utility of reducing cigarette consumption to increase quit readiness. Finally, it may also suggest the importance of emphasizing alternative and positive coping strategies in interventions to reduce urban stress, which may ultimately better position smokers experiencing homelessness to make a smoking quit attempt.

## Methods

### Participants

Participants were 457 homeless adult conventional cigarette smokers who were a part of a larger study about the health of homeless adults (Neisler et al., 2018; Reitzel et al., 2017). Inclusion criteria for the parent study included being age 18 or over, currently receiving services from at least 1 of 6 homeless serving agencies in Oklahoma City, OK, and > 6<sup>th</sup> grade English literacy. Additional inclusion criteria for the current study were self-identifying as being currently homeless and having smoked a cigarette in the last 30 days.

### Procedures

Enrolled participants completed questionnaires onsite at a service agency and

were compensated with a \$20 gift card. Data were collected from July-August 2016. The study was approved by affiliated Institutional Review Boards and participants provided informed consent (see Neisler et al., 2018; Reitzel et al., 2017 for more information).

## Measures

**Participant characteristics.** Participant characteristics included sex, age, race, and self-reported history of severe mental illness (major depression, post-traumatic stress or other anxiety disorders, bipolar disorder, and/or schizophrenia; coded 0 = “No” or 1 = “Yes”).

**Urban life stress.** Urban life stress was measured using the 21-item Urban Life Stress Scale (ULSS; Jaffee et al., 2005), which assesses potential sources of daily stress experienced by persons living in medium to large cities (e.g., money or finances, transportation, neighborhood environment). Items, rated on a 5-point Likert scale with 1 = “No stress” and 5 = “Extreme stress,” were summed for a total score. The reliability was  $\alpha = 0.91$ .

**Smoking level.** Smoking level was self-reported via status as a non-daily smoker or a daily smoker, which was further categorized using the average cigarettes smoked per day, as follows: light (1-10 cigarettes/day), moderate (11-20 cigarettes/day), or high (21+ cigarettes /day) level smokers. The categorization of smoking level was based on previous work demonstrating clinically significant differences in tobacco related outcomes, including nicotine dependence, between the groups (e.g., Nguyen et al., 2015; Savoy et al., 2014).

**Readiness to quit.** Readiness to quit was assessed using a modified Biener & Abrams

(1991)'s Contemplation Ladder. This ordinal measure has 8 levels with anchors where 1 = "I enjoy smoking and have decided not to quit smoking for my lifetime. I have no interest in quitting," and 8 = "I still smoke, but I have begun to change, like cutting back on the number of cigarettes I smoke. I am ready to set a quit date."

### Statistical Analyses

Of the sample of 457 participants, missing data ranged from 0% to 8.32% with no patterns related to missingness (Little's MCAR test  $X^2 = 4.581, p = .101$ ). Therefore, the analyzable sample ( $n = 411$ ) included participants with complete data. Chi-square score tests were used to examine whether the proportional odds assumptions were met for readiness to quit (McCullagh, 1980) with a  $p > 0.05$ . However, the assumptions of proportional odds were not met; therefore, the effect of urban life stress on smoking level and readiness to quit, respectively, was examined using two multinomial logistic regression analyses. Odds ratio (ORs) and 95% confidence intervals were calculated for each interpretable result. Lastly, mediation analysis was conducted using structural equation modeling to test the indirect effect on urban life stress and readiness to quit via smoking level in a single analysis. Due to the ordinal nature of the mediator and outcome, we conducted Bayesian analyses with 10,000 iterations in line with an ordered probit model and non-normal parameter distributions (Muthén, 2011). The total effect, direct effect, and indirect effects were derived. Age, sex, race, and serious mental illness diagnosis were included as covariates in the multinomial logistic regression models and mediation analysis. Significance was set at  $p < 0.05$ . Mediation analysis was conducted with the Mplus statistical package (version

7.4); other analyses were run using SAS 9.4 (SAS Institute, Cary, NC, USA).

## Results

### Sample Descriptive Statistics and Correlations

Of the 411 participants ( $M_{\text{age}} = 43.3 \pm 11.8$ ), 64.96% ( $n = 267$ ) were men. Nine percent ( $n = 37$ ) of participants were non-daily smokers, 28.71% ( $n = 118$ ) were light smokers, 46.72% ( $n = 192$ ) were moderate smokers, and 15.57% ( $n = 64$ ) were heavy smokers. See Table 1 (participant characteristics) and Table 2 (correlations between variables).

### Multinomial Logistic Regression

Two adjusted multinomial logistic regression analyses were conducted to examine the effect of urban life stress on smoking level (reference group: non-daily smokers) and readiness to quit (reference group: "[...] I have no interest in quitting."). The overall effect of urban life stress was significant for smoking level ( $p = 0.0192$ ), but not for readiness to quit ( $p = 0.7336$ ). Significant results showed that the odds ratio for a one-unit increase in urban life stress was 1.047 ( $CI_{.95} = 1.014, 1.082$ ) for being a heavy smoker vs. non-daily smoker. The odds ratio was not significant for other smoking levels (see Appendix Table 1).

Table 1

*Sociodemographic Characteristics of Adults Experiencing Homelessness Who Smoke (n = 411)*

Variables	Mean (SD)	% [n]
Age (years)	43.3 (11.8)	
Male		64.9 [267]
Mental Health Diagnosis (yes)		65.5 [269]
Race		
Non-Hispanic White		59.9 [246]
Black/African American		17.8 [73]
Native American/Alaskan Native		9.7 [40]
Hispanic/Latino		2.7 [11]
Multiracial/Other		9.9 [41]
Urban Life Stress	49.1 (14.5)	
Smoking Level		
Non-daily		9 [37]
Light (1-10/day)		28.7 [118]
Moderate (11-20/day)		46.7 [192]
Heavy (21+/day)		15.6 [64]
Readiness to Quit		
I enjoy smoking and have decided not to quit smoking for my lifetime. I have no interest in quitting.		7.8 [32]
I never think about quitting smoking, and I have no plans to quit.		6.1 [25]
I rarely think about quitting smoking, and I have no plans to quit.		11.2 [46]
I sometimes think about quitting smoking, but I have no plans to quit.		18.5 [76]
I often think about quitting smoking, but I have no plans to quit.		20.9 [86]
I definitely plan to quit smoking within the next 6 months.		16.3 [67]
I definitely plan to quit smoking within the next 30 days.		5.8 [24]
I still smoke, but I have begun to change, like cutting back on the number of cigarettes I smoke. I am ready to set a quit date.		13.4 [55]

Table 2

*Correlations Between Urban Stress, Smoking Level, and Readiness to Quit Smoking in a Sample of Adults Experiencing Homelessness Who Smoke (n = 411)*

	2	3	4	5	6
1. Readiness to Quit	-0.25***	0.03	-0.04	0.01	-0.04
2. Smoking Level		0.15**	0.04	-0.06	0.08
3. Urban Life Stress			-0.01	0.11*	0.23***
4. Age				-0.08	0.06
5. Sex (Female)					0.16**
6. Mental Health Diagnosis (Yes)					

Note. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\*  $p < 0.001$




Figure 1. Mediation analysis. Note: SD = standard deviation; CI = confidence interval.

### Mediation Analysis

In adjusted Bayesian mediation analyses, urban life stress was again positively associated with smoking level ( $a = 0.011$ ; CI<sub>.95</sub>: 0.004, 0.018), and smoking level was negatively associated with readiness to quit ( $b = -0.358$ , CI<sub>.95</sub>: -0.484, -0.232; see Figure 1). Results indicated a statistically significant indirect effect ( $ab = -0.005$ , CI<sub>.95</sub>: -0.010, -0.002), but a non-significant direct effect ( $c' = 0.007$ , CI<sub>.95</sub>: -0.005, 0.019) and total effect ( $c = 0.001$ , CI<sub>.95</sub>: -0.011, 0.013).

### Discussion

This study explored how urban stress, smoking level, and quit readiness were associated within a sample of smokers experiencing homelessness. Results supported that urban life stress was indirectly associated with readiness to quit through smoking level, whereby greater urban stress was related to heavier smoking, which was associated with less readiness to quit. Although longitudinal work is required to determine temporal relations, these results appear to be consistent with previous work suggesting that smoking operates as a means

of coping with high levels of stress experienced by adults who are homeless (Businelle et al., 2013; Okuyemi et al., 2006), and that experiencing more urban stressors may be linked with heavier smoking habits, which, in turn, may dampen readiness to quit smoking.

If this pattern of results is replicated in future longitudinal work, there may be several treatment implications to reduce the health impact of smoking on adults experiencing homelessness. Firstly, the experience of urban life stress can be intervened upon directly through the provision of practical aides to reduce its impact on affected persons. This might be achieved through the provision of assistance in navigating public transportation, obtaining needed health care and other public services, or behavioral psychotherapy for anxiety by professionals or trained peers. Additionally, efforts in the community and within shelter settings to improve perceptions of safety, reduce criminal activity, and curb the violence experienced by this vulnerable group are needed to reduce urban stress. Secondly, education and intervention efforts for smokers experiencing homelessness can



emphasize the use of healthy alternative approaches to coping with experienced stress (as opposed to smoking and/or smoking more cigarettes). Likewise, teaching emotional regulation coping strategies may be necessary for handling unavoidable stressors.

Study limitations include a cross-sectional design and the use of convenience sampling. Longitudinal studies are needed to shed light on the long-term effects of urban life stressors and smoking level on readiness to quit across time among this group. Studies are also needed to examine behavioral outcomes (e.g., quit attempts, cessation). Moreover, dual or poly-tobacco use was not accounted for in the analyses. Furthermore, this study was conducted among homeless-serving agencies in Oklahoma City and may not be representative of other homeless smokers across the United States. Additionally, the sample only represents homeless adults utilizing services and excluded homeless adults who did not access services, who may have different health risk factors and/or outcomes than this sample. Finally, the failure to meet the proportional odds assumptions in the present study may suggest that an alternate measurement of readiness to quit with fewer categories might be used in future research so that ordinal logistic regression can be employed.

### **Implications for Health Behavior Research**

There may be implications for a stepped reduction in cigarette consumption as a strategy toward eventual abstinence (Lindson et al., 2019), which would require provider and patient acknowledgement that smoking fewer cigarettes is an important and significant goal in the journey toward eventual abstinence. Initial research indicates

that a reduction in combustible cigarettes smoked amongst smokers experiencing homeless may be achieved through harm reduction strategies that rely on nicotine replacement (e.g., e-cigarettes; Scheibein et al., 2020). Successes in reducing combustible cigarette consumption might be used to reduce overall nicotine dependence and build self-efficacy to quit. Furthermore, cognitive intervention strategies, like motivational interviewing, can be used to directly bolster quit readiness (Fiore et al., 2008). As intent to quit smoking is associated with making quit attempts and achieving long-term abstinence (Martínez et al., 2015; Smit et al., 2010), this should have downstream effects on increasing smoking cessation rates in this population. Although multi-level interventions incorporating attention to each of these factors would be desirable, treatment resources are often limited in settings that provide services to adults experiencing homelessness; therefore, targeting smokers most likely to successfully quit may be important. These results might suggest that lighter smokers should be prioritized for abstinence-promoting programs and may be more successful due to greater quit readiness, whereas heavier smokers might benefit most from an initial focus on strategies to cut down on cigarette intake while building quit motivation before entering a formal cessation program. More work is needed, however, to confirm the veracity of these suppositions.

### **Discussion Question(s)**

1. How might the findings from this study guide clinicians and trained peers to provide the most effective smoking cessation treatment for adults who smoke and are homeless?

### Acknowledgments

Research reported in this publication was supported in part by the TSET Health Promotion Research Center at the University of Oklahoma Health Sciences Center and the Oklahoma Tobacco Settlement Endowment Trust (092-016-0002) (to M.S.B.), with additional support from the University of Houston (to L.R.R.). The execution of this manuscript was also supported by the NIMHD (U54MD015946, on which L.R.R. was supported, and T.A.C. and M.S.B. are supported), by a NIMHD/NIDA supplement award (U54MD015946-03S1, on which M.B. is MPI of the science and on which L.R.R. is supported), and by NIDA (R25DA054015 to L.R.R. as MPI, on which T.A.C. was supported and M.B. functioned as an early career scholar). C.W. was supported by the Cancer Prevention Research Training Program (CPRTP) at MD Anderson Cancer Center and the Cancer Prevention and Research Institute of Texas (CPRIT) Postdoctoral Fellowship in Cancer Prevention (RP 170259, Drs. Shine Chang and Sanjay Shete, MPIs), with research mentorship provided by L.H.M. and L.R.R. Training and education grants at MD Anderson benefit from the Cancer Center Support grant, P30CA016672. The study was approved by relevant institutions and informed consent was obtained from all participants. The authors have no conflicts of interest to declare, financial or otherwise.

### References

Agrawal, P., Neisler, J., Businelle, M. S., Kendzor, D. E., Hernandez, D. C., Odoh, C., & Reitzel, L. R. (2019). Exposure to violence and sleep inadequacies among men and women living in shelter

setting. *Health Behavior Research*, 2(4), Article 19. <https://doi.org/10.4148/2572-1836.1051>

Baggett, T. P., Hwang, S. W., O'Connell, J. J., Porneala, B. C., Stringfellow, E. J., Orav, E. J., Singer, D. E., & Rigotti, N. A. (2013a). Mortality among homeless adults in Boston: Shifts in causes of death over a 15-year period. *JAMA Internal Medicine*, 173(3), 189-195.

<https://doi.org/10.1001/jamainternmed.2013.1604>

Baggett, T. P., & Rigotti, N. A. (2010). Cigarette smoking and advice to quit in a national sample of homeless adults. *American Journal of Preventive Medicine*, 39(2), 164-172.

<https://doi.org/10.1016/j.amepre.2010.03.024>

Baggett, T. P., Tobey, M. L., & Rigotti, N. A. (2013b). Tobacco use among homeless people—addressing the neglected addiction. *The New England Journal of Medicine*, 369(3), 201-204.

<https://doi.org/10.1056/NEJMp1301935>

Biener, L., & Abrams, D. B. (1991). The Contemplation Ladder: Validation of a measure of readiness to consider smoking cessation. *Health Psychology*, 10(5), 360-365.

<https://doi.org/10.1037//0278-6133.10.5.360>

Brown, H. A., Roberts, R. D., Chen, T. A., Businelle, M. S., Obasi, E. M., Kendzor, D. E., & Reitzel, L. R. (2022). Perceived disease risk of smoking, barriers to quitting, and cessation intervention preferences by sex amongst homeless

- adult concurrent tobacco product users and conventional cigarette-only users. *International Journal of Environmental Research and Public Health*, 19(6), Article 3629. <https://doi.org/10.3390/ijerph19063629>
- Businelle, M. S., Cuate, E. L., Kesh, A., Poonawalla, I. B., & Kendzor, D. E. (2013). Comparing homeless smokers to economically disadvantaged domiciled smokers. *American Journal of Public Health*, 103 Suppl 2(Suppl 2), S218-S220. <https://doi.org/10.2105/AJPH.2013.301336>
- Creamer, M. R., Wang, T. W., Babb, S., Cullen, K. A., Day, H., Willis, G., Jamal, A., & Neff, L. (2019). Tobacco product use and cessation indicators among adults—United States, 2018. *MMWR Morbidity and Mortality Weekly Report*, 68(45), 1013-1019. <https://doi.org/10.15585/mmwr.mm6845a2>
- Fiore, M. C., Jaén, C. R., Baker, T. B., Bailey, W. C., Benowitz, N. L., Curry, S. J., Dorfman, S. F., Froelicher, E. S., Goldstein, M. G., Heaton, C. G., Henderson, P. N., Heyman, R. B., Koh, H. K., Kottke, T. E., Lando, H. A., Mecklenburg, R. E., Mermelstein, R. J., Mullen, P. D., Orleans, C. T., ... Wewers, M. E. (2008) *Treating tobacco use and dependence: 2008 update*. Clinical Practice Guideline. U.S. Department of Health and Human Services, Public Health Service.
- Jaffee, K. D., Liu, G. C., Canty-Mitchell, J., Qi, R. A., Austin, J., & Swigonski, N. (2005). Race, urban community stressors, and behavioral and emotional problems of children with special health care needs. *Psychiatric Services*, 56(1), 63-69. <https://doi.org/10.1176/appi.ps.56.1.63>
- Klemperer, E. M., West, J. C., Peasley-Miklus, C., & Villanti, A. C. (2020). Change in Tobacco and Electronic Cigarette Use and Motivation to Quit in Response to COVID-19. *Nicotine & Tobacco Research*, 22(9), 1662-1663. <https://doi.org/10.1093/ntr/ntaa072>
- Lawless, M. H., Harrison, K. A., Grandits, G. A., Eberly, L. E., & Allen, S. S. (2015). Perceived stress and smoking-related behaviors and symptomatology in male and female smokers. *Addictive Behaviors*, 51, 80-83. <https://doi.org/10.1016/j.addbeh.2015.07.011>
- Lecic-Tosevski, D. (2019). Is urban living good for mental health?. *Current Opinion in Psychiatry*, 32(3), 204-209. <https://doi.org/10.1097/YCO.00000000000000489>
- Lindson, N., Klemperer, E., Hong, B., Ordóñez-Mena, J. M., & Aveyard, P. (2019). Smoking reduction interventions for smoking cessation. *The Cochrane Database of Systematic Reviews*, 9(9), CD013183. <https://doi.org/10.1002/14651858.CD013183.pub2>
- Lindson-Hawley, N., Hartmann-Boyce, J., Fanshawe, T. R., Begh, R., Farley, A., & Lancaster, T. (2016). Interventions to reduce harm from continued tobacco use. *The Cochrane Database of Systematic Reviews*, 10(10), Article CD005231.

- <https://doi.org/10.1002/14651858.CD005231.pub3>
- Martínez, C., Guydish, J., Le, T., Tajima, B., & Passalacqua, E. (2015). Predictors of quit attempts among smokers enrolled in substance abuse treatment. *Addictive Behaviors, 40*, 1-6.  
<https://doi.org/10.1016/j.addbeh.2014.08.005>
- McCullagh, P. (1980). Regression models for ordinal data. *Journal of the Royal Statistical Society: Series B (Methodological), 42*(2), 109-127.  
<https://doi.org/10.1111/j.2517-6161.1980.tb01109.x>
- Murphy, E. R. (2019). Transportation and homelessness: A systematic review. *Journal of Social Distress and the Homeless, 28*(2), 96-105.  
<https://doi.org/10.1080/10530789.2019.1582202>
- Muthén, B. (2011). Applications of causally defined direct and indirect effects in mediation analysis using SEM in Mplus. Paper available at:  
<https://www.statmodel.com/download/causalmediation.pdf>
- Nagawa, C. S., Fukunaga, M. I., Faro, J. M., Liu, F., Anderson, E., Kamberi, A., Orvek, E. A., Davis, M., Pbert, L., Cutrona, S. L., Houston, T. K., & Sadasivam, R. S. (2022). Characterizing pandemic-related changes in smoking over time in a cohort of current and former smokers. *Nicotine & Tobacco Research, 25*(2), 203-210.  
<https://doi.org/10.1093/ntr/ntac033>
- Neisler, J., Reitzel, L. R., Garey, L., Kendzor, D. E., Hébert, E. T., Vijayaraghavan, M., & Businelle, M. S. (2018). The moderating effect of perceived social support on the relation between heaviness of smoking and quit attempts among adult homeless smokers. *Drug and Alcohol Dependence, 190*, 128-132.  
<https://doi.org/10.1016/j.drugalcdep.2018.06.007>
- Nguyen, M.-A. H., Reitzel, L. R., Kendzor, D. E., & Businelle, M. S. (2015). Perceived cessation treatment effectiveness, medication preferences, and barriers to quitting among light and moderate/heavy homeless smokers. *Drug and Alcohol Dependence, 153*, 341-345.  
<https://doi.org/10.1016/j.drugalcdep.2015.05.039>
- Okuyemi, K. S., Caldwell, A. R., Thomas, J. L., Born, W., Richter, K. P., Nollen, N., Braunstein, K., & Ahluwalia, J. S. (2006). Homelessness and smoking cessation: insights from focus groups. *Nicotine & Tobacco Research, 8*(2), 287-296.  
<https://doi.org/10.1080/14622200500494971>
- Palepu, A., Hubley, A. M., Russell, L. B., Gadermann, A. M., & Chinni, M. (2012). Quality of life themes in Canadian adults and street youth who are homeless or hard-to-house: A multi-site focus group study. *Health and Quality of Life Outcomes, 10*, Article 93.  
<https://doi.org/10.1186/1477-7525-10-93>
- Popova, L., Henderson, K., Kute, N., Singh-Looney, M., Ashley, D. L., Reynolds, R. M., Nayak, P., & Spears, C. A. (2023). "I'm Bored and I'm Stressed": A Qualitative Study of Exclusive Smokers, ENDS Users, and Transitioning Smokers

- or ENDS Users in the Time of COVID-19, *Nicotine & Tobacco Research*, 25(2), 185-192. <https://doi.org/10.1093/ntr/ntab199>
- Reitzel, L. R., Short, N. A., Schmidt, N. B., Garey, L., Zvolensky, M. J., Moisiuc, A., Reddick, C., Kendzor, D. E., & Businelle, M. S. (2017). Distress tolerance links sleep problems with stress and health in homeless. *American Journal of Health Behavior*, 41(6), 760-774. <https://doi.org/10.5993/AJHB.41.6.10>
- Rigotti, N. A., Chang, Y., Regan, S., Lee, S., Kelley, J. H. K., Davis, E., Levy, D. E., Singer, D. E., & Tindle, H. A. (2021). Cigarette smoking and risk perceptions during the COVID-19 pandemic reported by recently hospitalized participants in a smoking cessation trial. *Journal of General Internal Medicine*, 36(12), 3786-3793. <https://doi.org/10.1007/s11606-021-06913-3>
- Savoy, E., Reitzel, L. R., Scheuermann, T. S., Agarwal, M., Mathur, C., Choi, W. S., & Ahluwalia, J. S. (2014). Risk perception and intention to quit among a tri-ethnic sample of nondaily, light daily, and moderate/heavy daily smokers. *Addictive Behaviors*, 39(10), 1398-1403. <https://doi.org/10.1016/j.addbeh.2014.05.002>
- Scheibein, F., McGirr, K., Morrison, A., Roche, W., & Wells, J. S. G. (2020). An exploratory non randomized study of a 3-month electronic nicotine delivery system (ENDS) intervention with people accessing a homeless supported temporary accommodation service (STA) in Ireland. *Harm Reduction Journal*, 17(1), Article 73. <https://doi.org/10.1186/s12954-020-00406-y>
- Smit, E. S., Fidler, J. A., & West, R. (2010). The role of desire, duty, and intention in predicting attempts to quit smoking. *Addiction*, 106(4), 844-851. <https://doi.org/10.1111/j.1360-0443.2010.03317.x>
- Soar, K., Dawkins, L., Robson, D., & Cox, S. (2020). Smoking amongst adults experiencing homelessness: A systematic review of prevalence rates, interventions, and the barriers and facilitators to quitting and staying quit. *Journal of Smoking Cessation*, 15(2), 94-108. <https://doi.org/10.1017/jsc.2020.11>
- Swayampakala, K., Thrasher, J., Carpenter, M. J., Shigematsu, L. M., Cupertio, A. P., & Berg, C. J. (2013). Level of cigarette consumption and quit behavior in a population of low-intensity smokers—longitudinal results from the International Tobacco Control (ITC) survey in Mexico. *Addictive Behaviors*, 38(4), 1958-1965. <https://doi.org/10.1016/j.addbeh.2012.12.007>
- U.S. Department of Housing and Urban Development (HUD)/Office of Community Planning (2021). *The 2020 annual homeless assessment report (AHAR) to congress*. Housing and Urban Development.
- Vijayaraghavan, M., Elser, H., Frazer, K., Lindson, N., & Apollonio, D. (2020). Interventions to reduce tobacco use in people experiencing homelessness. *The Cochrane Database of Systematic Reviews*, 12(12), CD013413.

<https://doi.org/10.1002/14651858.CD013413.pub2>

Wrighting, Q., Reitzel, L. R., Chen, T. -A., Kendzor, D. E., Hernandez, D. C., Obasi, E. M., Shree, S., & Businelle, M. S. (2019). Characterizing discrimination experiences by race among homeless adults. *American Journal of Health Behavior*, 43(3), 531-542.

<https://doi.org/10.5993/AJHB.43.3.8>

## Appendix

### Supplementary Table 1

*Multinomial Logistic Regression Analyses: The Effect of Urban Life Stress on Smoking Level/Readiness to Quit Among a Sample of Adults Who Smoke and Are Homeless (n = 411)*

Variables	Odds Ratio (95% CI)
<b>Smoking Level (reference: Non-daily smoker)</b>	
Light	1.014 (0.986, 1.044)
Moderate	1.026 (0.997, 1.056)
Heavy	1.047 (1.014, 1.082)**
<b>Readiness to Quit (reference: I enjoy smoking and have decided not to quit smoking for my lifetime. I have no interest in quitting.)</b>	
I definitely plan to quit smoking within the next 30 days.	1.029 (0.991, 1.068)
I definitely plan to quit smoking within the next 6 months.	1.012 (0.981, 1.044)
I never think about quitting smoking, and I have no plans to quit.	0.997 (0.959, 1.037)
I often think about quitting smoking, but I have no plans to quit.	1.012 (0.982, 1.043)
I rarely think about quitting smoking, and I have no plans to quit.	1.005 (0.972, 1.04)
I sometimes think about quitting smoking, but I have no plans to quit.	1.005 (0.975, 1.036)
I still smoke, but I have begun to change, like cutting back on the number of cigarettes I smoke. I am ready to set a quit date.	0.999 (0.967, 1.032)

*Note.* \*\*  $p < 0.01$ ; analyses controlled for age, sex, race, and mental health diagnosis.