Journal of International Agricultural and Extension Education

Volume 18 | Issue 3

Article 1

10-1-2011

Determinants of Adoption of Improved Maize Varieties and Chemical Fertilizers in Mozambique

Eunice Cavane Eduardo Mondlane University, cavaneeu@msu.edu

Cynthia Donavan Department of Agricultural, Food and Resource Economics Michigan State University, donavanc@anr.msu.edu

Follow this and additional works at: https://newprairiepress.org/jiaee

Part of the Growth and Development Commons, and the University Extension Commons

Recommended Citation

Cavane, E., & Donavan, C. (2011). Determinants of Adoption of Improved Maize Varieties and Chemical Fertilizers in Mozambique. *Journal of International Agricultural and Extension Education*, *18*(3), 5-22. DOI: https://doi.org/10.5191/jiaee.2011.18301

This Research Article is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Journal of International Agricultural and Extension Education by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Determinants of Adoption of Improved Maize Varieties and Chemical Fertilizers in Mozambique

Abstract

In Mozambigue, adoption of improved maize seed and chemical fertilizers is still limited. This study assessed farmers' attitudes towards hybrid maize SC513, Nitrogenous (N) Phosphorous (P) Potassium (K), (NPK 12-24-12) and urea fertilizers in highlands and lowlands of the Manica District. The study determined the influence of farmers' characteristics, attitudes, sources of information, and agroecological conditions on adoption of these technologies during 1995 through 2005. A questionnaire was administered during April and May 2006 with a randomly selected sample of 293 households. In general, farmers held positive attitudes towards improved maize varieties and chemical fertilizers, but the strength of attitudes towards fertilizers, in particular, varied by source of information. Farmers who learned about fertilizers from extension had stronger positive attitudes than farmers who learned about fertilizers from neighbors, although with hybrid seeds adoption, there was no significant difference between key sources. Overall, the number of farmers using SC513 was higher than the number of farmers using NPK and urea. Farmers' decision to adopt SC513 was positively associated with agro-ecological conditions, knowledge, production traits and marketability of the maize. Agro-ecological conditions, knowledge of fertilizer application, and extension contact influenced adoption of chemical fertilizers. The results differentiate a simpler process of adoption of new seed from a more complex process of adoption of fertilizers which demands greater knowledge of timing and soils as well as basic computational skills. Factors determining adoption of hybrid maize varieties and chemical fertilizers should be considered when designing extension programs for these technologies.

Keywords

Adoption, attitudes, chemical fertilizer, extension, improved maize varieties

Funding Source

I am deeply grateful to the Ford Foundation and the Department of Community, Agriculture, Recreation and Resource Studies at Michigan State University for providing the financial support for my PhD dissertation thesis, from which this paper is based From the Executive and Managing Editors

DOI: 10.5191/jiaee.2011.18301

Determinants of Adoption of Improved Maize Varieties and Chemical Fertilizers in Mozambique

Eunice Cavane Assistant Lecturer, Faculty of Agronomy and Forestry Engineering Department of Crop Production and Protection Eduardo Mondlane University ecavane@uem.mz or cavaneeu@msu.edu

Cynthia Donavan Assistant Professor, International Development Department of Agricultural, Food and Resource Economics Michigan State University donavanc@anr.msu.edu

Abstract

In Mozambique, adoption of improved maize seed and chemical fertilizers is still limited. This study assessed farmers' attitudes towards hybrid maize SC513, Nitrogenous (N) Phosphorous (P) Potassium (K), (NPK 12-24-12) and urea fertilizers in highlands and lowlands of the Manica District. The study determined the influence of farmers' characteristics, attitudes, sources of information, and agro-ecological conditions on adoption of these technologies during 1995 through 2005. A questionnaire was administered during April and May 2006 with a randomly selected sample of 293 households. In general, farmers held positive attitudes towards improved maize varieties and chemical fertilizers, but the strength of attitudes towards fertilizers, in particular, varied by source of information. Farmers who learned about fertilizers from extension had stronger positive attitudes than farmers who learned about fertilizers from neighbors, although with hybrid seeds adoption, there was no significant difference between key sources. Overall, the number of farmers using SC513 was higher than the number of farmers using NPK and urea. Farmers' decision to adopt SC513 was positively associated with agroecological conditions, knowledge, production traits and marketability of the maize. Agroecological conditions, knowledge of fertilizer application, and extension contact influenced adoption of chemical fertilizers. The results differentiate a simpler process of adoption of new seed from a more complex process of adoption of fertilizers which demands greater knowledge of timing and soils as well as basic computational skills. Factors determining adoption of hybrid maize varieties and chemical fertilizers should be considered when designing extension programs for these technologies.

Keywords: Adoption, attitudes, chemical fertilizer, extension, improved maize varieties

Introduction

Improving maize production is one of the most important strategies for food security in Mozambique. However, chemical fertilizers and improved maize varieties are not yet widely adopted in Mozambique. In 2008 less than 5 percent of approximately three million farmers used chemical fertilizers and about 10 percent used improved maize varieties (Ministry of Agriculture [MINAG], 2009). Hence a widespread adoption of improved seed varieties and chemical fertilizers is a challenge for agricultural development policy in Mozambique.

Improved maize varieties include hybrids and open pollinated varieties (OPVs) whose traits have been improved for selected characteristics such as drought tolerance, disease resistance, short maturity rate, increased vield per unit of land, and quality protein (Byerlee, 1994). Adoption of improved maize varieties can help farmers face labor constraints, food insecurity and lack of income. Research in Southern Africa (Byerlee, 1994; Kaliba et al., 2000) indicated that farmers prefer early maturing maize to better deal with labor constraints, risk considerations and crop rotations. Sometimes farm households would improve food security by planting an early maturing variety that can be consumed in the "hungry season," before the main harvest. In areas with small scale irrigation and near market places for fresh maize, early maturing varieties provide sources of income for farmers (Rotter and Keulen, 1996).

Expanded use of fertilizer in Sub-Saharan Africa has been stressed as one of the solutions to alleviate production shortfalls and land degradation in the region (Pinstrup-Andersen, Pandya-Lorch, & Rosegrant, 1997). Farmers can benefit from applying chemical fertilizer to maize. When used in optimal amounts, chemical fertilizers increase production and farm efficiency. Optimal applications of commercial fertilizers replenish the nutrients removed by the crop and in some cases exceed levels of nutrients found in the soils before they were farmed. In Kenya, Rotter and Keulen (1996) found that there was a tremendous potential for increasing maize yields, and hence national production, by applying moderate amounts of nitrogen (N) and phosphorous (P) fertilizers in the lowlands and midlands.

For widespread adoption of improved varieties and chemical fertilizer by farmers, extension educators need to understand the factors affecting technology adoption (Abebaw & Belay, 2001). Adoption of technology is influenced by physical, socio-economic, and mental factors including agro-ecological conditions, age of farmer, family size, education of farmer, how-to-knowledge, source of information, and farmer's attitudes towards the technology (Feder et al., 1985; Byerlee & Polanco, 1986; Neupane et al., 2002; Rogers, 2003). High levels of adoption of improved maize varieties and chemical fertilizers are more likely to be found among farmers located in agro-ecological regions with high rainfall (Kaliba et al., 2000: Hintze et al., 2003). Young farmers are more likely to adopt a new technology because they have had more schooling and are more open to attitude change than older farmers (International Maize and Wheat Improvement Center [CIMMYT], 1993; Visser & Krosnick, 1998). Education is expected to enhance the decision making and the adoption of agricultural technologies. Family size plays a role on labor provision. Adoption of new varieties often requires more labor inputs (Feder et al., 1985). It is assumed that large families provide the labor required for improved maize production practices. Access to sources of information, such as extension, market, and neighbors, enhances the adoption of technology (Abebaw & Belay, 2001). Knowledge influences adoption. Farmers who have adequate knowledge of technology use are more likely to adopt it (Abebaw & Belay, 2001; Rogers, 2003).

Journal of International Agricultural and Extension Education

Knowledge of technology application acts as an intervening variable between "attitudes towards the technology" and "use of the technology" (Rogers & Havens, 1961). Farmers' attitudes are important in determining adoption of improved technology. Attitudes are evaluative responses towards the technology, and are formed as farmers gain information about it. Adopters tend to hold positive attitudes towards the technology (Chilonda & Van Huylenbroeck, 2001). The literature suggests that attitudes towards improved maize varieties place great emphasis on two factors: production characteristics and income factors. Similarly, for chemical fertilizers, the literature suggests two factors: effect of fertilizer and costs factors (Envong et al., 1999; Thompson, 1992).

The objectives of this study were twofold: firstly, to assess farmers' attitudes towards adoption of hybrid maize SC513, NPK, and urea fertilizers in highlands and lowlands of the Manica District; and secondly, to determine the influence of farmers' characteristics, attitudes, source of information, and agro-ecological conditions on adoption of hybrid maize SC513, NPK, and urea fertilizers.

Materials and Methods

This study involved a cross-sectional survey with 293 randomly selected households growing maize in the highlands of Machipanda and lowlands of Vanduzi in the Manica District, central region of Mozambique. Machipanda and Vanduzi were selected purposely because of the importance of maize in the farming systems and the availability of maize technology dissemination programs in the two areas. Vanduzi and Machipanda represent two distinct agro-ecological zones, R4 and R10 respectively. Machipanda is located at higher altitude (900-1500 m), and gets higher annual rainfall (900-1500 mm) than Vanduzi which is located at 200-1000 m above sea level with annual rainfall of about 900-1000 mm.

Data were collected between April and May, 2006, using personal interviews with 120 farmer household heads from Machipanda, and 173 fromVanduzi. The explanatory variables measured in the questionnaire are presented in Table 1.

The Logistic Model

In the field of agriculture, adoption of technologies is measured as a dichotomous response variable (0 = non adoption of innovation and 1 = adoption of innovation). The logistic model is the standard method of analysis, when the outcome variable is dichotomous (Hosmer & Lemeshow, 2000). The logistic regression model characterizing adoption of SC513, NPK or urea by the sample of households is specified as follows:

$$\pi = \frac{e^{\alpha + \beta 1x1 + \beta 2x2 + \dots + \beta pXp + \varepsilon}}{1 + e^{\alpha + \beta 1x1 + \beta 2x2 + \dots + \beta pXp + \varepsilon}}$$
(1.0)

Where:

π, is the actual proportion of farmers adopting the technology for particular values of independent variables X₁, X₂,..., X_p, that influence adoption of SC513, NPK or urea. $β_1, β_2,..., β_p$ denote the regression coefficients associated with independent variables X₁, X₂, ..., X_p. ε is the error term. From the equation (1.0), we arrive at a simple linear regression equation through *logit transformation* (Chatterjee et al., 2000; Hosmer and Lemeshow, 2000):

 $\log[\pi/(1-\pi)] = \alpha + \beta 1x1 + \beta 2x2 + \dots + \beta pXp + \varepsilon \quad (2.0)$

Variable name	Variable	Description
	type	
Agro ecological region	Dichotomous	0 = Lowlands (Vanduzi), $1 =$ Highlands
(X_1)		(Machipanda)
Age (X ₂)	Continuous	Farmer's age (years)
Family size (X ₃)	Continuous	Number of members in the family
Level of education (X ₄)	Dichotomous	0 = illiterate, $1 =$ some schooling
Knowledge of advantages and disadvantages of improved maize and knowledge of fertilizer application(X ₅)	Dichotomous	0 = not knowledgeable, 1 = knowledgeable
Source of information (X ₆)	Multiple category variable	For improved maize SC513: 1 = neighbors, 2 = market, 3 = extension For chemical fertilizers: 1= market, 2 = extension, 3 = neighbors
Attitude (X7)	Interval	Factor scores for attitude toward marketability of grain of improved maize SC513 Factor scores for attitude toward production traits of improved maize SC513 Factor scores for attitude toward NPK Factor scores for attitude toward urea

 Table 1

 Description of Explanatory Variables Measured in the Ouestionnaire.

Data Analysis

Data were analyzed using the Statistical Package for Social Sciences SPSS version 15 (SPSS Inc. 2006). Attitude statements were measured using a 5-point Likert scale with 1 = Strongly Agree, 2 =Agree, 3 = Neutral, 4 = Disagree, and 5 =Strongly Disagree. Exploratory and confirmatory factor analyses were used to create summated scales of attitudes towards SC513, NPK and urea. A two-factor model for improved maize variety SC513 had an excellent fit, $X^2 = 7.550$, p > 0.05. Comparative index (CFI) was greater than 0.90 and the root mean square error of approximation (RMSEA) was lower than 0.05. A one-factor model for NPK had also a good fit, $X^2 = 7.285$, p > 0.01. Comparative

index (CFI) was greater than 0.90. Although the RMSEA was greater than 0.05, the 90% confidence interval for RMSEA was 0.030 to 0.184. This interval includes 0.05 criteria which indicate "reasonable" errors of approximation in the population. A onefactor model for urea had a reasonable fit. Comparative index (CFI) was slightly greater than 0.90. This means that although X^2 and RMSEA had values of 31.267 and 0.237 respectively, which suggest a poor fit, the one-factor model is still better than the complete or baseline model for urea.

Each of the two factors of the scale on attitudes towards maize variety SC513 had three items. Hence, for each factor the summed score across items for a given respondent ranged from 3 to 15, with scores lower than 9 indicating favorable attitudes, and scores greater than 9 indicating unfavorable attitudes. The scale of attitudes towards NPK /Urea had four items. The summed score across items for a given respondent ranged from 4 to 20, with scores lower than 12 indicating favorable attitudes, and scores greater than 12 indicating unfavorable attitudes. One-Way ANOVA was used to compare mean attitude scores for the three groups of sources of information on the technology, neighbors, market, and extension.

To create attitude scores for subsequent use in logistic regression, factor scores were estimated through principal component and varimax procedure (Hair et al., 2005). Two-factor solutions for the attitude scale on SC513, explained about 59.3 percent of the variance in attitudes toward SC513. One factor solution for the attitude scale on NPK explained about 48 percent of the variance in attitude toward it. One factor solution for the attitude scale on urea explained about 56.4 percent of the variance in attitudes toward urea. Variance Inflation Factor (VIF) estimates were examined for collinearity diagnostic in Multiple Logistic Regression. Most of VIF estimates had values less than 2, which indicate no serious problems of collinearity.

Results and Discussion

Characteristics of farmers

The last section of the questionnaire measured the general characteristics of the farmers. The results are presented in Table 2.

Table 2

Characteristics		Stud	Study Location			
	Machip	anda	Van	Iduzi		
	Frequency	Percent	Frequency	Percent		
Age (Years)	(n = 120)		(n = 173)			
Less than 44	65	54.2	89	51.4		
45 to 60	38	31.7	59	34.1		
61 to 76	12	10.0	21	12.1		
More than 76	5	4.2	4	2.3		
Level of education	(n = 120)		(n = 173)			
Illiterate	17	14.2	53	30.6		
Primary School	95	79.2	109	63.0		
Secondary/High School	8	6.7	11	6.4		
Family size	(n = 120)		(n = 173)			
Less than 8	79	65.8	112	64.7		
9 to 12	30	25.0	44	25.4		
More than 13	11	9.2	17	9.8		

Farmers were similar regarding age and family size. In both study locations the majority of farmers had an age < 44 years and family size < 8 members. However, for level of education the respondents showed differences. Vanduzi had a larger number (30.6 %) of illiterate farmers than Machipanda (14.2 %). It is possible that farmers in Machipanda had more access to school due to proximity to Zimbabwe, than farmers in Vanduzi.

The results on family size indicate that about 25 percent of households in both study areas have between 9 and 12 members. These numbers are higher than the average family size in

rural areas of Mozambique which is approximately six members. Assuming that adoption of new varieties requires more labor inputs (Feder et al., 1985), we would think that 25 percent of the rural households in Machipanda and Vanduzi, have relatively large families to rely upon for labor demands. However, the total number of family members does not always mean availability of labor, because some families may have higher dependency rates than others. Thus while the relatively large family size may suggest more labor for cultivation, more precise results on labor availability would need to be provided by information on dependency ratios.

Farmers' Attitudes Towards Improved Maize SC513, NPK and Urea Fertilizers

To capture farmers' attitudes, farmers were asked the extent of their agreement with six statements on characteristics and marketability of improved maize variety SC513, four statements on chemical fertilizer NPK, and four statements on chemical fertilizer urea. Tables 3, 4 and 5 present farmers' attitudes towards improved maize SC513, NPK and urea fertilizers. Mean score was based on a 5-point interval scale where *1 equals Strongly Agree, 2 equals Agree, 3 equals Neutral, 4 equals Disagree, and 5 equals Strongly Disagree*. The scale was reversed for negative statements.

Ţ	
al.	
$\underline{\succ}$	
CP.	
ω	

Ы	
$\boldsymbol{\omega}^{1}$	
7	
2	
9	
armers'	
<u>ب</u>	
1	
Ξ	
ť	
ã	
\tilde{o}	
Ś	
H	
2	
Ż.	
3	
à	
$\mathcal{S}_{\mathcal{I}}$	
5	
3	
ā.	
ç.	
Z.	
6	
~	
\leq	
2	
ត	
0	
4	
2	
Ξ.	
0	
2	
'Attitudes Towards Improved Maize Variety SC513	
Õ.	
S.	
-	
ω	

Farmers' Attitudes Towards Improved Maize Variety SC513	Impro-	ved Mai	ze Varie	etv SC5	13									
		Percent of Response Categories in	of Resp	onse Ca	ategor	ies in		Pe	ercent o	Percent of Response Categories in	nse Ca	tegories	s in	
			Macl	Machipanda	(Van	Vanduzi	(
			(n =	(n = 115)						= n)	(n = 91)			
	SA	° A	۲	R D				SA	A	۲	D	SD		3
Attitude toward	0/7	70	/0	0/7	/0	IVI	UC	70	70	0/2	0/7	70	IM	SU U
marketability of improved maize variety SC513.														
Grain from SC 513 is easy to sell.	7.8	66.1	19.1	6.1	.9	2.3	.7	7.7	62.6	29.7	ı	ı	2.2	6.
Fresh maize (roasting cobs) of SC 513 is easy	7.0	51.3 38.3		2.6	.9	2.4	.7	2.2	50.5	41.8	4.4	1.1	2.5	.7
to sell. Planting SC 513 is a waste of time and	4.3	77.4 5.2		11.3	1.7 2.3	2.3	.∞	3.3	3.3 67.0 24.2	24.2	4.4	1.1	2.3	.7
money. Attitude toward														
Attitude toward production characteristics of SC513.														
good germination.	12.2	78.3	7.0	1.7	.9	2.0	.6	20.9	46.2	33.0	ı	ı	2.1	.7
Grain from SC 513 is good for milling.	28.7	65.2	3.5	2.6	ı	1.8	.6	26.4	42.9	28.6	1.1	1.1	2.1	.8
When rainfall is low, Chimanhica has better production than SC 513.	7.0	7.0 73.9 7.0	7.0	12.2		2.2	.∞	5.5	5.5 53.8 27.5	27.5	13.2	ı	2.5	.∞

II Journal of International Agricultural and Extension Education

Farmers' Attitudes Towards NPK % NPK is good for maize. % % % NPK is not good for the soil. 5. NPK is a waste of time and money. 1.	SA % 5.1 6.0 2.6	Percent of Response Categories in Machipanda Machipanda $(n = 117)$ A N D SD $%$ $%$ $%$ M 80.3 6.8 6.8 .9 2.2 78.6 10.3 5.1 - 2.1 16.2 11.1 59.0 11.1 3.6 61.5 6.0 29.1 1.7 2.7	of Resj Mao (n N % 6.8 10.3 11.1 6.0	Response Ca Machipanda $(n = 117)$ D N D D N_6 9_6 0.3 5.1 0.3 5.1 1.1 59.0 1.1 59.0 29.1	Categor la SD % - 11.1 1.7	ries in <u>M</u> 2.1 3.6 2.7	<i>SD</i> .7 .6 1.0	Pe SA 5.2 2.8 4.1	Percent of: A A A 5 % 8 78.6 2 79.3 8 24.8 1 56.6	rf Respo Var (n = N 15.2 14.5 34.5 25.5	Response Categories in Vanduzi (n = 145) N D SD % % % <u>//</u> 15.2 1.4 - 2. 15.2 1.4 - 2. 14.5 - 2. 34.5 34.5 3.4 3. 25.5 12.4 1.4 2.	1.4	es in <u>M</u> 2.1 3.1 2.5	
Table 5														
Turmers Allunes Iowards	Jrea													
<u>r'urmers</u> Aunuues rowurus Orea	Jrea	Percent of Response Categories in Machipanda	ofRes	ponse (Categor	ries in		Pé	Percent of		Response Categories in Vanduzi	utegori	es in	
runners Annuaes Lowaras o	Jrea	Percent	of Res Maa (n	Response Ca Machipanda $(n = 118)$	Categor	ies in		Pé	Arcent o		Response C $_{i}$ Vanduzi (n = 144)	utegori	es in	
runners Annuaes rowards o	Jrea SA	Percent A	of Res Mau (n N	ponse (chipanci = 118) D %	Categor la SD	ies in	GS	Pe SA	A A		nduzi 144) %	utegori SD	es in	
Urea is good for maize	Jrea SA %	Percent A %	of Res Mao (n N %	ponse (chipand = 118) D %	Categor la SD %	$\frac{1}{1.9}$.4 <i>SD</i>	Pe 6.3	A 74.3		onse C _z nduzi 144) D 1.4	utegori SD .7	$\frac{1}{2.2}$ is in	
Urea is good for maize Urea increases maize yield.	Jrea SA % 15.3 17.8	Percent A % 82.2 78.0	of Res Mao (n N % 1.7 3.4	ponse (chipanci = 118) D % .8	Categor la SD -	$\frac{M}{1.9}$	SD 4. 2.	Pe SA 9% 9.0	A 9% 74.3 74.3		onse C₂ nduzi 144) D 0% 1.4 .7	utegori SD % .7	es in <u>M</u> 2.2	
Urea is good for maize Urea increases maize yield. Urea is not good for the soil	Jrea SA % 15.3 17.8 7.6	Percent A % 82.2 78.0 64.4	of Res Mao (n N % 1.7 3.4	ponse (chipanci = 118) D % .8 .8 .8	Lategor la SD -	ies in <i>M</i> 1.9 2.4	.9 .5 .5	Pe SA 9.0 3.5	A A 74.3 74.3 31.9		onse C <i>ε</i> nduzi = 144) D % 1.4 .7 .7 .7	utegori SD .7 .7 .7	M 2.1 2.9	

Journal of International Agricultural and Extension Education

Volume 18, Number 3

12

Farmers' attitudes towards SC513 consisted mostly of cognitive responses. expressed as beliefs by farmers about the relationship between the seed and the production and marketability characteristics that describe the seed. In general, respondents from Machipanda and Vanduzi held positive attitudes towards the marketability of improved maize variety SC513, production characteristics of SC513, and use of NPK and urea for maize production. For improved maize variety SC513, most farmers in Machipanda (93.9%) and Vanduzi (63.3%) appreciated the quality of the seed and maize meal obtained with maize variety SC513. Other studies found that farmers are favorable toward production characteristics of improved maize varieties (Hintze et al., 2003).

For chemical fertilizers most farmers (>80%) in Machipanda and Vanduzi appreciated the effect of NPK and urea on increasing maize yield. Farmers agreed that NPK and urea were good for increasing maize yield. Farmers' attitudes towards use of chemical fertilizer on maize consisted of both cognitive and affective responses. The cognitive responses were expressed as beliefs farmers have about the relationship between the fertilizer and its capacity of increasing maize yield as well as the relationship between the fertilizer and its value for maize production.

The affective response was expressed as a good or bad feeling

experienced by farmers when fertilizers are paired with their effect on the crop and soil. For example, farmers, particularly in Machipanda, expressed a negative attitude towards the effect of NPK on the soil. These farmers might have learned the affective response "Chemical fertilizer is not good for the soil" through direct emotional experiences offered by those instances when farmers stopped applying fertilizer and obtained low yields. As farmers explained when asked why they thought chemical fertilizer was not good for the soil, "NPK kills the soil. Once you use it, you should keep using it. The soil becomes dependent on it. If you stop using NPK your soil will not produce as much as before" (Farmers in Machipanda, 2006). This suggests that some farmers in Machipanda view the application of NPK as an intrusion of plant nutrients into a balanced environment. Another aspect explicit in farmers' explanations is a fear of their maize production becoming dependent on an input (NPK) which they feel cannot afford buying it regularly. These results highlight the need of increasing farmers' knowledge of optimal fertilizer use, and improving access to fertilizer.

Influence of Source of Information on Farmers' Attitudes Towards Improved Maize SC513, NPK and Urea Fertilizers

This study also performed comparisons between mean attitude scores and sources of information within each study location. The results, of ANOVA and Scheffe's post hoc multiple comparison method, are presented in Table 6.

18, N
\mathbf{a}
Jumber 3

2	
С,	
1	
CD	
6	

Sample ANOVA Table for Mean Attitude by Sources of Information

Machipanda	و	Μ	Machipanda	da				Vanduzi	IZ.
	Source								
	Information	Ν	Mean	F	Sig	Ν	Mean	F	Sig.
Attitude toward	Neighbors	72	6.81			28	7.43		
marketability of	Market	35	7.37	2.02	.138	26	7.15	2.05 .135	.135
improved maize variety	Extension	8	6.25			37	6.73		
Attitude toward	Neighbors	72	5.99			28	7.04		
production	Market	35	6.26	.661	.661 .518	26	7.08	2.79	.067
characteristics of SC513.	Extension	8	5.75			37	6.14		
Attitude toward NPK.	Market	41	10.7			25	9.52		
	Extension*	19	10.2	.512	.600	32	9.03	5.76	.004
	Neighbors*	57	10.6			88	10.2		
Attitude toward urea.	Market	41	8.63			26	9.46		
	Extension**	19	8.47	1.34	.265	30	8.33	8.53	.000
	Neighbors**	85	8.05			88	10.06		

In the lowlands, the strength of positive attitudes towards fertilizers differed significantly (P<.01 and P<.001) between farmers who learned about fertilizers from extension and those who learned from neighbors. Farmers who learned about fertilizers from extension had stronger positive attitudes than farmers who learned about fertilizers from neighbors. Personal contacts between extension agents and farmers are more effective for delivering specific information to farmers and eventually persuade farmers to use fertilizers. The results show that while all three sources of information, neighbors, market and extension can be used by farmers to learn and raise awareness about fertilizers, particular attention should be given to public extension to strengthen farmers' attitudes towards fertilizers.

Pattern of Adoption of Improved Maize Variety SC513 and Chemical Fertilizers NPK and Urea in Machipanda and Vanduzi

The results on patterns of adoption of SC513, NPK and urea in Machipanda and Vanduzi are presented in Figures 1, 2, and 3. The curves are based on respondents' recall of the year in which they first used the technology.




Figure 1. Pattern of Adoption of Hybrid Maize SC513 in Machipanda and Vanduzi from 1990 – 2005.




Figure 2. Pattern of Adoption of Fertilizer NPK in Machipanda and Vanduzi from 1970 – 2005.




Figure 3. Pattern of Adoption of Fertilizer urea in Machipanda and Vanduzi from 1970 – 2005.

Farmers in Machipanda and Vanduzi, have been using improved maize variety SC513 since early 1990, while chemical fertilizers for maize production started much earlier, before 1990. Improved maize variety SC513 and chemical fertilizers NPK and urea were formally introduced in the study area in 1995. Since then new farmers joined each year increasing the percentage of usage (i.e. the number of farmers who have used the technology). This increase was more rapid in highlands of Machipanda than in lowlands of Vanduzi. A possible explanation is that hybrid maize seeds do better than local varieties in high rainfall regions and in these regions higher incidence of fertilizer use are also expected than in the lowland regions (Kaliba et al., 2000). In addition, Machipanda is much closer to Zimbabwe and farmers had access to inputs there. Since the increase of percentage of usage was different between the two agro ecological regions, these findings call attention to the evaluation of the suitability and accessibility

of improved maize technology in a particular agro-ecological zone.

Discontinuance of SC513, NPK and Urea

The pattern of adoption provides a positive picture in the sense that it expresses that through the years the number of farmers who had used the technologies was accumulating. Nevertheless, the current adoption was not high. For example, in both study areas the percentage of farmers cultivating improved maize variety SC513 and applying chemical fertilizer on maize during the growing season 2004 - 05 was lower than the cumulative percentage (Table 7). The current adoption (i.e. adoption in years 2004 and 2005) indicates that discontinuance occurred throughout the time since the first year farmers started using the technologies. One obvious reason for reduced adoption is discontinuance. After farmers had adopted the technologies, they may discontinue their use for various reasons including replacement, dissatisfaction, and misuse of the technology (Rogers, 2003).

Table 7

Percentage of Farmers Ever Used and Currently Using the Technology

Technologies		Study l	ocation	
	Mac	chipanda	V	anduzi
	Ever used	Using 2004-05	Ever used	Using 2004-05
Hybrid maize SC 513	88% (105)	37.7% (45)	28% (48)	5.8% (10)
Fertilizers (NPK/urea)	87% (102)	33.3% (39)	33% (48)	2.8% (4)

Reasons for Discontinuance of SC513, NPK and urea

The reasons for discontinuance of SC513, NPK and urea, were also explored. The discontinuance of improved maize SC513 was mainly due to dissatisfaction with husk cover, susceptibility to storage insects, and replacement of hybrid variety SC513 with OPV and local maize varieties. The discontinuance of fertilizers was, mainly, due to lack of money to purchase NPK and urea. The reasons for discontinuance have important implications for maize breeders and social scientists. Breeders need to improve the husk cover and enhance resistance to storage insects to make the hybrid variety SC513 attractive to farmers. Social researchers need to investigate ways by which farmers can be provided with fertilizers at accessible costs including provision of fertilizer on credit basis. Factors associated with adoption of improved maize SC513, NPK and urea fertilizers

Tables 8, 9 and 10 present the

maximum likelihood estimates of the logistic models for factors associated with adoption of SC513, NPK and urea.

Table 8

Maximum Likelihood Estimates of Logistic Model for Factors Affecting Adoption of SC513

Variables	β	S.E.	Wald	Sig.	Exp(β)
Agro-ecological zone	2.51	.512	24.0	.000	12.2
Knowledge of advantages and disadvantages of SC513	1.99	.732	7.40	.007	7.32
Attitude toward marketability of produce from SC513	497	.234	4.50	.034	.608
Attitude toward traits of SC513	812	.228	12.6	.000	.444
Constant	704	1.33	.279	.598	.495
Likelihood ratio chi-square df(10)	87.03			.000	

Table 9

Maximum Likelihood Estimates of Logistic Model for Factors Affecting Adoption of NPK

Variables	β	S.E.	Wald	Sig.	Exp(β)
Agro-ecological zone	.794	.392	4.10	.043	2.21
Knowledge of application methods for NPK	2.61	.753	12.0	.001	13.6
Information sources on NPK			14.5	.001	
Market (1)	.154	.442	.122	.727	1.17
Extension (2)	1.57	.430	13.3	.000	4.81
Constant	-4.32	1.11	15.1	.000	.013
Likelihood ratio chi-square df(8)	64.3			.000	

Table 10

Maximum Likelihood Estimates of Logistic Model for Factors Affecting Adoption of Urea

	5			~ 1				
Variables	β	S.E.	Wald	Sig.	$Exp(\beta)$			
Agro-ecological zone	1.43	.369	14.9	.000	4.16			
Knowledge of application methods for urea	2.05	.521	15.5	.000	7.74			
Information sources on urea			9.99	.007				
Market (1)	211	.418	.256	.613	.810			
Extension (2)	1.23	.434	8.01	.005	3.41			
Constant	-2.86	.964	8.81	.003	.057			
Likelihood ratio chi-square df(8)	81.6			.000				
The fit of the models was satisfactory. The estimated coefficients for the likelihood ratio chi-square were	of 87.0,	significant (P<.001), with chi-square value of 87.0, 64.3, and 81.6. The models accounted (R ² _{Logistic} .) for 39, 24 and 27						

Journal of International Agricultural and Extension Education

percent of the variation between adopters and non-adopters of SC513, NPK, and Urea respectively.

As expected, the agro - ecological zone and how-to knowledge were positively (P < .05) associated with the adoption of improved maize variety SC513, NPK, and urea fertilizers. Being in the highlands increased the probability of adoption of the three technologies, by a factor of 12, 2 and 4 respectively. These results can be related to the effect of high annual rainfall in the highlands of Machipanda. This argument is supported by other studies (Kaliba et al., 2000; Hintze et al., 2003). Hybrid maize might be more adaptable to the highlands of Machipanda than to the lowlands of Vanduzi. Likewise, fertilizer use is expected to be higher in highlands where rainfall is relatively higher than lowlands.

Feeling knowledgeable about advantages and disadvantages of improved maize and about application of chemical fertilizers increased the probability of adoption of the three technologies, by a factor of 7, 14, and 8. Positive association between knowledge of fertilizer application and adoption of fertilizer was also mentioned by Rogers and Havens (1961). These researchers found that knowledge of fertilizer (i.e. how to use fertilizer, and what nutrients the crop needs) acted as an intervening variable, between "attitude toward fertilizer" and the "use of fertilizer." CIMMYT (1993) also highlights the importance of basic computation skills for proper use of fertilizers.

Extension was a significant factor (P<.05) for adoption of fertilizers. The probability of adoption of NPK and urea, for farmers who learned about these technologies from extension services, increased 5 and 3 times more than the probability of adoption by farmers who learned from neighbors. Other studies had found positive relationships between extension and chemical fertilizers (Abebaw and Belay, 2001; Kaliba et al., 2000). The

results indicate that strengthening the provision of information on the use of fertilizers through extension (i.e. results from demonstration plots, individual or group meetings) may improve the levels of adoption of chemical fertilizers in Machipanda and Vanduzi. Moreover, given the relevant role played by extension in the adoption of chemical fertilizers, one option would be to concentrate extension resources to train farmers in the optimal use of fertilizers. Another option would be to promote vocational training on improved maize varieties technology, including chemical fertilizers, in the rural schools. For improved maize variety, contrary to what was expected, extension had no significant effect on adoption of SC513. As a source of information, extension had a positive but insignificant impact on the adoption decisions of SC513. It is plausible that all sources of information (market, neighbors, extension, and market) provide farmers with useful information.

Farmers' attitudes were significant factors of adoption of improved maize. As expected, holding negative attitudes toward traits of SC513 and negative attitudes toward the marketability of produce from SC513 had a significant (P<.05) negative effect on the logarithm of the odds of adoption. Other studies found that farmers' perceptions of production characteristics (yield, maturity rate, drought resistance, insect resistance lodging resistance, grain weight) determine variety selection and adoption (Hintze et al., 2003). These results imply that researchers should continue and strengthen research on production characteristics of improved maize varieties, and public extension services as well as the private sector (seed companies, wholesalers and retailers) should emphasize messages on the production characteristics of the improved maize varieties, including seed quality, drought tolerance and maize meal quality. Positive attitudes toward selling grain and fresh maize of SC513 were also a

significant factor of adoption. This association indicates that farmers obtain some surplus from SC513 that can be used for household income. Thus, in addition to production characteristics, and benefits associated with improved maize varieties, extension messages should provide information on prices and demand for it. Breeders should develop maize varieties for which output is marketable. The private sector should provide farmers with marketing information.

Overall the three most important factors associated with adoption of improved maize SC513, in decreasing order were: agro ecological region, attitude toward production traits of improved maize SC513, and knowledge about advantages and disadvantages of improved maize varieties. Extension on NPK and urea, knowledge about fertilizer application, and agroecological conditions had significant influence on adoption of chemical fertilizers.

Conclusion

This study showed that, in general, farmers in Machipanda and Vanduzi do support the use of improved maize varieties and chemical fertilizers, but attitude strength varied according to sources of information. In the lowlands farmers who learned about fertilizers from extension had stronger positive attitudes than farmers who learned about fertilizers from neighbors. Different patterns of adoption were found in this study. The number of farmers using improved maize SC513 was higher than the number of farmers using NPK and urea for maize production. For each of the three technologies the rate of usage differed between the highlands and lowlands. The increase was more rapid in highlands than in lowlands. Hence these findings call attention to the evaluation of the suitability and accessibility of improved maize production technology in a particular agro-ecological zone.

Regarding factors of adoption, this study has confirmed earlier research which showed that adoption of improved maize varieties and chemical fertilizers technologies, is influenced by agro ecological conditions, attitude toward production traits and marketability of improved maize, how-to-knowledge to apply the technology, and the role of extension in dissemination of improved technology. Therefore, these factors should be considered when planning, implementing and evaluating extension programs for dissemination of improved maize varieties and chemical fertilizers. Farmers' characteristics such as age, level of formal education, and family size did not influence significantly the adoption of improved maize variety SC513 and chemical fertilizers NPK and urea, in the highlands of Machipanda and lowlands of Vanduzi in the Manica District.

Acknowledgments

I am deeply grateful to the Ford Foundation and the Department of Community, Agriculture, Recreation and Resource Studies at Michigan State University for providing the financial support for my PhD dissertation thesis, from which this paper is based.

References

- Abebaw, D., & Belay, K. (2001). Factors influencing adoption of high yielding maize varieties in Southwestern Ethiopia: An application of logit. *Quarterly Journal of International Agriculture, 401,* 49-167.
- Byerlee, D. (1994). Maize research in Sub-Saharan Africa: an Overview of past impact and future prospects.
 Economics. Working Paper 94-03.
 Mexico, D. F: CIMMYT.
- Byerlee, D., & de Polanco, H. E. (1986). Farmers' stepwise adoption of technological packages: Evidence from the Mexican Altiplano.

Journal of International Agricultural and Extension Education

American Journal of Agricultural Economics, 68, 519-527.

- Chatterjee, S., Hadi, A. S., & Price, B. (2000). *Regression analysis by example* (Third Edition). New York: Wiley Series in Probability and Statistics.
- Chilonda, P., & Van Huylenbroeck, G. (2001). Attitudes towards and uptake of veterinary services by small-scale cattle farmers in Eastern province Zambia. *Outlook on Agriculture, 30,* 231-218.
- Enyong, L. A., Debrah, S. K., & Batione, A. (1999). Farmers' perceptions and attitudes toward introduced soil-fertility enhancing technologies in Western Africa. *Nutrient Cycling in Agroecosystems*, 53, 177-187.
- Feder, G., Just, R. E., & Zilberman, D. (1985). Adoption of agricultural innovation in developing countries: A Survey. *Economic Development* and Cultural Change, 33, 225-298.
- Hair, J. F., Black, W. C., Babin, B. J.,
 Anderson, R. E., & Tatham, R. L.
 (2005). *Multivariate data analysis*(Sixth Edition). New Jersey: Pearson
 Prentice Hall.
- Hintze, L. H., Renkow, M., & Sain, G. (2003). Variety characteristics and maize adoption in Honduras. *Journal* of Agricultural Economics, 29, 307-317.
- Hosmer, D., & Lemeshow, S. (2000). *Applied logistic regression* (Third Edition). New York: A Wiley– Interscience Publication.
- International Maize and Wheat Improvement Center (CIMMYT). (1993).*The adoption of agricultural technology. A guide for survey design.* Mexico, D. F.: CIMMYT: Author.
- Kaliba, R. M. A., Verkuijl, H., & Mwangi, W. (2000). Factors affecting adoption of improved maize seeds and use of inorganic fertilizer for

maize production in the intermediate and lowland zones of Tanzania. *Journal of Agricultural and Applied Economics*, *32*, 35-47.

- Ministry of Agriculture (MINAG). (2009). Agricultural Survey 2008. Maputo, Mozambique. Ministry of Agriculture: Author.
- Neupane, R. P., Sharma, K. R., & Thapa, G. B. (2002). Adoption of agroforestry in the hills of Nepal: a logistic regression analysis. *Journal of Agricultural Systems*, 72, 177-196.
- Pinstrup-Andersen, P., Pandya-Lorch, R. & Rosegrant, M.W. (1997). *The World Food*
- Situation: Recent Development, Emerging Issues, and Long-Term Prospects, Food Policy Report, 2020 vision. The International Food Policy Research Institute, Washington, DC, USA.
- Rotter, R., & van Keulen, H. (1997).Variations in response to fertilizer application in the tropics: II. Risks and opportunities for smallholders cultivating maize on Kenya's arable land. *Journal of Agricultural Systems*, 53, 69-95.
- Rogers, E. M., & Havens, E. (1961). The Impact of Demonstrations on Farmers' Attitudes Toward Fertilizers (Research Bulletin No. 898). Ohio Agricultural Experiment Station. Wooster, Ohio.
- Rogers, E. M. (2003). *Diffusion of Innovations* (Fourth Edition), New York: Free Press.
- SPSS Inc (2006). SPSS for Windows. Release 15.0.0. SPSS Inc., Chicago, USA.
- Thompson, P. T. (1992). *Perceptions and attitudes of farmers about a new fertilizer technology*. United States: International fertilizer development center.
- Visser, S. P., & Krosnick, J. A. (1998). Development strength over the life