Kansas Agricultural Experiment Station Research Reports

Volume 2
Issue 4 *Turfgrass Research*

Article 7

January 2016

Late Pre-Emergent Control of Annual Bluegrass with Flazasulfuron & Indaziflam

J. Reeves

Kansas State University, jakereeves@ksu.edu

J. Hoyle

Kansas State University, jahoye@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Horticulture Commons](#), and the [Weed Science Commons](#)

Recommended Citation

Reeves, J. and Hoyle, J. (2016) "Late Pre-Emergent Control of Annual Bluegrass with Flazasulfuron & Indaziflam," *Kansas Agricultural Experiment Station Research Reports*: Vol. 2: Iss. 4. <https://doi.org/10.4148/2378-5977.1213>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2016 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

TURFGRASS RESEARCH 2016

JULY 2016

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

K-State Research and Extension is an equal
opportunity provider and employer.

Late Pre-Emergent Control of Annual Bluegrass with Flazasulfuron & Indaziflam

Jake Reeves and Jared Hoyle

Summary. All applications of flazasulfuron or indaziflam resulted in acceptable control of annual bluegrass (*Poa annua*), only allowing <3% visual weed cover (VWC) in all treatments compared to the non-treated (26% VWC) up to 133 days after application (DAA).

Rationale. There are limited chemical options for controlling annual bluegrass (*Poa annua*) when the initial timing for pre-emergence is missed. Glyphosate is the standard chemical removal option, but resistance is becoming an issue in turfgrass systems. Flazasulfuron and indaziflam are new chemistries that provide pre- and post-emergent control, which could be a viable option for a late pre-emergent application.

Objectives. Examine efficacy of a late pre-emergent application of flazasulfuron and indaziflam for the control of annual bluegrass.

Study Description. Experiments were initiated in 2014 in Fort Riley, KS at Custer Hill Golf Course (CHGC). Research at CHGC was conducted on hybrid 'Midiron' bermudagrass [*Cynodon dactylon* L. Pers. × *C. transvaalensis* Burt-Davy]. Turfgrass was maintained at 1.0 in. All treatments were applied December 9, 2014. Individual treatments are shown in Table 1. Experimental design was a randomized complete block with four replications. Herbicides were applied to 5 × 5 ft plots in 374 L/ha water at 275 kPa with a CO₂ pressurized boom sprayer with XR8004VS flat-fan nozzles. Collected data included visual annual bluegrass cover (0-100%). Data were collected every 14 days after initiation (DAI). Means were separated using Fisher's Protected LSD ($P \leq 0.05$).

View all turfgrass research reports online at: <http://newprairiepress.org/kaesrr>

Results. All treatments, except for the non-treated, began showing post control of annual bluegrass at 83 DAA and continued to show pre-emergent control (Table 2) through trial completion at 133 DAA. All flazasulfuron and indaziflam treatments were not significantly different in the study as both controlled crabgrass pressure resulting in 3% or less annual bluegrass breakthrough. Both chemicals are viable annual bluegrass pre-emergent and post-emergent control options for turfgrass managers.

Table 1. Treatment regime for late pre-emergent control of *Poa annua* in bermudagrass with fluzasulfuron and indaziflam.

Treatment number	Chemistry	Rate
1	Fluzasulfuron ^z	1.5 oz/a
2	Flazasulfuron ^z	2.0 oz/a
3	Flazasulfuron ^z	2.5 oz/a
4	Flazasulfuron ^z	3.0 oz/a
5	Indaziflam	9.0 fl oz/a
6	Untreated	-

^z Treatments included a Non-Ionic Surfactant at 0.25% v/v.

Table 2. Means separation for visual weed cover with late pre-emergent control of *Poa annua* in bermudagrass with fluzasulfuron and indaziflam.

Treatment number	0 DAA ^z	83 DAA	133 DAA
1	2.5 a ^y	0.5 b	1.5 b
2	9.0 a	1.3 b	3.0 b
3	3.5 a	0.3 b	0.5 b
4	3.3 a	0.0 b	0.3 b
5	2.0 a	0.0 b	0.3 b
6	4.3 a	5.5 a	26.3 a

^z Days after application.

^y Means in columns for percent visual weed cover followed by the same letter are not significantly different according to Fisher's LSD at the .05 significance level.

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service