Inversion in the published genetic map of linkage group VII

E Kouzminova
University of Oregon

E U. Selker
University of Oregon

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Kouzminova, E., and E.U. Selker (1999) "Inversion in the published genetic map of linkage group VII," *Fungal Genetics Reports*: Vol. 46, Article 6. <https://doi.org/10.4148/1941-4765.1234>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Inversion in the published genetic map of linkage group VII

Abstract

In the course of cloning the *dim-2* gene of *Neurospora crassa* we found that the published map of LG VII has an inversion of a segment extending from *for* to *un-10K*. Direct physical mapping confirmed that the gene order in this region should be *wc-1*, *for*, *frq*, *oli*, *un-10*.

Inversion in the published genetic map of linkage group VII

Elena Kouzminova and Eric U. Selker* University of Oregon, Institute of Molecular Biology Eugene, Oregon

*corresponding author

In the course of cloning the *dim-2* gene of *Neurospora crassa* we found that the published map of LG VII has an inversion of a segment extending from *for* to *un-10*. Direct physical mapping confirmed that the gene order in this region should be *wc-1, for, frq, oli, un-10*.

In preparation to clone the *dim-2* gene (Foss *et al.* 1993 Science 262:1737-1741) of *Neurospora crassa*, we mapped this gene to the right arm of linkage group VII (LGVII) between *wc-1* and *arg-10*. To map the *dim-2* gene more precisely, three other crosses were performed (Table 1). We used strains with genetic markers *wc-1, frq* and *for* on LGVII and initially relied on a published genetic map (Figure 1a) to interpret our results. Data from the cross between *un-10* and *wc-1 dim-2* strains suggested that *dim-2* is left of *un-10* and between *wc-1* and *un-10*; about half of the recombinants (21/36) in the *wc-1* to *un-10* interval were *dim-2* (Table 1, cross 1). In two additional crosses in which one of the parents carried either a *for* or a *frq* mutation, however, we could not find *dim-2* recombinant progeny of the genotype *wc-1 dim-2 for* and *wc-1 dim-2 frq* (Table 1, crosses 2 and 3); other markers in these crosses segregated as expected from the published genetic map (data not shown). Therefore, *dim-2* appeared to be tightly linked to both *for* and *frq*. This result was unexpected since *frq* and *for* were reported to be to the right of *un-10* by approximately 9 and 12 map units, respectively (Figure 1a). To explain our genetic data we considered the possibility that the segment including *un-10, oli, frq* and *for* is inverted (Figure 1b). This orientation accounted for all our data and was subsequently confirmed by physical mapping (data not shown). Results of chromosomal walks from *wc-1* and *un-10* showed that *for* is proximal to *wc-1* followed by *frq, oli, dim-2* and *un-10* (Kouzminova and Selker, unpublished). Thus we suggest that the order of the genes *for, frq, oli* and *un-10* is as shown in Figure 1b.

Table 1. Map data.

Cross	Zygote genotype	Genotype and number of recombinant progeny analyzed for <i>dim-2</i>	Genotype and number of <i>dim-2</i> recombinant progeny
1 ^a	$\frac{+}{wc-1} \frac{+}{dim-2} \frac{un-10}{+}$	<i>wc-1⁺ un-10⁻</i> 36	<i>wc-1⁻ dim-2⁺ un-10⁻</i> 21 <i>wc-1⁺ dim-2⁻ un-10⁻</i> 15
2 ^b	$\frac{+}{wc-1} \frac{for (+)}{+ (dim-2)^d}$	random progeny 100	<i>wc-1 for dim-2⁺</i> 4 <i>wc-1⁻ for⁻ dim-2</i> 2
3 ^c	$\frac{+}{wc-1} \frac{frq::hph (+)}{+ (dim-2)^d}$	<i>wc-1 frq::hph</i> 10	<i>wc-1 frq::hph dim-2⁺</i> 10 <i>wc-1 frq::hph dim-2</i> 0

^aRecombinant progeny were selected on minimal medium at 34°C to select against *un-10* progeny. The *wc-1* mutation was scored under constant light at 34°C.

^b Progeny were scored without selection.

^c Progeny were selected on plates supplemented with hygromycin and scored for *wc-1* under constant light at 34°C.

^d Parentheses indicate that the position of *dim-2* relative to either *for* or *frq* could not be determined from the cross.

Figure 1. (a) Previous map and percentages of recombination between some genetic markers on LGVII (Perkins *et al.* 1982 Microbiol Reviews, 46:426-570). (b) Revised genetic map based on our findings. Genetic distance between *wc-1* and *for* was calculated from the cross 2 data (Table 1). the centromeric region is indicated by the circle on the left and genes in the disputed region are in bold.