

Kansas Agricultural Experiment Station Research Reports

Volume 2
Issue 6 *Kansas Fertilizer Research*


Article 7

January 2016

Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/Double-Crop Soybean Rotation

D. W. Sweeney
Kansas State University, dsweeney@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Sweeney, D. W. (2016) "Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/Double-Crop Soybean Rotation," *Kansas Agricultural Experiment Station Research Reports*: Vol. 2: Iss. 6. <https://doi.org/10.4148/2378-5977.1241>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright January 2016 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/Double-Crop Soybean Rotation

Abstract

(Abstract only. Link to: <http://newprairiepress.org/kaesrr/vol1/iss4/2/>) Article is nearly identical to [Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/Double-Crop Soybean Rotation](#) previously published in Southeast Agricultural Research Center 2015.

Overall in 2014, adding nitrogen (N) improved average wheat yields, but different N placement methods resulted in similar yields. Double-crop soybean yields were unaffected by tillage or the residual from N treatments that were applied to the previous wheat crop.

Keywords

Tillage, Nitrogen, Corn, Wheat, Soybean

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](#).

Kansas Agricultural Experiment Station Research Reports

Volume 1

Issue 4 Southeast Agricultural Research Center
Reports

Article 2


2015

Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/Double-Crop Soybean Rotation

D. W. Sweeney

Kansas State University, dsweeney@ksu.edu

Follow this and additional works at: <http://newprairiepress.org/kaesrr>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Sweeney, D. W. (2015) "Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/Double-Crop Soybean Rotation," *Kansas Agricultural Experiment Station Research Reports*: Vol. 1: Iss. 4. <http://dx.doi.org/10.4148/2378-5977.1052>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2015 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. K-State Research and Extension is an equal opportunity provider and employer.



Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/Double-Crop Soybean Rotation

Abstract

Overall in 2014, adding nitrogen (N) improved average wheat yields, but different N placement methods resulted in similar yields. Double-crop soybean yields were unaffected by tillage or the residual from N treatments that were applied to the previous wheat crop.

Keywords

tillage, nitrogen, placement, rotation

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/Double-Crop Soybean Rotation

D.W. Sweeney

Summary

Overall in 2014, adding nitrogen (N) improved average wheat yields, but different N placement methods resulted in similar yields. Double-crop soybean yields were unaffected by tillage or the residual from N treatments that were applied to the previous wheat crop.

Introduction

Many crop rotation systems are used in southeastern Kansas. This experiment is designed to determine the long-term effects of selected tillage and N fertilizer placement options on yields of short-season corn, wheat, and double-crop soybean in rotation.

Experimental Procedures

A split-plot design with four replications was initiated in 1983 with tillage system as the whole plot and N treatment as the subplot. In 2005, the rotation was changed to begin a short-season corn/wheat/double-crop soybean sequence. Use of three tillage systems (conventional, reduced, and no-till) continues in the same areas as during the previous 22 years. The conventional system consists of chiseling, disking, and field cultivation. Chiseling occurs in the fall preceding corn or wheat crops. The reduced-tillage system consists of disking and field cultivation prior to planting. Glyphosate is applied to the no-till areas prior to planting. The four N treatments for the crop are: no N (control), broadcast urea ammonium nitrate (UAN; 28% N) solution, dribble UAN solution, and knife UAN solution at 4 in. deep. The N rate for the corn crop grown in odd-numbered years is 125 lb/a. The N rate of 120 lb/a for wheat is split as 60 lb/a applied preplant as broadcast, dribble, or knifed UAN. All plots except for the controls are top-dressed in the spring with broadcast UAN at 60 lb/a N.

Results and Discussion

In 2014, wheat yields were low, averaging less than 30 bu/a (data not shown). Fertilizing with N increased wheat yield by approximately 70%, but preplant application method (broadcast, dribble, or knife) did not affect yields. Tillage had no effect on wheat yields. Average yield of soybean planted double-crop after wheat harvest exceeded 45 bu/a in 2014, but yield was not affected by tillage systems or the residual from N fertilizer treatments that were applied to the wheat.