Kansas Agricultural Experiment Station Research Reports

Volume 2
Issue 7 *Southwest Research-Extension Center
Reports*

Article 24

January 2016

Efficacy of Preemergence or Early Postemergence Weed Control with Keystone NXT, Hornet WDG, Atrazine, SureStart II, Lumax EZ, and Resicore

R. Currie
Kansas State University, rscurrie@ksu.edu

P. Geier
Kansas State University, pgeier@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Agronomy and Crop Sciences Commons](#), and the [Weed Science Commons](#)

Recommended Citation

Currie, R. and Geier, P. (2016) "Efficacy of Preemergence or Early Postemergence Weed Control with Keystone NXT, Hornet WDG, Atrazine, SureStart II, Lumax EZ, and Resicore," *Kansas Agricultural Experiment Station Research Reports*: Vol. 2: Iss. 7. <https://doi.org/10.4148/2378-5977.1269>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2016 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Efficacy of Preemergence or Early Postemergence Weed Control with Keystone NXT, Hornet WDG, Atrazine, SureStart II, Lumax EZ, and Resicore

R. Currie and P. Geier

Summary

Control of buffalobur was complete regardless of herbicide used. Velvetleaf and puncturevine control, although not perfect, was excellent by all herbicides. The premix of SureStart II (acetochlor + flumetsulam + clopyralid) with atrazine and Durango DMA (glyphosate) applied early postemergence and the preemergence herbicides Resicore (acetochlor + mesotrione + clopyralid) with atrazine and Lumax EZ (*S*-metolachlor + atrazine + mesotrione) provided excellent Palmer amaranth control. The early postemergence treatment of SureStart II plus atrazine and Durango DMA was the only treatment to provide excellent control of green foxtail.

Introduction

There are many products that provide an excellent foundation treatment that are supposed to be followed by a subsequent, postemergence application. It was the objective of this test to evaluate such compounds.

Procedures

An experiment at the Kansas State University Southwest Research-Extension Center near Garden City, KS determined the efficacy of preemergence and early postemergence herbicides in irrigated corn. Preemergence herbicides were applied June 2, 2015, with early postemergence treatments applied June 16, 2015. Corn was 5 to 7 inches tall when early postemergence treatments were made, and weeds were 1 to 2 inches tall. Herbicides were applied using a CO₂-compressed tractor-mounted or backpack sprayer delivering 20 gpa at 3 mph and 27 or 30 psi. Soil was a Ulysses silt loam with 1.4% organic matter, pH of 8.0, and cation exchange capacity of 18.4. Plots were 10 by 35 feet and arranged in a randomized complete block with four replications. Visual weed control was determined July 28, 2015, which was 42 days after early postemergence application (42 DAPT). Grain yields were determined October 14, 2015 by mechanically harvesting the center two rows of each plot and adjusting weights to 15.5% moisture.

Results and Discussion

Control of buffalobur was complete at 42 DAPT regardless of herbicide. Velvetleaf and puncturevine control was 99 to 100 and 95 to 99%, respectively, at the same date by all herbicides. The premix of SureStart II with atrazine and Durango DMA applied early postemergence and the preemergence herbicides Resicore with atrazine and Lumax EZ controlled Palmer amaranth 96 to 99%. The early postemergence treatment of SureStart II plus atrazine and Durango DMA was the only treatment to control green foxtail more than 95% at 42 DAPT. Grain yields did not differ between herbicide treatments or the untreated controls.

Table 1. Application information.

Application timing	Preemergence	Postemergence
Application date	June 2, 2015	June 16, 2015
Air temperature (°F)	75	82
Relative humidity (%)	62	44
Soil temperature (°F)	67	76
Wind speed (mph)	5 to 8	3 to 5
Wind direction	North	North
Soil moisture	Good	Fair

Table 2. Efficacy of preemergence or early postemergence herbicides in irrigated corn.

Treatment ^a	Rate	Timing ^b	42 days after early postemergence application					Yield
			Palmer amaranth	Velvetleaf	Puncturevine	Buffalobur	Green foxtail	
			----- % visual -----					bu/a
Keystone NXT	2.4 qt	PRE	95	100	95	100	91	134.0
Hornet WDG	4.0 oz	PRE						
SureStart II	2.0 pt	PRE	90	100	98	100	84	122.1
Atrazine	1.0 qt	PRE						
SureStart II	2.0 pt	EPOST	99	100	99	100	97	141.9
Atrazine	1.0 qt	EPOST						
Durango DMA	32 oz	EPOST						
AMS	2.5 %	EPOST						
Resicore	2.5 qt	PRE	99	99	96	100	89	124.1
Atrazine	1.0 qt	PRE						
Resicore	1.5 qt	PRE	96	100	96	100	85	135.6
Atrazine	1.0 qt	PRE						
Lumax EZ	2.7 qt	PRE	99	100	98	100	85	148.7
Untreated			0	0	0	0	0	130.2
LSD (0.05)			4.1	1.4	5.5	NS	4.6	NS

^a AMS is ammonium sulfate.

^b PRE is preemergence, EPOST is early postemergence.

Figure 1. Untreated control.

Figure 2. SureStart II 2 pt + atrazine 1 qt + Durango DMA 32 oz + AMS early postemergence, 50 days after early postemergence application.

Figure 3. Resicore 2.5 qt + atrazine 1 qt preemergence, 64 days after preemergence application.

Figure 4. Resicore 1.5 qt + atrazine 1 qt preemergence, 64 days after preemergence application.

Figure 5. Lumax EZ 2.7 qt preemergence, 64 days after preemergence application.