

## Mind the buffering capacity of citric acid

S. R. Nozawa  
*FFCLRP-USP*

I. C. Rigoli  
*FFCLRP-USP*

G. Thedei Jr.  
*FFCLRP-USP*

*See next page for additional authors*

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Nozawa, S. R., I.C. Rigoli, G. Thedei, and A. Rossi (1995) "Mind the buffering capacity of citric acid," *Fungal Genetics Reports*: Vol. 42, Article 16. <https://doi.org/10.4148/1941-4765.1346>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## Mind the buffering capacity of citric acid

### Abstract

Many microbial cultures are buffered with citric acid over a pH range of 2.5 to 7.0 since the pKa values for this triprotic acid are 3.13, 4.76 and 6.40, as shown in The Merck Index (pp 330-331, 10th Edition, Martha Windholz, ed.). However, the information about the buffering range of this weak acid is controversial since the pKa3 value may be 5.40, as specified in Day and Underwood (Quantitative Analysis, 6th Edition, 1991, p. 662. New Jersey: Prentice Hall). With this in mind, we determined the pKa values of citric acid at concentrations ranging from 5 mM to 50 mM, concentrations which are the most employed in buffers for the culture of many microorganisms.

### Authors

S. R. Nozawa, I. C. Rigoli, G. Thedei Jr., and A. Rossi


## Mind the buffering capacity of citric acid

*S.R. Nozawa, I.C. Rigoli, G. Thedei Jr. and A. Rossi - Departamento de Química, FFCLRP-USP, 14040-901 Ribeirão Preto, SP, Brazil.*

Many microbial cultures are buffered with citric acid over a pH range of 2.5 to 7.0 since the pKa values for this triprotic acid are 3.13, 4.76 and 6.40, as shown in The Merck Index (pp 330-331, 10th Edition, Martha Windholz, ed.). However, the information about the buffering range of this weak acid is controversial since the pKa3 value may be 5.40, as specified in Day and Underwood (Quantitative Analysis, 6th Edition, 1991, p. 662. New Jersey: Prentice Hall). With this in mind, we determined the pKa values of citric acid at concentrations ranging from 5 mM to 50 mM, concentrations which are the most employed in buffers for the culture of many microorganisms.

The experimental values obtained were 3.10, 4.80 and 5.40 for pKa1, pKa2 and pKa3, respectively (Figure 1A). Furthermore, citric acid does not seem to be a good buffer even considering pKa values ranging from 3.1 to 5.4 (Fig. 1A). This is because the values of the three dissociation constants are too close to permit distinction of the three proton receptor phases. The titration curve for sodium citrate with HCl (Figure 1B) confirmed the observation that citric acid has no buffering capacity at pH higher than 6.2. Figure 1A also shows the titration curve of 20 mM 2-[N- Morpholino]ethanesulfonic acid (MES) with 100 mM sodium hydroxide, clearly confirming its buffering capacity at pH 6.1, which corresponds to its pKa value.

This work was supported by a FAPESP grant to A. Rossi.



**Figure 1.** (A) Titration curves of 5 mM citric acid (50 ml) with 100 mM NaOH (p) and 20 mM MES (50 ml) with 100 mM NaOH (p). (B) Titration curve of 20 mM sodium citrate (50 ml) with 100 mM HCl. All determinations were made at room temperature and the pH was measured with a Mettrom Herisau E-510 pH meter.