Some observations concerning *sp* and *ure-2* in *Neurospora*

F. J. Bowring
Flinders University

David E. A. Catcheside
Flinders University

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Bowring, F. J., and D.E. Catcheside (1994) "Some observations concerning *sp* and *ure-2* in *Neurospora*," *Fungal Genetics Reports*: Vol. 41, Article 3. <https://doi.org/10.4148/1941-4765.1365>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Some observations concerning sp and ure-2 in Neurospora

Abstract

We are currently seeking rec-2 which is proximal to am on linkage group V and are gleaning incidental information on genes in this region and elsewhere. So far, the following is apparent: sp is recessive: each of 15 heterokaryons forced between his-3 K874; sp B132 and am1 32213 strains by growth on Vogel's + glycine medium had wild-type morphology.

Some observations concerning sp and ure-2 in Neurospora

F.J. Bowring & D.E.A. Catcheside - School of Biological Sciences, Flinders University, G.P.O. Box 2100, Adelaide, South Australia, 5001.

We are currently seeking rec-2 which is proximal to am on linkage group V and are gleaning incidental information on genes in this region and elsewhere. So far, the following is apparent:

sp is recessive: each of 15 heterokaryons forced between his-3 K874; sp B132 and am1 32213 strains by growth on Vogel's + glycine medium had wild-type morphology.

The gene order sp, ure-2, am inferred from two point crosses is confirmed by three point data: FGSC 3809 (a; ure-2 D74) was crossed to F11089 (A; sp B132, am B501). 320 random spores were isolated, 79% germinated.

Zygote genotype

and

recombination percent			Parental combinations			Singles Region I			Singles Region II
sp	+	am	sp + am	+ ure-2 +	sp ure-2 +	+ + am	sp + +	+	
ure-2	am								
+	ure-2	+	135	92	9	11	5		
0									
7.9	2.0								

A gene in FGSC 4299 interferes with Kolmark's urease test: While seeking to order sp, ure-2 and am, we set up a cross between FGSC 4299 (a; ure-2 47, am 32213) and T9043 (A; his-3 K874; sp B132). Of 320 random spores, 92% germinated and were scored for urease activity by dabbing conidia onto paper soaked in urea and a pH indicator to detect ammonia production (Kolmark 1969, Mutation Res. 8:51-63). By this test, T9043 was urease positive and FGSC 4299 urease negative. However, only 23% of the progeny were urease positive, suggesting the segregation of a second mutation unlinked to ure-2. There is preliminary evidence that this additional locus could be linked to his-3.