

# Kansas Agricultural Experiment Station Research Reports

---

Volume 3

Issue 2 Southeast Agricultural Research Center  
Reports

Article 10

---

January 2017

## Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/ Double-Crop Soybean Rotation

D. W. Sweeney

Kansas State University, dsweeney@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Agronomy and Crop Sciences Commons](#)

---

### Recommended Citation

Sweeney, D. W. (2017) "Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/ Double-Crop Soybean Rotation," *Kansas Agricultural Experiment Station Research Reports*: Vol. 3: Iss. 2. <https://doi.org/10.4148/2378-5977.1379>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright January 2017 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

# Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/ Wheat/ Double-Crop Soybean Rotation

## **Abstract**

In 2016, adding nitrogen (N) greatly improved average wheat yields, but the response to tillage and different N placement methods was minimal. Double-crop soybean yields were unaffected by tillage or the residual from N treatments that were applied to the previous wheat crop.

## **Keywords**

tillage, no-till, nitrogen placement, corn, wheat, soybean

## **Creative Commons License**



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

## **Tillage and Nitrogen Placement Effects on Yields in a Short-Season Corn/Wheat/ Double-Crop Soybean Rotation**

*D.W. Sweeney*

### **Summary**

In 2016, adding nitrogen (N) greatly improved average wheat yields, but the response to tillage and different N placement methods was minimal. Double-crop soybean yields were unaffected by tillage or the residual from N treatments that were applied to the previous wheat crop.

### **Introduction**

Many crop rotation systems are used in southeastern Kansas. This experiment is designed to determine the long-term effect of selected tillage and N fertilizer placement options on yields of short-season corn, wheat, and double-crop soybean in rotation.

### **Experimental Procedures**

A split-plot design with four replications was initiated in 1983 with tillage system as the whole plot and N treatment as the subplot. In 2005, the rotation was changed to begin a short-season corn/wheat/double-crop soybean sequence. Use of three tillage systems (conventional, reduced, and no-till) continues in the same areas as the previous 22 years. The conventional system consists of chiseling, disking, and field cultivation. Chiseling occurs in the fall preceding corn or wheat crops. The reduced-tillage system consists of disking and field cultivation prior to planting. Glyphosate is applied to the no-till areas prior to planting. The four N treatments for the crop are: no-N (control), broadcast urea-ammonium nitrate (UAN; 28% N) solution, dribble UAN solution, and knife UAN solution at 4 inches deep. The N rate for the corn crop grown in odd-numbered years is 125 lb/a. The N rate of 120 lb/a for wheat is split as 60 lb/a applied preplant as broadcast, dribble, or knifed UAN. All plots except for the no-N controls are top-dressed in the spring with broadcast UAN at 60 lb/a N.

### **Results and Discussion**

In 2016, conventional tillage resulted in 2 bu/a greater yield than with no-till (Table 1). Overall, fertilizing with N quadrupled wheat yield, but preplant application method (broadcast, dribble, or knife) did not affect yields. Average yield of soybean planted doublecrop after wheat harvest was nearly 40 bu/a in 2016, but was not affected by tillage systems or the residual from N fertilizer treatments that were applied to the wheat.

**Table 1. Effect of tillage and fall N fertilization on yield of wheat and following double-crop soybean in 2016**

Treatment	Wheat yield	Double-crop soybean yield
	----- bu/a -----	
Tillage		
Conventional	37.8	39.0
Reduced	36.8	39.9
No-till	35.8	39.7
LSD (0.05)	1.1	NS
N Fertilization		
No-N control	9.8	40.0
Broadcast UAN†	45.5	39.4
Dribble UAN	45.1	39.7
Knife UAN	46.4	39.0
LSD (0.05)	1.4	NS

†UAN: urea-ammonium nitration solution, 28% N.