

# Kansas Agricultural Experiment Station Research Reports

---

Volume 3

Issue 2 Southeast Agricultural Research Center  
Reports

Article 11

---

January 2017

## Timing of Side-Dress Applications of Nitrogen for Corn in Conventional and No-Till Systems

D. W. Sweeney

Kansas State University, [dsweeney@ksu.edu](mailto:dsweeney@ksu.edu)

D. E. Shoup

Kansas State University, [dshoup@ksu.edu](mailto:dshoup@ksu.edu)

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Agronomy and Crop Sciences Commons](#)

---

### Recommended Citation

Sweeney, D. W. and Shoup, D. E. (2017) "Timing of Side-Dress Applications of Nitrogen for Corn in Conventional and No-Till Systems," *Kansas Agricultural Experiment Station Research Reports*: Vol. 3: Iss. 2. <https://doi.org/10.4148/2378-5977.1380>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright January 2017 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

# Timing of Side-Dress Applications of Nitrogen for Corn in Conventional and No-Till Systems

## Abstract

Corn yields were affected by tillage and nitrogen (N) side-dress options in 2016. Corn yields were 12% greater with conventional tillage than with no-till. Side-dress applications of N at V10 resulted in greater corn yield than side-dress N applications at V6.

## Keywords

nitrogen, side-dress, corn, tillage, no-till

## Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

## **Timing of Side-Dress Applications of Nitrogen for Corn in Conventional and No-Till Systems**

*D.W. Sweeney and D. Shoup*

### **Summary**

Corn yields were affected by tillage and nitrogen (N) side-dress options in 2016. Corn yields were 12% greater with conventional tillage than with no-till. Side-dress applications of N at V10 resulted in greater corn yield than side-dress N applications at V6.

### **Introduction**

Environmental conditions vary widely in the spring in southeastern Kansas. As a result, much of the N applied prior to corn planting may be lost before the time of maximum plant N uptake. Side-dress or split applications to provide N during rapid growth periods may improve N use efficiency while reducing potential losses to the environment. The objective of this study was to determine the effect of timing of side-dress N fertilization compared with preplant N applications for corn grown on a claypan soil.

### **Experimental Procedures**

The experiment was established in spring 2015 on a Parsons silt loam soil at the Parsons unit of the Kansas State University Southeast Agricultural Research Center. The experiment was a split-plot arrangement of a randomized complete block design with four blocks (replications). Whole plot tillage treatments were conventional tillage (chisel, disk, and field cultivate) and no-till. Sub-plot nitrogen treatments were six preplant/side-dress N application combinations that include 1) a no-N control, 2) 150 lb N/a applied preplant, 3) 100 lb N/a applied preplant with 50 lb N/a applied at the V6 (six-leaf) growth stage, 4) 100 lb N/a applied preplant with 50 lb N/a applied at the V10 (ten-leaf) growth stage, 5) 150 lb N/a applied preplant with 50 lb N/a applied at the V6 growth stage, and 6) 150 lb N/a applied preplant with 50 lb N/a applied at the V10 growth stage. The N source for all treatments was liquid urea-ammonium nitrate (28% N) fertilizer. Preplant N fertilizer was applied on April 4, 2016, side-dress N at V6 on May 22, 2016, and side-dress N at V10 on June 6, 2016 to appropriate plots. Corn was planted on April 5 and harvested on August 29, 2016.

### **Results and Discussion**

In 2016, corn yielded 12 bu/a more with conventional tillage than with no-till (Table 1). Even though yield components were not significantly affected by tillage, the com-

bined trend for greater stand, kernel weight, and kernels/ear likely accounted for the yield response to tillage. Adding N fertilizer more than tripled yields obtained in the no-N control. Applying 100 lb N/a preplant followed by 50 lb N/a at the V6 growth state did not improve yields above that obtained with all 150 lb N/a applied preplant. However, delaying the 50 lb N/a side-dress application to the V10 stage improved yield by 8.4 bu/a compared to all N preplant. A similar increase in yield was found by delaying N side-dress to the V10 stage instead of the V6 stage when adding 50 lb N/a extra to a 150 lb N/a preplant application. These effects of N timing on corn yield in 2016 appeared to be related to responses in kernel weight and kernels/ear.

**Table 1. Tillage and N side-dress application effects on yield and yield components of corn in 2016**

Treatment	Yield	Stand	Kernel weight	Ears/plant	Kernels/ear
	bu/a	#/a	mg		
Tillage (T) <sup>1</sup>					
Conventional	110.5	22600	231	0.99	529
No-till	98.3	21700	222	0.99	508
LSD (0.05)	6.3	NS	NS	NS	NS
Nitrogen Timing (N) <sup>2</sup>					
No-N control	34.4	22000	172	0.99	235
150 PP	111.2	22000	219	0.99	592
100 PP/50 V6	112.0	22200	234	0.99	553
100 PP/50 V10	119.6	22400	240	0.99	570
150 PP/50 V6	118.9	22200	240	1.00	569
150 PP/50 V10	130.3	22100	255	0.99	594
LSD (0.05)	7.0	NS	15	NS	34

<sup>1</sup>Conventional tillage: chisel, disk, and field cultivate.

<sup>2</sup>Nitrogen treatments: Control, no N fertilizer; 150 PP, 150 lb N/a applied preplant with no side-dress N; 100 PP/50 V6, 100 lb N/a applied preplant with 50 lb N/a side-dress applied at V6 (six-leaf) growth stage; 100 PP/50 V10, 100 lb N/a applied preplant with 50 lb N/a side-dress applied at V10 (ten-leaf) growth stage; 150 PP/50 V6, 150 lb N/a applied preplant with 50 lb N/a side-dress applied at V6 growth stage; and 150 PP/50 V10, 150 lb N/a applied preplant with 50 lb N/a side-dress applied at V10 growth stage.