Production of mutants of *Gaeumannomyces graminis* var. *tritici* and var. *avenae* by 4- nitroquinolene-oxide treatment of protoplasts.

P. Bowyer
The Sainsbury Laboratory

R. Musker
University of Birmingham

A. E. Osbourn
The Sainsbury Laboratory

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Bowyer, P., R. Musker, A.E. Osbourn, B. Clarke, C. Caten, and M.J. Daniels (1992) "Production of mutants of *Gaeumannomyces graminis* var. *tritici* and var. *avenae* by 4- nitroquinolene-oxide treatment of protoplasts.," *Fungal Genetics Reports*: Vol. 39, Article 3. <https://doi.org/10.4148/1941-4765.1429>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Production of mutants of *Gaeumannomyces graminis* var. *tritici* and var. *avenae* by 4- nitroquinolene-oxide treatment of protoplasts.

Abstract

The ascomycete fungus *Gaeumannomyces graminis* is the causative agent of take-all disease of cereals. Much information about the physiology and pathology of this organism has been generated (Asher and Shipton (Eds.) "Biology and Control of Take-All", Academic Press, 1981), but genetic studies such as the production of mutants have been hindered by problems in obtaining viable propagules suitable for mutagenesis (Blanch et al. 1981. Trans. Brit. Mycol. Soc. 77:391-399). The fungus is homothallic but many strains cannot be induced to form perithecia in culture and even the fertile strains produce insufficient numbers of ascospores for use in mutagenesis. It is, however, possible to produce and regenerate large numbers of protoplasts and Rochefrette et al.

Authors

P. Bowyer, R. Musker, A. E. Osbourn, B. Clarke, C. Caten, and M. J. Daniels

Production of mutants of *Gaeumannomyces graminis* var. *tritici* and var. *avenae* by 4-nitroquinolene-oxide treatment of protoplasts.

P. Bowyer, R. Musker*, A.E. Osbourn, B. Clarke, C. Caten* and M.J. Daniels - The Sainsbury Laboratory, The Norwich Research Park, Colney, Norwich, NR4 7UH, U.K. *Dept. of Genetics, University of Birmingham, Birmingham B15 2TT, U.K.

The ascomycete fungus *Gaeumannomyces graminis* is the causative agent of take-all disease of cereals. Much information about the physiology and pathology of this organism has been generated (Asher and Shipton (Eds.) "Biology and Control of Take-All", Academic Press, 1981), but genetic studies such as the production of mutants have been hindered by problems in obtaining viable propagules suitable for mutagenesis (Blanch et al. 1981. Trans. Brit. Mycol. Soc. 77:391-399). The fungus is homothallic but many strains cannot be induced to form perithecia in culture and even the fertile strains produce insufficient numbers of ascospores for use in mutagenesis. It is, however, possible to produce and regenerate large numbers of protoplasts and Rochefrette et al. (1979. Ann. Phytopath. 11:43-51) reported the mutagenesis of *G. graminis* protoplasts using nitrosoguanidine (NTG). Here we report the mutagenesis of protoplasts by 4-nitroquinolene oxide (NQO), reportedly a safer and more disposable mutagen than NTG (Bal et al. 1977. Mutat. Res. 56:153-156).

Protoplasts were prepared by standard methods (e.g. Ballance and Turner 1985. Gene 36:321-331) with the following modifications. Protoplasts were produced from washed 2-day old liquid cultures (potato dextrose broth) by shaking (100 rpm) in 0.6 M KCl, 5 mg Novozyme 234/ml for 4 h at 25 C. Mycelial debris was filtered out with four layers of Miracloth (Calbiochem Inc, USA) and the resulting suspension pelleted by centrifugation at 3000 x g. The supernatant was removed and the protoplasts were washed by resuspending them in 0.6 M KCl and repelleting them three times.

Approximately 108 protoplasts were incubated for 1 h at room temperatures in 5 ml 0.6 M KCl containing NQO (1 ug/ml Sigma) from a 10 mg/ml stock in acetone (this treatment was predicted to kill approximately 99% of protoplasts; see Figure 1). After treatment, the NQO was neutralized by the addition of an equal volume of 5% sodium thiosulphate in 0.6 M KCl, protoplasts were washed three times, as previously described, in protoplast regeneration medium (per liter: NaNO₃, 4 g; KCl, 44g; KH₂PO₄, 1.52 g; MgSO₄.7H₂O, 1 g; glucose, 10 g; yeast extract, 0.5 g; casamino acids, 0.2 g; FeSO₄.7H₂O, 1 mg; ZnSO₄.7H₂O, 10 mg; CuSO₄.5H₂O, 1 mg; Na₂B₄O₇.10H₂O, 1 mg; and (NH₄)₆Mo₇O₂₄.4H₂O, 50 mg; adjusted to pH 6.5) and added to 5 ml molten regeneration medium containing 0.5% agar and the appropriate inhibitor, which had been cooled to 48 C. The protoplast/agar mix was then overlaid onto protoplast regeneration agar plates.

The proportion of protoplasts containing nuclei was estimated in a separate experiment by bis-benzimide staining and visualization under UV light irradiation. Approximately 16% contained one nucleus and 6% contained two or more nuclei. Regeneration frequencies for protoplasts of both *avenae* and *tritici* varieties were normally between 5 and 10%. Plates were incubated at 22 C for 1-3 weeks and fast growing colonies were picked and retested on the appropriate selective media. Rates of transformation of protoplasts to phleomycin resistance can be enhanced by regenerating protoplasts for a few days before adding the antibiotic to the plates. In our hands, however, it was found that regeneration of mutated protoplasts before the addition of inhibitor resulted in high levels of background growth which interfered with selection of resistant colonies. The following mutants resistant to these inhibitors were selected:

<u>Fungus</u>	<u>Selection</u>	<u>Mutants/10⁸ protoplasts</u> <u>+ NQO</u>	<u>Mutants/10⁸ protoplasts</u> <u>-NQO</u>
<u>G. graminis</u> var. <u>tritici</u> (Strain T5)	25 mM fluoroacetate	22	1
	10 mM canavanine*	1	0
	100 mg ethionine/ml*	3	0
	2 mM deoxyglucose	17	2
	1 mg fluororotic acid/ml*	1	0
	2 mg carboxin/ml	16	0
	2 µg benomyl/ml	5	0
<u>G. graminis</u> var. <u>avenae</u> (Strain A3)	25 mM fluoroacetate	0	0
	100 mg ethionine/ml*	1	0
	2 mM deoxyglucose	112	6
	1 mg fluororotic acid/ml*	0	0

* The figures here represent the mean from two or more experiments

1280 *G. graminis* var. *tritici* colonies surviving NQO treatment were screened for auxotrophy; 1 methionine and 1 arginine auxotroph were isolated together with a mutant requiring NH₂⁺ as a nitrogen source. In addition, 6 color and 5 morphological mutants were isolated from the *tritici* isolate, and 2 color and one morphological mutant were isolated from the *avenae* isolate.

In conclusion, this method has provided us with an effective means of mutagenizing this organism: 195 mutants were produced using NQO mutagenesis compared with only 7 mutants reported for NTG mutagenesis (Rocheffette et al.) although differences in selection procedures and strains used may render this comparison invalid. No direct comparison of NTG and NQO mutagenesis rates was done. Work is in progress to characterize the mutants so far isolated.

Figure 1. Protoplast survival after NQO treatment for 1 hour (Regeneration frequency is expressed as a percentage of untreated protoplast regeneration - typically 5-10% of untreated protoplasts regenerated)