

## Construction of testers for reversion assay.

M. Kimura

N. Osawa

C. Ishii

H. Inoue

Follow this and additional works at: <http://newprairiepress.org/fgr>

---

### Recommended Citation

Kimura, M., N. Osawa, C. Ishii, and H. Inoue (1986) "Construction of testers for reversion assay," *Fungal Genetics Reports*: Vol. 33, Article 6. <https://doi.org/10.4148/1941-4765.1581>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

# Construction of testers for reversion assay.

## **Abstract**

Construction of testers for reversion assay.

## **Creative Commons License**



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Kimura, M., N. Osawa, C. Ishii

and H. Inoue

Construction of testers for  
reversion assay.

The ad-3 forward mutation assay system developed by de Serres and coworkers has been used in our laboratory for research on DNA repair end mutagenesis. This system gives much information on mutagenesis, but is expensive to use and is labor intensive. Therefore, we desired to develop a reversion assay system which is more convenient.

Dr. T. Ishikawa (University of Tokyo) had isolated and kept many ad-8 mutants in his laboratory. We screened revertability of seventy eight ad-8 mutants by the spot test method of Ong (1978, Mutation Res. 53:297-308) and selected two mutants on the basis of reversion specificity. One of them, ad-8(E193), was highly reverted by MNNG but not by ICR170 and the other, ad-8(E146) was highly reverted by ICR170 but not by MNNG. Each of these mutants was back-crossed to standard strain 74-OR31-14a(al-2 cot-1 pan-2 a) twice and two testers (T26 and T28) were constructed. Their genotypes are shown in Table 1. Responses of both testers to several mutagens are shown in Table 2. These data are from the plate test method of Ong. The results indicate that T26 responds to frameshift mutagens and T28 to basepair substitution mutagens. These testers may be useful along with Ong's strains N23 and N26, which had been constructed for the assay of environmental mutagens and are useful for quantitative and qualitative comparison of DNA repair ability and mutagen specificity.

Table 1. Tester strains

Tester	Strain #	FGSC #	Genotype (allele)			
T26	C3-T26-14a	5071	<u>ad-8(E146)</u>	<u>al-2(Y112M38)</u>	<u>cot-1(C102(t))</u>	<u>pan-2(Y387-15.7)</u>
T28	C3-T28-39a	5071	<u>ad-8(E193)</u>	<u>al-2(Y112M38)</u>	<u>cot-1(C102(t))</u>	<u>pan-2(Y387-15.7)</u>

Table 2. Induction of reverse mutations with plate test in T26 and T28

Muta- gen	Concen- tration	No. of revertants per plate		Muta- gen	Concen- tration	No. of revertants per Plate	
		T26	T28			T26	T28
MNNG	0 (ug/plate)	1	2	EMS	0 (ul/plate)	0	<b>1</b>
	1	0	60		10	0	<b>51</b>
	5	1	1070		20	2	290
	10	3	>1000				
ICR170	0 (ug/plate)	1	1	4NQO	0 (ug/plate)	0	2
	<b>5</b>	<b>84</b>	<b>3</b>		0.2	2	13
	10	130	3		<b>0.5</b>	<b>4</b>	<b>32</b>
	20	145	1	UV	0 (erg/mm2)	0	1
MMS	0 (ul/plate)	0	1		<b>1000</b>	<b>2</b>	<b>15</b>
	1	10	29	2000		8	23
	<b>2</b>	<b>2</b>	<b>8</b>	4000		11	20

Spontaneous reversion of T26 and T28 was 0.5 and 1.5 per plate, respectively, on average