

DNA polymerase in temperature-sensitive mutants of *Neurospora*

A. L. Schroeder

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Schroeder, A. L. (1984) "DNA polymerase in temperature-sensitive mutants of *Neurospora*," *Fungal Genetics Reports*: Vol. 31, Article 21. <https://doi.org/10.4148/1941-4765.1615>

This One-Liner is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

DNA polymerase in temperature-sensitive mutants of *Neurospora*

Abstract

DNA polymerase in temperature-sensitive mutants of *Neurospora*

ONE-LINER

Schroeder, A. L.

DNA Polymerase in temperature-sensitive mutants of Neurospora.

at 37° C for 5-60 min. None of the strains showed temperature-sensitivity of DNA polymerase activity.
Program in Genetics and Cell Biology, Washington State University, Pullman, WA 99164-4350.

I have tested DNA polymerase activity in the temperature-sensitive mutant un-3 and 11 temperature-sensitive mutants described by Stadler (Neurospora Newsletter 28: 18) which stop DNA synthesis soon after being shifted to the non-permissive temperature, 37° C (Stadler, personal communication). These

are 3B, 47D, 67, 6V, 145, 121, 19, 6B, 58E, 16t, 134C. Assays of crude extracts were at 37° C for 30 min after preincubation