

## Germination of microconidia from selected Neurospora strains

S. C. Hedman

S. Vanderschmidt

Follow this and additional works at: <http://newprairiepress.org/fgr>

---

### Recommended Citation

Hedman, S. C., and S. Vanderschmidt (1981) "Germination of microconidia from selected Neurospora strains," *Fungal Genetics Reports*: Vol. 28, Article 9. <https://doi.org/10.4148/1941-4765.1654>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

# Germination of microconidia from selected Neurospora strains

## **Abstract**

Germination of microconidia from selected Neurospora strains

## **Creative Commons License**



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Germination of microconidia from  
selected *Neurospora* strains.

These latter strains are similar to the F<sub>2</sub> lines of pe, fl previously described by Mnkres (1977 *Neurospora* Newsl. 24: 9).

Cultures were grown on either minimal medium (1X Vogel's minimal, 2% sucrose, 1.5% agar or on complete medium (1X Vogel's minimal, 2% sucrose, 0.1% yeast extract, 0.1% malt extract, 1.5% agar) in 125 ml Erlenmeyer flasks at 30°C under constant illumination. This complete medium was similar to that described by Baylis and DeBusk (1965 *Neurospora* Newsl. 7: 7) except that liver extract was omitted. All of the strains yielded almost exclusively microconidia as verified by scanning electron microscopy of hyphae. These microconidia were generally ellipsoidal with a major axis length of  $2.1 \pm 0.3 \mu\text{m}$ . This was in contrast to macroconidia obtained from STA4 (FGSC #262) which had a major axis length of  $5.8 \pm 1.8 \mu\text{m}$ . The number of microconidia produced by cultures grown on complete medium was five to seven times greater than that from cultures grown on minimal medium.

Microconidial suspensions were prepared by vigorously shaking each culture with 15 ml of water followed by filtration through two layers of cheesecloth; filtered microconidial suspensions were examined with a hemocytometer to determine concentrations of microconidia and to verify the absence of hyphal fragments. Germination percentages of microconidia were ascertained by suspending a diluted aliquot in 45°C molten agar and overlaying upon sorbose medium (1X Vogel's minimal, 1% sorbose, 0.05% fructose, 0.05% glucose, 1.5% agar). Microconidia from the pe, fl strain and the pe, fl derived strains of Mnkres produced colonies within three days of incubation at 30°C; colonies from the fl; dn strains were not observed until after four to five days of incubation. We observed no significant differences in percent germination between microconidia derived from cultures grown on minimal medium and those derived from cultures grown on complete medium. Using these same techniques, macroconidia from STA4 showed greater than 80% germination.

TABLE 1\*

Strain	Total Microconidial Yield	Percent Germinating Microconidia
<u>pe, fl</u> (#569)	$1.65 \times 10^9$	26%
<u>pe, fl</u> (3073)	$6.75 \times 10^8$	34%
F2 <u>pe, fl</u> (14-8a)	$1.45 \times 10^8$	26%
<u>pe, fl</u> (14.12A)	$8.40 \times 10^8$	24%
<u>pe, fl</u> (14-13a)	$1.02 \times 10^9$	39%
<u>pe, fl</u> (17-3a)	$7.60 \times 10^7$	35%
<u>pe, fl</u> (17-8A)	$6.80 \times 10^8$	34%
<u>fl; dn</u> (#3518)	$1.32 \times 10^9$	6%

\*Table 1 summarizes the data for cultures grown on complete medium

may be highly fertile as a female parent (Perkins 1979 *Neurospora* Newsl. 26: 9) its total usefulness may be diminished due to the low viability of its microconidia.

In addition, both microconidial yield and viability were significantly reduced when cultures were grown at 35°C. Also culturing beyond seven days at 30°C increased total microconidial yields, gave reduced microconidia germination from pe, fl strains, but no significant change in percentage microconidia germination of fl; dn. The technical assistance of Ms. Doria Harris is gratefully acknowledged. - - - Department of Biology, University of Minnesota, Duluth, Minnesota 55812.

In contrast to macroconidia, microconidia are generally uninucleate and possess fewer mitochondria (Lowry et al. 1967 *J. Bacteriology* 94: 1757). During our studies of mitochondrial inheritance, we have investigated various parameters affecting microconidial production and the viability of pe, fl (FGSC #569 and #3073), fl; dn (FGSC #3518), and several microconidiating strains kindly provided us by Ken Mnkres (14-8a, 14-12A, 14-13a, 17-3a, and 17-8A).

Regarding the data in Table 1, the following points are noted: (1) The viabilities of microconidia from certain pe, fl strains were greater than those reported by Barratt (1964 *Neurospora* Newsl. 6: 6); (2) The viabilities of microconidia of the pe, fl-derived strains of Mnkres were considerably lower than his reported value of 85 - 95% (1977 *Neurospora* Newsl. 24: 9); and (3) While fl; dn