Surfactant resistance in natural *Neurospora* populations.

A. Al-Saqr

B. R. Smith

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Al-Saqr, A., and B.R. Smith (1980) "Surfactant resistance in natural *Neurospora* populations.," *Fungal Genetics Reports*: Vol. 27, Article 2. <https://doi.org/10.4148/1941-4765.1665>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Surfactant resistance in natural *Neurospora* populations.

Abstract

Surfactant resistance in natural *Neurospora* populations.

Surfactant resistance in natural
Neurospora populations.

Seven wild type strains collected in different countries were obtained from the Fungal Genetics Stock Center and screened for resistance to four surfactants (Table 1). Five of the wild type strains show resistance to dequalinium chloride (DC), cetyltrimethyl ammonium bromide (CTAB), and benzalkonium chloride (BAC). The North African strain, isolated by Dr. Bill Hansen from leaves of *Oryza* and *Cyperus* species in the Ivory Coast (personal commun-

ication with FGSC), shows higher levels of resistance than the mutant induced in the laboratory (AL-Sagur, A.M 1976 Ph.D. Thesis, Aberdeen University, Scotland). Honduras wild type is resistant to CTAB and BAC but sensitive to DC and polymyxin, similar to the pattern of resistance shown by one of the UV-induced mutants. The Philippine Islands 4 shows a higher level of resistance to DC and BAC but is sensitive to the other surfactants. This is a new pattern of resistance not encountered among the laboratory induced mutants.

TABLE 1

Cross-resistant studies on wild type strains
(++ = very good growth; + = good growth; - = no growth)

Wild type strain	FGSC number	m.t.	Resistance to ($\mu\text{g/ml}$)*			
			DC	CTAB	BAC	Polymyxin B Sulphate 150
<i>N. crassa</i>						
North Africa I	430	A	+ 20	++ 10	++ 12	
Panama CZ30.4	1130	a	++ 5	+ 10	++ 6	
Panama CZ30.6	1131	A	++ 2.5	++ 5	++ 6	
<i>N. intermedia</i>						
Java	431	a	++ 5	++ 5	+ 6	
Fiji	432	a	++ 5	++ 5	++ 6	
Honduras	1300	a		++ 5	++ 6	
Philippine Islands 4	433	a	++ 20	"	+ 15	

*Drugs are incorporated in 2ml slants of Vogel's medium

Exact details of the isolation of the wild types tested here are unknown, and it is impossible to say whether the resistance to surfactant developed following the application of fungicides or as a response to naturally occurring surfactants. Department of Agriculture and Biology, Nuclear Research Center, Tuwaitha, Baghdad, Iraq and University of Aberdeen, Scotland.