Guanine-requiring mutants

O. C. Yoder
Cornell University

Follow this and additional works at: <https://newprairiepress.org/fgr>

This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Yoder, O. C. (1979) "Guanine-requiring mutants," *Fungal Genetics Reports*: Vol. 26, Article 6.
<https://doi.org/10.4148/1941-4765.1694>

This New Mutants and Stocks is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Guanine-requiring mutants

Abstract

Guanine-requiring mutants

Yoder, O.C.

Guanine-requiring mutants.

UV, survivors were concentrated by filtration enrichment in high-sorbose minimal medium, and mutants were selected on minimal medium containing guanine (0.2 mg/ml). Four survivors (out of 191) grew up promptly on guanine and not on minimal or complete medium.

Three of these mutants behaved identically. Although initially clearly negative, they started growing after 3 to 4 days on minimal or complete medium and within 2 additional days achieved growth comparable to that of wild type. Subcultures behaved like the original isolates. The mutants grew more readily on guanosine than on guanine, presumably because guanosine is the more soluble; the threshold level of guanosine required for normal growth was 1 to 2 μ M. Growth on guanosine-medium was inhibited by adenine, adenine or adenylic acid. The inhibition was apparently competitive since there was complete inhibition when adenosine and guanosine were at equimolar concentrations but inhibition was abolished if the guanosine concentration exceeded that of adenine. Cytidine was also inhibitory but at a concentration 5-fold higher than that of guanosine. Other compounds were not inhibitory: thymine, hypoxanthine, xanthine, uracil, thymidine, uridine, cytosine, xanthurenic acid. Adenine (1 mg/ml) prevented adaptation on minimal or complete.

In heterokaryon tests on minimal medium, the three mutants did not complement each other, but all three complemented the fourth mutant. Recombination among the three was < 0.1% and they were therefore assigned to a locus designated gua. Results of five crosses indicated that gua is linked to mating type (probably left) in linkage group 1. Stocks of gua have been deposited with FGSC and assigned Nos. 3524 (A) and 3525 (a).

The remaining mutant, gua(OY304), grew slowly on guanosine-medium and not at all on minimal or complete; it did not adopt. In crosses, it did not act as a female parent but was highly fertile as a male. Progeny (54) of Oak Ridge X gua(OY304) were all fast growing prototrophs. gua(OY304); al-2 formed a fast growing, orange heterokaryon when combined with in on minimal medium. Most colonies derived from plated conidia of the heterokaryon were orange, but a few were white. Ten white and three orange colonies were isolated and transferred to minimal, complete, and minimal plus guanosine. Of the white isolates, 1 was fast-growing gua⁺, 2 were slow-growing gua, 1 was slow-growing gua⁺, and 6 grew at an intermediate rate; 4 of these were gua and 2 were gua⁺. All three orange isolates grew moderately fast; 2 were gua and 1 was gua⁺. The gua(OY304) mutant appears to display quantitative effects and may be cytoplasmic. The guanine mutants, obtained at Stanford University, will not be studied further in this laboratory. - - - Department of Plant Pathology, Cornell University, Ithaca, NY 14853.