

## A new culture method for biochemical studies of the circadian clock

J. Perlman

*University of California at Santa Cruz*

J. F. Feldman

*University of California at Santa Cruz*

Follow this and additional works at: <http://newprairiepress.org/fgr>

---

### Recommended Citation

Perlman, J., and J.F. Feldman (1979) "A new culture method for biochemical studies of the circadian clock," *Fungal Genetics Reports*: Vol. 26, Article 18. <https://doi.org/10.4148/1941-4765.1706>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

# A new culture method for biochemical studies of the circadian clock

## **Abstract**

A new culture method for biochemical studies of the circadian clock

## **Creative Commons License**



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

A new culture method for biochemical  
studier of the circadian clock.

and standing liquid cultures. Methods and results ore presented only for the latter because that system has been characterized the most extensively and potentially may be of the most use.

Cultures were maintained on Horowitz complete slants. All other medium, both liquid and solid, was Vogel's salts containing 1.2% sodium acetate and 0.05% casamino acids [solid medium contained 1.5% agar]. bd A conidia from 6-B day old slants were suspended in distilled water or (in later experiments) liquid medium, and filtered through glass wool. The concentration of conidia in the filtrate was measured (Klett-Summerson colorimeter, blue filter), and an aliquot immediately added to a large volume of stirring liquid medium to give a final concentration of  $2 \times 10^5$  spores/ml. Using an automatic pipetter, 25ml of the stirring suspension were added to each of several dozen 100x15mm plastic disposable Petri dishes. Six growth tuber with solid medium were inoculated at one end with about 50 microliters of undiluted filtrate. All plates and growth tubes were put in constant light at 25°C. After about a day, they were transferred to constant darkness in on environmental growth chamber, also at 25°C. The Petri dish cultures hod visible growth by 24 hours, and subsequently formed a mycelial mat which covered the surface. At several different times after the cultures were placed in the dark, six pieces of mycelium from each of three plates were cut with a cork borer (1.1 mm diam.) and transferred to fresh growth tuber. (The size of the transferred piecer has been varied greatly with identical results.) The growth fronts of control and experimental growth tuber were marked in red safelight (G. E. BCJ, 60 watt) at the same time each subsequent day. After about 7 days in the dark, the timer of occurrence, or phases, of the first conidial bands of the experimental growth tuber were determined by linear regression analysis and compared to the corresponding band of the control tuber.

The phases observed in the growth tubes inoculated with pieces of the standing liquid cultures were very close to those of the controls at all timer sampled. (In some experiments small and consistent phase advances were seen in the experimental tubes.) The sampling manipulations therefore da not affect the phase of the clock, and it may be concluded that the liquid cultures have a normal circadian clock whose phase is ret, like that of the controls, by the light-to-dark transition. The phases of experimental growth tubes start to differ from that of the controls at approximately the time when the liquid cultures reach stationary phase (about 55-60 hours of age). It seems likely that either the clock of older cultures "runs down," or that older cultures are susceptible to phase resetting when cut and transferred. This appears to be largely independent of the length of time they spend in constant light. Preliminary experiments suggest that a 3-hour period in constant light (possibly even less) suffices to set the phase of these cultures.

Experiments are underway to relate the age of the cultures and nuclear division timer to the functioning of the clock. Other types of experiments, particularly the addition of various agents to determine their effects on the clock, ore now possible.

• • • Thimonn Laboratories, University of California at Santa Cruz, Santa Cruz, CA 95064.