

## Sex-hormones in *Neurospora crassa*: hormone action is affected by genetic background

M. S. Islam  
*Atomic Energy Centre*

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Islam, M. S. (1977) "Sex-hormones in *Neurospora crassa*: hormone action is affected by genetic background," *Fungal Genetics Reports*: Vol. 24, Article 4. <https://doi.org/10.4148/1941-4765.1720>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## Sex-hormones in *Neurospora crassa*: hormone action is affected by genetic background

### Abstract

Sex-hormones in *Neurospora crassa*: hormone action is affected by genetic background

Islam, M.S. Sex-hormones in Neurospora crassa: Hormone action is affected by genetic background.

Sex-hormones (sex and fertility-inducing substances) have been isolated from single strain cultures as well as established cross of Neurospora crassa (Em A and Em a) and their biological properties determined (Islam and Weijer 1969 Neurospora Newsletter 15:14; Vigfusson et al 1971 Folia Microbiol. 16: 166; Islam 1973 Mycopath. et Mycol. App. 51: 87). To determine if hormone action is affected by the genetic

background of compatible strains, pair-wise crosses were made with different tester strains (fl<sup>P</sup> a, St. L. a, Em a) to a sterility mutant (7232 A). Inocula were placed at diagonally opposite sides of the petri plates containing 15ml of the crossing medium of Westergaard and Mitchell (1947 Am. J. Botany 34: 573-577), supplemented with a sex-hormone-containing extract from Em A. Five replicate plates were made for both extracted-treated and control (no extract) plates. The number of perithecia per plate was determined at weekly intervals during 4 weeks of incubation at 24-25° C.

The results showed that there was an improvement of fertility (increase in number of perithecia) in extracted-treated plates of crosses fl<sup>P</sup> a x 7232 A and St. L. a x 7232 A as compared with controls. However, in the case of the cross Em a x 7232 A no significant improvement of fertility was noticed. These results indicate that the genetic backgrounds of compatible strains are also a factor for the effective action of Neurospora sex-hormones. - - - Division of Genetics, Atomic Energy Centre, P.O. Box 164, Ramna, Bangladesh.