

## Sex-hormones in *Neurospora crassa*: hormone action is affected by genetic background

M. S. Islam  
*Atomic Energy Centre*

Follow this and additional works at: <http://newprairiepress.org/fgr>

---

### Recommended Citation

Islam, M. S. (1977) "Sex-hormones in *Neurospora crassa*: hormone action is affected by genetic background," *Fungal Genetics Reports*: Vol. 24, Article 4. <https://doi.org/10.4148/1941-4765.1720>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

# Sex-hormones in *Neurospora crassa*: hormone action is affected by genetic background

## **Abstract**

Sex-hormones in *Neurospora crassa*: hormone action is affected by genetic background

## **Creative Commons License**



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Islam, M.S. Sex-hormones in Neurospora crassa: Hormone action is affected by genetic background.

Sex-hormones (sex and fertility-inducing substances) have been isolated from single strain cultures as well as established cross of Neurospora crassa (Em A and Em a) and their biological properties determined (Islam and Weijer 1969 Neurospora Newsletter 15:14; Vigfusson et al 1971 Folia Microbiol. 16: 166; Islam 1973 Mycopath. et Mycol. App. 51: 87). To determine if hormone action is affected by the genetic

background of compatible strains, pair-wise crosses were made with different tester strains (fl<sup>P</sup> a, St. L. a, Em a) to a sterility mutant (7232 A). Inocula were placed at diagonally opposite sides of the petri plates containing 15ml of the crossing medium of Westergaard and Mitchell (1947 Am. J. Botany 34: 573-577), supplemented with a sex-hormone-containing extract from Em A. Five replicate plates were made for both extracted-treated and control (no extract) plates. The number of perithecia per plate was determined at weekly intervals during 4 weeks of incubation at 24-25° C.

The results showed that there was an improvement of fertility (increase in number of perithecia) in extracted-treated plates of crosses fl<sup>P</sup> a x 7232 A and St. L. a x 7232 A as compared with controls. However, in the case of the cross Em a x 7232 A no significant improvement of fertility was noticed. These results indicate that the genetic backgrounds of compatible strains are also a factor for the effective action of Neurospora sex-hormones. - - - Division of Genetics, Atomic Energy Centre, P.O. Box 164, Ramna, Bangladesh.