

## A method for obtaining high specific activity radioactive extracts from perithecia

T. E. Johnson  
*University of Colorado*

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Johnson, T. E. (1978) "A method for obtaining high specific activity radioactive extracts from perithecia," *Fungal Genetics Reports*: Vol. 25, Article 4. <https://doi.org/10.4148/1941-4765.1721>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## A method for obtaining high specific activity radioactive extracts from perithecia

### Abstract

A method for obtaining high specific activity radioactive extracts from perithecia

Johnson, T. E.      A method for obtaining high specific activity radioactive extracts from perithecia.

One way of dissecting a biological process is by isolating mutants which are blocked at different points in that process. These mutants are then used to determine what alterations have occurred in the molecular components of that process.

Many mutants have been isolated which block perithecial development as either the male or female component of a cross (review in Johnson, T.E., (1978) Genetics 88: 27-47). One very general approach to understanding the molecular alterations in these mutants is to determine what changes have occurred in the proteins which are synthesized by those mutants. This report presents a reliable way to label perithecial proteins with radioactive amino acids. The proteins can then be displayed by one or two dimensional electrophoresis and autoradiography.

The desired maternal component is inoculated on crossing medium (Westergaard and Mitchell (1947) Am. J. Botany 34: 573-577) supplemented with 2% sucrose and appropriate growth requirements. Prior to inoculation the petri plates should be covered with filter papers so that the whole surface of the plate is covered (VWR grade no. 613 is especially suitable because the paper withstands long periods of wetting without shredding).

Conditions for growth and fertilization have been given (Johnson (1978) Genetics 88: 27-47). At appropriate time after fertilization protoperithecia and perithecia can be isolated by scraping the surface of the paper with a blunt spatula. The mat is pressed dry, weighed and suspended in crossing medium with no sucrose (5ml per plate harvested) and the desired radioactive label. The culture is then incubated on a rotary shaker for six hours and harvested by filtration through a .6µ Millipore filter. The sample is




Figure 1. Cultures were harvested and incubated, for six hours, total activity taken up i.e., average of three repeats  $\pm$  Standard Deviation,; total radioactivity incorporated into TCA precipitable counts (o-o-o).

then ground (Nasrallah and Srb (1973) Proc. No. Acad. Sci. USA 70: 1891-1893), centrifuged at 5,000 g for 5 minutes and the supernatant frozen at  $-60^{\circ}$  C. Before use, the sample is thawed and centrifuged again at 5,000 g for 5 minutes.

The amount of label incorporated into TCA precipitable material plateaus at about four hours. Labelled proteins can be obtained at any point in development by using this technique; there is, however, a drop off both in uptake and in incorporation at later points in the development.3 program (Figure 1). Using  $^{35}$ S-methionine (2.2 $\mu$  Ci, 700mCi/ $\mu$ g), I have obtained extracts containing as much as 1000 cpm/ $\mu$ g protein. This technique provides a very efficient uptake of label with minimum handling after labeling.

Department of Molecular, Cellular and Developmental Biology, University of Colorado, Boulder, Colorado 80309.