

## Ammonium transport using $^{14}\text{C}$ -methylamine as substrate

J. E. Maruniak  
*Florida State University*

A. G. DeBusk

Follow this and additional works at: <http://newprairiepress.org/fgr>

---

### Recommended Citation

Maruniak, J. E., and A.G. DeBusk (1977) "Ammonium transport using  $^{14}\text{C}$ -methylamine as substrate," *Fungal Genetics Reports*: Vol. 24, Article 9. <https://doi.org/10.4148/1941-4765.1730>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

# Ammonium transport using $^{14}\text{C}$ -methylamine as substrate

## Abstract

Ammonium transport using  $^{14}\text{C}$ -methylamine as substrate

## Creative Commons License



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Marunjak, J. F. and A. G. DeBusk. Ammonium

transport using  $^{14}\text{C}$ -methylamine as substrate.

a  $K_m$  of about  $1.6 \times 10^{-5}$  molar. The determination of Michaelis-Menten constants was done over a range of  $2 \times 10^{-4}$  M to  $5 \times 10^{-7}$  M methylamine. The velocities (Table 1), were determined by initial rates determined from 1, 2, 3 and 4 minutes sample points with conidia that had been reincubated in nitrogen-deficient medium containing 1% glucose for 2 hours prior to the addition of  $^{14}\text{C}$ -methylamine.

Table 1

Methylamine transport rates at various substrate concentrations

Substrate conc. (molar)	$2 \times 10^{-4}$	$1 \times 10^{-4}$	$5 \times 10^{-5}$	$2 \times 10^{-6}$	$1 \times 10^{-6}$	$5 \times 10^{-7}$
Velocity (moles/g/min)	14.89	12.65	10.95	1.66	0.85	0.46

Transport rate greatly increases with nitrogen starvation, with or without glucose present. However, if glucose is absent, there is no saturation of transport even up to 2 mM methylamine. Inhibition kinetics show that  $\text{NH}_4\text{Cl}$  competitively inhibits ( $K_i = 4.7 \times 10^{-4}$  M)  $^{14}\text{C}$ -methylamine uptake, showing that methylamine is a suitable ammonium analog. Transport at pH 5.6 in nitrogen-deficient medium shows that prior development of transport capability is optimal at pH 3 to 4. However, if developed at pH 5.6 in nitrogen deficient medium containing glucose, subsequent transport increases with a pH increase up to pH 9. In addition to protein synthesis (as exemplified by cycloheximide inhibition), energy is required to maintain or produce uptake capability as demonstrated by sodium azide and 2,4-dinitrophenol sensitivity. Furthermore, transport is temperature dependent with a  $Q_{10}$  of 1.9 between 20 and 30°C.


Accumulation assays (in nitrogen-deficient medium containing glucose) using  $^{14}\text{C}$ -methylamine, were performed with Tatum  $\alpha$ , 74-OR8-1a, am1 (FGSC #521), and pmg:mtr:bat (Pm-nbg) (FGSC #2606) (deficient in neutral, basic, and general amino acid transport; Rao and DeBusk 1975 B.B.A. 413:45; 1976 Neurospora Newsl. 22: 12-13) and can be compared in Figure 1. It is interesting to note that the absence of the NADP-linked glutamate dehydrogenase does not prevent methylamine (and presumably ammonium) transport since am1, which lacks this enzyme, is capable of  $^{14}\text{C}$ -methylamine transport. It has been reported (Dubois, Grenson and Wiame 1973 B.B.R.C. 50:967) that glutamate dehydrogenaseless mutants in yeast are derepressed in the presence of ammonium for ammonium repressible activities. The data implies that such phenomena in Neurospora may not be due to lack of transport of ammonium since am1 transports methylamine even better than wild type. Also, pmg:mtr:bat (Pm-nbg) transports  $^{14}\text{C}$ -methylamine almost as well as the parental strain, Tatum  $\alpha$ , which shows the separate identity of amino acid and ammonium transport systems.

A three day growth test at 25°C revealed that methylamine cannot serve as a carbon or nitrogen source for Neurospora. Instead, methylamine, at 10mM, is toxic to Neurospora on a nitrate medium (Vogel's medium N substituting  $\text{KNO}_3$  for  $\text{NH}_4\text{NO}_3$ ) but not on an ammonium medium (Vogel's medium N with  $\text{NH}_4\text{Cl}$  instead of  $\text{NH}_4\text{NO}_3$ ), thus supporting the hypothesis that methylamine and ammonium ions utilize the same transport system.

- - - Genetics Group, Department of Biological Science, Florida State University, Tallahassee, Florida 32306.

Neurospora crassa wild type, Tatum  $\alpha$  (5Y4fg  $\alpha$ ) conidia, possess an active transport system for ammonium which can be characterized using  $^{14}\text{C}$ -methylamine as an ammonium analog. In nitrogen-deficient medium (Vogel's minimal medium with  $\text{NH}_4\text{NO}_3$  omitted but containing 1% glucose), the initial rate of uptake of  $^{14}\text{C}$ -methylamine shows Michaelis-Menten saturation kinetics with a  $V_{max}$  of about 15 moles/g/min and

Figure 1



Conidial transport of  $^{14}\text{C}$ -methylamine at 25°C by several strains of Neurospora crassa in nitrogen-deficient medium containing 1% glucose. The isotope concentration was  $1 \times 10^{-4}$  M at a specific activity of 0.1  $\mu\text{Ci}/0.1$  mole/ml.